

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
27 December 2001 (27.12.2001)

PCT

(10) International Publication Number
WO 01/98468 A2

(51) International Patent Classification⁷: C12N 9/00

(21) International Application Number: PCT/US01/19178

(22) International Filing Date: 13 June 2001 (13.06.2001)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:

60/212,336	16 June 2000 (16.06.2000)	US
60/213,955	22 June 2000 (22.06.2000)	US
60/215,396	29 June 2000 (29.06.2000)	US
60/216,821	7 July 2000 (07.07.2000)	US
60/218,946	14 July 2000 (14.07.2000)	US

233 Coy Drive, San Jose, CA 95123 (US). PATTERSON, Chandra [US/US]; 490 Sherwood Way #1, Menlo Park, CA 94025 (US). TANG, Y., Tom [US/US]; 4230 Ranwick Court, San Jose, CA 95118 (US). WALSH, Roderick, T. [IE/US]; 8 Boundary Court, St. Lawrence Road, Canterbury, Kent CT1 3EZ (GB). AZIMZAI, Yalda [US/US]; 5518 Boulder Canyon Drive, Castro Valley, CA 94552 (US). LU, Yan [CN/US]; 3885 Corrina Way, Palo Alto, CA 94303 (US). RAMKUMAR, Jayalaxmi [IN/US]; 34359 Maybird Circle, Fremont, CA 94555 (US). XU, Yuming [US/US]; 1739 Walnut Drive, Mountain View, CA 94040 (US). REDDY, Roopa [IN/US]; 1233 W. McKinley Avenue #3, Sunnyvale, CA 94086 (US). DAS, Debopriya [IN/US]; 1179 Bonita Avenue, Apt. 3, Mountain View, CA 94040 (US). KEARNEY, Liam [IE/US]; 50 Woodside Avenue, San Francisco, CA 94127 (US). KALLICK, Deborah, A. [US/US]; 900 Olive Street, Menlo Park, CA 94025 (US).

(71) Applicant (for all designated States except US): INCYTE GENOMICS, INC. [US/US]; 3160 Porter Drive, Palo Alto, CA 94304 (US).

(74) Agents: HAMLET-COX, Diana et al.; Incyte Genomics, Inc., 3160 Porter Drive, Palo Alto, CA 94304 (US).

(72) Inventors; and

(75) Inventors/Applicants (for US only): YUE, Henry [US/US]; 826 Lois Avenue, Sunnyvale, CA 94087 (US). ELLIOTT, Vicki, S. [US/US]; 3770 Polton Place Way, San Jose, CA 95121 (US). GANDHI, Ameena, R. [US/US]; 837 Roble Avenue, #1, Menlo Park, CA 94025 (US). LAL, Preeti [IN/US]; P.O. Box 5142, Santa Clara, CA 95056 (US). AU-YOUNG, Janice [US/US]; 233 Golden Eagle Lane, Brisbane, CA 94005 (US). TRIBOULEY, Catherine, M. [FR/US]; 1121 Tennessee Street, #5, San Francisco, CA 94107 (US). DELEGEANE, Angelo, M. [US/US]; 594 Angus Drive, Milpitas, CA 95035 (US). BAUGHN, Mariah, R. [US/US]; 14244 Santiago Road, San Leandro, CA 94577 (US). NGUYEN, Danniell, B. [US/US]; 1403 Ridgewood Drive, San Jose, CA 95118 (US). LEE, Ernestine, A. [US/US]; 624 Kains Street, Albany, CA 94706 (US). HAFALIA, April [US/US]; 2227 Calle de Primavera, Santa Clara, CA 95054 (US). KHAN, Farrah, A. [IN/US]; 3617 Central Road #102, Glenview, IL 60025 (US). WALIA, Narinder, K. [US/US]; 890 Davis Street #205, San Leandro, CA 94577 (US). YAO, Monique, G. [US/US]; 111 Frederick Court, Mountain View, CA 94043 (US). LU, Dyung, Aina, M. [US/US];

(81) Designated States (national): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.

(84) Designated States (regional): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).


Published:

— without international search report and to be republished upon receipt of that report

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: PROTEASES

(57) Abstract: The invention provides human proteases (PRTS) and polynucleotides which identify and encode PRTS. The invention also provides expression vectors, host cells, antibodies, agonists, and antagonists. The invention also provides methods for diagnosing, treating, or preventing disorders associated with aberrant expression of PRTS.



WO 01/98468 A2

PROTEASES

TECHNICAL FIELD

This invention relates to nucleic acid and amino acid sequences of proteases and to the use of these sequences in hydrolysis of peptide bonds and in the diagnosis, treatment, and prevention of gastrointestinal, cardiovascular, autoimmune/inflammatory, cell proliferative, developmental, epithelial, neurological, and reproductive disorders, and in the assessment of the effects of exogenous compounds on the expression of nucleic acid and amino acid sequences of proteases.

BACKGROUND OF THE INVENTION

Proteases cleave proteins and peptides at the peptide bond that forms the backbone of the protein or peptide chain. Proteolysis is one of the most important and frequent enzymatic reactions that occurs both within and outside of cells. Proteolysis is responsible for the activation and maturation of nascent polypeptides, the degradation of misfolded and damaged proteins, and the controlled turnover of peptides within the cell. Proteases participate in digestion, endocrine function, and tissue remodeling during embryonic development, wound healing, and normal growth. Proteases can play a role in regulatory processes by affecting the half life of regulatory proteins. Proteases are involved in the etiology or progression of disease states such as inflammation, angiogenesis, tumor dispersion and metastasis, cardiovascular disease, neurological disease, and bacterial, parasitic, and viral infections.

Proteases can be categorized on the basis of where they cleave their substrates. Exopeptidases, which include aminopeptidases, dipeptidyl peptidases, tripeptidases, carboxypeptidases, peptidyl-dipeptidases, dipeptidases, and omega peptidases, cleave residues at the termini of their substrates. Endopeptidases, including serine proteases, cysteine proteases, and metalloproteases, cleave at residues within the peptide. Four principal categories of mammalian proteases have been identified based on active site structure, mechanism of action, and overall three-dimensional structure. (See Beynon, R.J. and J.S. Bond (1994) Proteolytic Enzymes: A Practical Approach, Oxford University Press, New York NY, pp. 1-5.)

Serine Proteases

The serine proteases (SPs) are a large, widespread family of proteolytic enzymes that include the digestive enzymes trypsin and chymotrypsin, components of the complement and blood-clotting cascades, and enzymes that control the degradation and turnover of macromolecules within the cell and in the extracellular matrix. Most of the more than 20 subfamilies can be grouped into six clans, each with a common ancestor. These six clans are hypothesized to have descended from at least four evolutionarily distinct ancestors. SPs are named for the presence of a serine residue found in the active

catalytic site of most families. The active site is defined by the catalytic triad, a set of conserved asparagine, histidine, and serine residues critical for catalysis. These residues form a charge relay network that facilitates substrate binding. Other residues outside the active site form an oxyanion hole that stabilizes the tetrahedral transition intermediate formed during catalysis. SPs have a wide range of substrates and can be subdivided into subfamilies on the basis of their substrate specificity. The main subfamilies are named for the residue(s) after which they cleave: trypases (after arginine or lysine), aspases (after aspartate), chymases (after phenylalanine or leucine), metases (methionine), and serases (after serine) (Rawlings, N.D. and A.J. Barrett (1994) *Methods Enzymol.* 244:19-61).

Most mammalian serine proteases are synthesized as zymogens, inactive precursors that are activated by proteolysis. For example, trypsinogen is converted to its active form, trypsin, by enteropeptidase. Enteropeptidase is an intestinal protease that removes an N-terminal fragment from trypsinogen. The remaining active fragment is trypsin, which in turn activates the precursors of the other pancreatic enzymes. Likewise, proteolysis of prothrombin, the precursor of thrombin, generates three separate polypeptide fragments. The N-terminal fragment is released while the other two fragments, which comprise active thrombin, remain associated through disulfide bonds.

The two largest SP subfamilies are the chymotrypsin (S1) and subtilisin (S8) families. Some members of the chymotrypsin family contain two structural domains unique to this family. Kringle domains are triple-looped, disulfide cross-linked domains found in varying copy number. Kringles are thought to play a role in binding mediators such as membranes, other proteins or phospholipids, and in the regulation of proteolytic activity (PROSITE PDOC00020). Apple domains are 90 amino-acid repeated domains, each containing six conserved cysteines. Three disulfide bonds link the first and sixth, second and fifth, and third and fourth cysteines (PROSITE PDOC00376). Apple domains are involved in protein-protein interactions. S1 family members include trypsin, chymotrypsin, coagulation factors IX-XII, complement factors B, C, and D, granzymes, kallikrein, and tissue- and urokinase-plasminogen activators. The subtilisin family has members found in the eubacteria, archaeobacteria, eukaryotes, and viruses. Subtilisins include the proprotein-processing endopeptidases kexin and furin and the pituitary prohormone convertases PC1, PC2, PC3, PC6, and PACE4 (Rawlings and Barrett, *supra*). The prolyl oligopeptidase (S9) family includes enzymes from prokaryotes and eukaryotes with greatly differing specificities. Dipeptidyl peptidase IV (DPP-IV) is identical to CD26 and is implicated in the inactivation of peptide hormones, as well as in regulating T-cell growth (reviewed in Kahne, T. et al. (1999) *Int. J. Mol. Med.* 4:3-15; Mentlein, R. (1999) *Regul. Pept.* 85:9-24): Inhibition of DPP-IV has been suggested as a treatment for type 2 diabetes (Holst, J.J. and C.F. Deacon (1998) *Diabetes* 47:1663-1670), and lowered serum DPP-IV activity has been measured in anorexia and bulimia patients (van West, D. et al. (2000) *Eur. Arch. Psych. Clin. Neurosci.* 250:86-92).

SPs have functions in many normal processes and some have been implicated in the etiology or treatment of disease. Enterokinase, the initiator of intestinal digestion, is found in the intestinal brush border, where it cleaves the acidic propeptide from trypsinogen to yield active trypsin (Kitamoto, Y. et al. (1994) *Proc. Natl. Acad. Sci. USA* 91:7588-7592). Prolylcarboxypeptidase, a lysosomal serine
5 peptidase that cleaves peptides such as angiotensin II and III and [des-Arg⁹] bradykinin, shares sequence homology with members of both the serine carboxypeptidase and prolylendopeptidase families (Tan, F. et al. (1993) *J. Biol. Chem.* 268:16631-16638). The protease neuropsin may influence synapse formation and neuronal connectivity in the hippocampus in response to neural signaling (Chen, Z.-L. et al. (1995) *J. Neurosci.* 15:5088-5097). Tissue plasminogen activator is useful for acute
10 management of stroke (Zivin, J.A. (1999) *Neurology* 53:14-19) and myocardial infarction (Ross, A.M. (1999) *Clin. Cardiol.* 22:165-171). Some receptors (PAR, for proteinase-activated receptor), highly expressed throughout the digestive tract, are activated by proteolytic cleavage of an extracellular domain. The major agonists for PARs, thrombin, trypsin, and mast cell tryptase, are released in allergy and inflammatory conditions. Control of PAR activation by proteases has been suggested as a
15 promising therapeutic target (Vergnolle, N. (2000) *Aliment. Pharmacol. Ther.* 14:257-266; Rice, K.D. et al. (1998) *Curr. Pharm. Des.* 4:381-396). Prostate-specific antigen (PSA) is a kallikrein-like serine protease synthesized and secreted exclusively by epithelial cells in the prostate gland. Serum PSA is elevated in prostate cancer and is the most sensitive physiological marker for monitoring cancer progression and response to therapy. PSA can also identify the prostate as the origin of a metastatic
20 tumor (Brawer, M.K. and P.H. Lange (1989) *Urology* 33:11-16).

The signal peptidase is a specialized class of SP found in all prokaryotic and eukaryotic cell types that serves in the processing of signal peptides from certain proteins. Signal peptides are amino-terminal domains of a protein which direct the protein from its ribosomal assembly site to a particular cellular or extracellular location. Once the protein has been exported, removal of the signal
25 sequence by a signal peptidase and posttranslational processing, e.g., glycosylation or phosphorylation, activate the protein. Signal peptidases exist as multi-subunit complexes in both yeast and mammals. The canine signal peptidase complex is composed of five subunits, all associated with the microsomal membrane and containing hydrophobic regions that span the membrane one or more times (Shelness, G.S. and G. Blobel (1990) *J. Biol. Chem.* 265:9512-9519). Some of these subunits serve to fix the
30 complex in its proper position on the membrane while others contain the actual catalytic activity.

Another family of proteases which have a serine in their active site are dependent on the hydrolysis of ATP for their activity. These proteases contain proteolytic core domains and regulatory ATPase domains which can be identified by the presence of the P-loop, an ATP/GTP-binding motif (PROSITE PDOC00803). Members of this family include the eukaryotic mitochondrial matrix

proteases, Clp protease and the proteasome. Clp protease was originally found in plant chloroplasts but is believed to be widespread in both prokaryotic and eukaryotic cells. The gene for early-onset torsion dystonia encodes a protein related to Clp protease (Ozelius, L.J. et al. (1998) *Adv. Neurol.* 78:93-105).

The proteasome is an intracellular protease complex found in some bacteria and in all eukaryotic cells, and plays an important role in cellular physiology. Proteasomes are associated with the ubiquitin conjugation system (UCS), a major pathway for the degradation of cellular proteins of all types, including proteins that function to activate or repress cellular processes such as transcription and cell cycle progression (Ciechanover, A. (1994) *Cell* 79:13-21). In the UCS pathway, proteins targeted for degradation are conjugated to ubiquitin, a small heat stable protein. The ubiquitinated protein is then recognized and degraded by the proteasome. The resultant ubiquitin-peptide complex is hydrolyzed by a ubiquitin carboxyl terminal hydrolase, and free ubiquitin is released for reutilization by the UCS. Ubiquitin-proteasome systems are implicated in the degradation of mitotic cyclic kinases, oncoproteins, tumor suppressor genes (p53), cell surface receptors associated with signal transduction, transcriptional regulators, and mutated or damaged proteins (Ciechanover, *supra*). This pathway has been implicated in a number of diseases, including cystic fibrosis, Angelman's syndrome, and Liddle syndrome (reviewed in Schwartz, A.L. and A. Ciechanover (1999) *Annu. Rev. Med.* 50:57-74). A murine proto-oncogene, *Unp*, encodes a nuclear ubiquitin protease whose overexpression leads to oncogenic transformation of NIH3T3 cells. The human homologue of this gene is consistently elevated in small cell tumors and adenocarcinomas of the lung (Gray, D.A. (1995) *Oncogene* 10:2179-2183). Ubiquitin carboxyl terminal hydrolase is involved in the differentiation of a lymphoblastic leukemia cell line to a non-dividing mature state (Maki, A. et al. (1996) *Differentiation* 60:59-66). In neurons, ubiquitin carboxyl terminal hydrolase (PGP 9.5) expression is strong in the abnormal structures that occur in human neurodegenerative diseases (Lowe, J. et al. (1990) *J. Pathol.* 161:153-160). The proteasome is a large (~2000 kDa) multisubunit complex composed of a central catalytic core containing a variety of proteases arranged in four seven-membered rings with the active sites facing inwards into the central cavity, and terminal ATPase subunits covering the outer port of the cavity and regulating substrate entry (for review, see Schmidt, M. et al. (1999) *Curr. Opin. Chem. Biol.* 3:584-591).

Cysteine Proteases

Cysteine proteases (CPs) are involved in diverse cellular processes ranging from the processing of precursor proteins to intracellular degradation. Nearly half of the CPs known are present only in viruses. CPs have a cysteine as the major catalytic residue at the active site where catalysis proceeds via a thioester intermediate and is facilitated by nearby histidine and asparagine residues. A glutamine residue is also important, as it helps to form an oxyanion hole. Two important CP families include the

papain-like enzymes (C1) and the calpains (C2). Papain-like family members are generally lysosomal or secreted and therefore are synthesized with signal peptides as well as propeptides. Most members bear a conserved motif in the propeptide that may have structural significance (Karrer, K.M. et al. (1993) Proc. Natl. Acad. Sci. USA 90:3063-3067). Three-dimensional structures of papain family members show a bilobed molecule with the catalytic site located between the two lobes. Papains include cathepsins B, C, H, L, and S, certain plant allergens and dipeptidyl peptidase (for a review, see Rawlings, N.D. and A.J. Barrett (1994) Methods Enzymol. 244:461-486).

Some CPs are expressed ubiquitously, while others are produced only by cells of the immune system. Of particular note, CPs are produced by monocytes, macrophages and other cells which migrate to sites of inflammation and secrete molecules involved in tissue repair. Overabundance of these repair molecules plays a role in certain disorders. In autoimmune diseases such as rheumatoid arthritis, secretion of the cysteine peptidase cathepsin C degrades collagen, laminin, elastin and other structural proteins found in the extracellular matrix of bones. Bone weakened by such degradation is also more susceptible to tumor invasion and metastasis. Cathepsin L expression may also contribute to the influx of mononuclear cells which exacerbates the destruction of the rheumatoid synovium (Keyszer, G.M. (1995) Arthritis Rheum. 38:976-984).

Calpains are calcium-dependent cytosolic endopeptidases which contain both an N-terminal catalytic domain and a C-terminal calcium-binding domain. Calpain is expressed as a proenzyme heterodimer consisting of a catalytic subunit unique to each isoform and a regulatory subunit common to different isoforms. Each subunit bears a calcium-binding EF-hand domain. The regulatory subunit also contains a hydrophobic glycine-rich domain that allows the enzyme to associate with cell membranes. Calpains are activated by increased intracellular calcium concentration, which induces a change in conformation and limited autolysis. The resultant active molecule requires a lower calcium concentration for its activity (Chan, S.L. and M.P. Mattson (1999) J. Neurosci. Res. 58:167-190). Calpain expression is predominantly neuronal, although it is present in other tissues. Several chronic neurodegenerative disorders, including ALS, Parkinson's disease and Alzheimer's disease are associated with increased calpain expression (Chan and Mattson, *supra*). Calpain-mediated breakdown of the cytoskeleton has been proposed to contribute to brain damage resulting from head injury (McCracken, E. et al. (1999) J. Neurotrauma 16:749-761). Calpain-3 is predominantly expressed in skeletal muscle, and is responsible for limb-girdle muscular dystrophy type 2A (Minami, N. et al. (1999) J. Neurol. Sci. 171:31-37).

Another family of thiol proteases is the caspases, which are involved in the initiation and execution phases of apoptosis. A pro-apoptotic signal can activate initiator caspases that trigger a proteolytic caspase cascade, leading to the hydrolysis of target proteins and the classic apoptotic death

of the cell. Two active site residues, a cysteine and a histidine, have been implicated in the catalytic mechanism. Caspases are among the most specific endopeptidases, cleaving after aspartate residues. Caspases are synthesized as inactive zymogens consisting of one large (p20) and one small (p10) subunit separated by a small spacer region, and a variable N-terminal prodomain. This prodomain
5 interacts with cofactors that can positively or negatively affect apoptosis. An activating signal causes autoproteolytic cleavage of a specific aspartate residue (D297 in the caspase-1 numbering convention) and removal of the spacer and prodomain, leaving a p10/p20 heterodimer. Two of these heterodimers interact via their small subunits to form the catalytically active tetramer. The long prodomains of some caspase family members have been shown to promote dimerization and auto-processing of procaspases.
10 Some caspases contain a "death effector domain" in their prodomain by which they can be recruited into self-activating complexes with other caspases and FADD protein associated death receptors or the TNF receptor complex. In addition, two dimers from different caspase family members can associate, changing the substrate specificity of the resultant tetramer. Endogenous caspase inhibitors (inhibitor of apoptosis proteins, or IAPs) also exist. All these interactions have clear effects on the control of
15 apoptosis (reviewed in Chan and Mattson, supra; Salveson, G.S. and V.M. Dixit (1999) Proc. Natl. Acad. Sci. USA 96:10964-10967).

Caspases have been implicated in a number of diseases. Mice lacking some caspases have severe nervous system defects due to failed apoptosis in the neuroepithelium and suffer early lethality. Others show severe defects in the inflammatory response, as caspases are responsible for processing IL-
20 1b and possibly other inflammatory cytokines (Chan and Mattson, supra). Cowpox virus and baculoviruses target caspases to avoid the death of their host cell and promote successful infection. In addition, increases in inappropriate apoptosis have been reported in AIDS, neurodegenerative diseases and ischemic injury, while a decrease in cell death is associated with cancer (Salveson and Dixit, supra; Thompson, C.B. (1995) Science 267:1456-1462).

25 Aspartyl proteases

Aspartyl proteases (APs) include the lysosomal proteases cathepsins D and E, as well as chymosin, renin, and the gastric pepsins. Most retroviruses encode an AP, usually as part of the pol polyprotein. APs, also called acid proteases, are monomeric enzymes consisting of two domains, each domain containing one half of the active site with its own catalytic aspartic acid residue. APs are most
30 active in the range of pH 2-3, at which one of the aspartate residues is ionized and the other neutral. The pepsin family of APs contains many secreted enzymes, and all are likely to be synthesized with signal peptides and propeptides. Most family members have three disulfide loops, the first ~5 residue loop following the first aspartate, the second 5-6 residue loop preceding the second aspartate, and the third and largest loop occurring toward the C terminus. Retropepsins, on the other hand, are analogous

to a single domain of pepsin, and become active as homodimers with each retropepsin monomer contributing one half of the active site. Retropepsins are required for processing the viral polyproteins.

APs have roles in various tissues, and some have been associated with disease. Renin mediates the first step in processing the hormone angiotensin, which is responsible for regulating electrolyte

- 5 balance and blood pressure (reviewed in Crews, D.E. and S.R. Williams (1999) Hum. Biol. 71:475-503). Abnormal regulation and expression of cathepsins are evident in various inflammatory disease states. Expression of cathepsin D is elevated in synovial tissues from patients with rheumatoid arthritis and osteoarthritis. The increased expression and differential regulation of the cathepsins are linked to the metastatic potential of a variety of cancers (Chambers, A.F. et al. (1993) Crit. Rev. Oncol. 4:95-114).

Metalloproteases

Metalloproteases require a metal ion for activity, usually manganese or zinc. Examples of manganese metalloenzymes include aminopeptidase P and human proline dipeptidase (PEPD).

- Aminopeptidase P can degrade bradykinin, a nonapeptide activated in a variety of inflammatory responses. Aminopeptidase P has been implicated in coronary ischemia/reperfusion injury. Administration of aminopeptidase P inhibitors has been shown to have a cardioprotective effect in rats (Ersahin, C. et al (1999) J. Cardiovasc. Pharmacol. 34:604-611).

- Most zinc-dependent metalloproteases share a common sequence in the zinc-binding domain. The active site is made up of two histidines which act as zinc ligands and a catalytic glutamic acid C-terminal to the first histidine. Proteins containing this signature sequence are known as the metzincins and include aminopeptidases B and N, angiotensin-converting enzyme, neurolysin, the matrix metalloproteases and the adamalysins (ADAMS). An alternate sequence is found in the zinc carboxypeptidases, in which all three conserved residues – two histidines and a glutamic acid – are involved in zinc binding.

- 25 A number of the neutral metalloendopeptidases, including angiotensin converting enzyme and the aminopeptidases, are involved in the metabolism of peptide hormones. High aminopeptidase B activity, for example, is found in the adrenal glands and neurohypophyses of hypertensive rats (Prieto, I. et al. (1998) Horm. Metab. Res. 30:246-248). Oligopeptidase M/neurolysin can hydrolyze bradykinin as well as neurotensin (Serizawa, A. et al. (1995) J. Biol. Chem 270:2092-2098).
- 30 Neurotensin is a vasoactive peptide that can act as a neurotransmitter in the brain, where it has been implicated in limiting food intake (Tritos, N.A. et al. (1999) Neuropeptides 33:339-349).

The matrix metalloproteases (MMPs) are a family of at least 23 enzymes that can degrade components of the extracellular matrix (ECM). They are Zn^{+2} endopeptidases with an N-terminal catalytic domain. Nearly all members of the family have a hinge peptide and C-terminal domain which

can bind to substrate molecules in the ECM or to inhibitors produced by the tissue (TIMPs, for tissue inhibitor of metalloprotease; Campbell, I.L. et al. (1999) Trends Neurosci. 22:285). The presence of fibronectin-like repeats, transmembrane domains, or C-terminal hemopexinase-like domains can be used to separate MMPs into collagenase, gelatinase, stromelysin and membrane-type MMP subfamilies. In the inactive form, the Zn^{+2} ion in the active site interacts with a cysteine in the pro-sequence. Activating factors disrupt the Zn^{+2} -cysteine interaction, or "cysteine switch," exposing the active site. This partially activates the enzyme, which then cleaves off its propeptide and becomes fully active. MMPs are often activated by the serine proteases plasmin and furin. MMPs are often regulated by stoichiometric, noncovalent interactions with inhibitors; the balance of protease to inhibitor, then, is very important in tissue homeostasis (reviewed in Yong, V.W. et al. (1998) Trends Neurosci. 21:75). Ehlers-Danlos syndrome type VII C is caused by mutations in the procollagen I N-proteinase gene (Colige, A. et al. (1999) Am. J. Hum. Genet. 65:308-317).

MMPs are implicated in a number of diseases including osteoarthritis (Mitchell, P. et al. (1996) J. Clin. Invest. 97:761), atherosclerotic plaque rupture (Sukhova, G.K. et al. (1999) Circulation 99:2503), aortic aneurysm (Schneiderman, J. et al. (1998) Am. J. Path. 152:703), non-healing wounds (Saarialho-Kere, U.K. et al. (1994) J. Clin. Invest. 94:79), bone resorption (Blavier, L. and J.M. Delaisse (1995) J. Cell Sci. 108:3649), age-related macular degeneration (Steen, B. et al. (1998) Invest. Ophthalmol. Vis. Sci. 39:2194), emphysema (Finlay, G.A. et al. (1997) Thorax 52:502), myocardial infarction (Rohde, L.E. et al. (1999) Circulation 99:3063) and dilated cardiomyopathy (Thomas, C.V. et al. (1998) Circulation 97:1708). MMP inhibitors prevent metastasis of mammary carcinoma and experimental tumors in rat, and Lewis lung carcinoma, hemangioma, and human ovarian carcinoma xenografts in mice (Eccles, S.A. et al. (1996) Cancer Res. 56:2815; Anderson et al. (1996) Cancer Res. 56:715-718; Volpert, O.V. et al. (1996) J. Clin. Invest. 98:671; Taraboletti, G. et al. (1995) J. NCI 87:293; Davies, B. et al. (1993) Cancer Res. 53:2087). MMPs may be active in Alzheimer's disease. A number of MMPs are implicated in multiple sclerosis, and administration of MMP inhibitors can relieve some of its symptoms (reviewed in Yong, *supra*).

Another family of metalloproteases is the ADAMs, for A Disintegrin and Metalloprotease Domain, which they share with their close relatives the adamalysins, snake venom metalloproteases (SVMPs). ADAMs combine features of both cell surface adhesion molecules and proteases, containing a prodomain, a protease domain, a disintegrin domain, a cysteine rich domain, an epidermal growth factor repeat, a transmembrane domain, and a cytoplasmic tail. The first three domains listed above are also found in the SVMPs. The ADAMs possess four potential functions: proteolysis, adhesion, signaling and fusion. The ADAMs share the metzincin zinc binding sequence and are inhibited by some MMP antagonists such as TIMP-1.

ADAMs are implicated in such processes as sperm-egg binding and fusion, myoblast fusion, and protein-ectodomain processing or shedding of cytokines, cytokine receptors, adhesion proteins and other extracellular protein domains (Schl ndorff, J. and C.P. Blobel (1999) J. Cell. Sci. 112:3603-3617). The Kuzbanian protein cleaves a substrate in the NOTCH pathway (possibly NOTCH itself),
5 activating the program for lateral inhibition in Drosophila neural development. Two ADAMs, TACE (ADAM 17) and ADAM 10, are proposed to have analogous roles in the processing of amyloid precursor protein in the brain (Schl ndorff and Blobel, supra). TACE has also been identified as the TNF activating enzyme (Black, R.A. et al. (1997) Nature 385:729). TNF is a pleiotropic cytokine that is important in mobilizing host defenses in response to infection or trauma, but can cause severe
10 damage in excess and is often overproduced in autoimmune disease. TACE cleaves membrane-bound pro-TNF to release a soluble form. Other ADAMs may be involved in a similar type of processing of other membrane-bound molecules.

The ADAMTS sub-family has all of the features of ADAM family metalloproteases and contain an additional thrombospondin domain (TS). The prototypic ADAMTS was identified in mouse,
15 found to be expressed in heart and kidney and upregulated by proinflammatory stimuli (Kuno, K. et al. (1997) J. Biol. Chem. 272:556-562). To date eleven members are recognized by the Human Genome Organization (HUGO; <http://www.gene.ucl.ac.uk/users/hester/adamts.html#Approved>). Members of this family have the ability to degrade aggrecan, a high molecular weight proteoglycan which provides cartilage with important mechanical properties including compressibility, and which is lost during the
20 development of arthritis. Enzymes which degrade aggrecan are thus considered attractive targets to prevent and slow the degradation of articular cartilage (See, e.g., Tortorella, M.D. (1999) Science 284:1664; Abbaszade, I. (1999) J. Biol. Chem. 274:23443). Other members are reported to have antiangiogenic potential (Kuno et al., supra) and/or procollagen processing (Colige, A. et al. (1997) Proc. Natl. Acad. Sci. USA 94:2374).

25 The discovery of new proteases and the polynucleotides encoding them satisfies a need in the art by providing new compositions which are useful in hydrolysis of peptide bonds and in the diagnosis, prevention, and treatment of gastrointestinal, cardiovascular, autoimmune/inflammatory, cell proliferative, developmental, epithelial, neurological, and reproductive disorders, and in the assessment of the effects of exogenous compounds on the expression of nucleic acid and amino acid sequences of
30 proteases.

SUMMARY OF THE INVENTION

The invention features purified polypeptides, proteases, referred to collectively as "PRTS" and individually as "PRTS-1," "PRTS-2," "PRTS-3," "PRTS-4," "PRTS-5," "PRTS-6," "PRTS-7,"

"PRTS-8," "PRTS-9," "PRTS-10," "PRTS-11," "PRTS-12," "PRTS-13," "PRTS-14," "PRTS-15," "PRTS-16," "PRTS-17," "PRTS-18," "PRTS-19," "PRTS-20," and "PRTS-21." In one aspect, the invention provides an isolated polypeptide selected from the group consisting of a) a polypeptide comprising an amino acid sequence selected from the group consisting of SEQ ID NO:1-21, b) a

5 polypeptide comprising a naturally occurring amino acid sequence at least 90% identical to an amino acid sequence selected from the group consisting of SEQ ID NO:1-21, c) a biologically active fragment of a polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO:1-21, and d) an immunogenic fragment of a polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO:1-21. In one alternative, the invention provides an isolated polypeptide

10 comprising the amino acid sequence of SEQ ID NO:1-21.

The invention further provides an isolated polynucleotide encoding a polypeptide selected from the group consisting of a) a polypeptide comprising an amino acid sequence selected from the group consisting of SEQ ID NO:1-21, b) a polypeptide comprising a naturally occurring amino acid sequence at least 90% identical to an amino acid sequence selected from the group consisting of SEQ ID NO:1-

15 21, c) a biologically active fragment of a polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO:1-21, and d) an immunogenic fragment of a polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO:1-21. In one alternative, the polynucleotide encodes a polypeptide selected from the group consisting of SEQ ID NO:1-21. In another alternative, the polynucleotide is selected from the group consisting of SEQ ID NO:22-42.

20 Additionally, the invention provides a recombinant polynucleotide comprising a promoter sequence operably linked to a polynucleotide encoding a polypeptide selected from the group consisting of a) a polypeptide comprising an amino acid sequence selected from the group consisting of SEQ ID NO:1-21, b) a polypeptide comprising a naturally occurring amino acid sequence at least 90% identical to an amino acid sequence selected from the group consisting of SEQ ID NO:1-21, c) a biologically

25 active fragment of a polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO:1-21, and d) an immunogenic fragment of a polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO:1-21. In one alternative, the invention provides a cell transformed with the recombinant polynucleotide. In another alternative, the invention provides a transgenic organism comprising the recombinant polynucleotide.

30 The invention also provides a method for producing a polypeptide selected from the group consisting of a) a polypeptide comprising an amino acid sequence selected from the group consisting of SEQ ID NO:1-21, b) a polypeptide comprising a naturally occurring amino acid sequence at least 90% identical to an amino acid sequence selected from the group consisting of SEQ ID NO:1-21, c) a biologically active fragment of a polypeptide having an amino acid sequence selected from the group

consisting of SEQ ID NO:1-21, and d) an immunogenic fragment of a polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO:1-21. The method comprises a) culturing a cell under conditions suitable for expression of the polypeptide, wherein said cell is transformed with a recombinant polynucleotide comprising a promoter sequence operably linked to a polynucleotide encoding the polypeptide, and b) recovering the polypeptide so expressed.

Additionally, the invention provides an isolated antibody which specifically binds to a polypeptide selected from the group consisting of a) a polypeptide comprising an amino acid sequence selected from the group consisting of SEQ ID NO:1-21, b) a polypeptide comprising a naturally occurring amino acid sequence at least 90% identical to an amino acid sequence selected from the group consisting of SEQ ID NO:1-21, c) a biologically active fragment of a polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO:1-21, and d) an immunogenic fragment of a polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO:1-21.

The invention further provides an isolated polynucleotide selected from the group consisting of a) a polynucleotide comprising a polynucleotide sequence selected from the group consisting of SEQ ID NO:22-42, b) a polynucleotide comprising a naturally occurring polynucleotide sequence at least 90% identical to a polynucleotide sequence selected from the group consisting of SEQ ID NO:22-42, c) a polynucleotide complementary to the polynucleotide of a), d) a polynucleotide complementary to the polynucleotide of b), and e) an RNA equivalent of a)-d). In one alternative, the polynucleotide comprises at least 60 contiguous nucleotides.

Additionally, the invention provides a method for detecting a target polynucleotide in a sample, said target polynucleotide having a sequence of a polynucleotide selected from the group consisting of a) a polynucleotide comprising a polynucleotide sequence selected from the group consisting of SEQ ID NO:22-42, b) a polynucleotide comprising a naturally occurring polynucleotide sequence at least 90% identical to a polynucleotide sequence selected from the group consisting of SEQ ID NO:22-42, c) a polynucleotide complementary to the polynucleotide of a), d) a polynucleotide complementary to the polynucleotide of b), and e) an RNA equivalent of a)-d). The method comprises a) hybridizing the sample with a probe comprising at least 20 contiguous nucleotides comprising a sequence complementary to said target polynucleotide in the sample, and which probe specifically hybridizes to said target polynucleotide, under conditions whereby a hybridization complex is formed between said probe and said target polynucleotide or fragments thereof, and b) detecting the presence or absence of said hybridization complex, and optionally, if present, the amount thereof. In one alternative, the probe comprises at least 60 contiguous nucleotides.

The invention further provides a method for detecting a target polynucleotide in a sample, said target polynucleotide having a sequence of a polynucleotide selected from the group consisting of a) a

polynucleotide comprising a polynucleotide sequence selected from the group consisting of SEQ ID NO:22-42, b) a polynucleotide comprising a naturally occurring polynucleotide sequence at least 90% identical to a polynucleotide sequence selected from the group consisting of SEQ ID NO:22-42, c) a polynucleotide complementary to the polynucleotide of a), d) a polynucleotide complementary to the polynucleotide of b), and e) an RNA equivalent of a)-d). The method comprises a) amplifying said target polynucleotide or fragment thereof using polymerase chain reaction amplification, and b) detecting the presence or absence of said amplified target polynucleotide or fragment thereof, and, optionally, if present, the amount thereof.

The invention further provides a composition comprising an effective amount of a polypeptide selected from the group consisting of a) a polypeptide comprising an amino acid sequence selected from the group consisting of SEQ ID NO:1-21, b) a polypeptide comprising a naturally occurring amino acid sequence at least 90% identical to an amino acid sequence selected from the group consisting of SEQ ID NO:1-21, c) a biologically active fragment of a polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO:1-21, and d) an immunogenic fragment of a polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO:1-21, and a pharmaceutically acceptable excipient. In one embodiment, the composition comprises an amino acid sequence selected from the group consisting of SEQ ID NO:1-21. The invention additionally provides a method of treating a disease or condition associated with decreased expression of functional PRTS, comprising administering to a patient in need of such treatment the composition.

The invention also provides a method for screening a compound for effectiveness as an agonist of a polypeptide selected from the group consisting of a) a polypeptide comprising an amino acid sequence selected from the group consisting of SEQ ID NO:1-21, b) a polypeptide comprising a naturally occurring amino acid sequence at least 90% identical to an amino acid sequence selected from the group consisting of SEQ ID NO:1-21, c) a biologically active fragment of a polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO:1-21, and d) an immunogenic fragment of a polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO:1-21. The method comprises a) exposing a sample comprising the polypeptide to a compound, and b) detecting agonist activity in the sample. In one alternative, the invention provides a composition comprising an agonist compound identified by the method and a pharmaceutically acceptable excipient. In another alternative, the invention provides a method of treating a disease or condition associated with decreased expression of functional PRTS, comprising administering to a patient in need of such treatment the composition.

Additionally, the invention provides a method for screening a compound for effectiveness as an antagonist of a polypeptide selected from the group consisting of a) a polypeptide comprising an

amino acid sequence selected from the group consisting of SEQ ID NO:1-21, b) a polypeptide comprising a naturally occurring amino acid sequence at least 90% identical to an amino acid sequence selected from the group consisting of SEQ ID NO:1-21, c) a biologically active fragment of a polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO:1-21, and
5 d) an immunogenic fragment of a polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO:1-21. The method comprises a) exposing a sample comprising the polypeptide to a compound, and b) detecting antagonist activity in the sample. In one alternative, the invention provides a composition comprising an antagonist compound identified by the method and a pharmaceutically acceptable excipient. In another alternative, the invention provides a method of
10 treating a disease or condition associated with overexpression of functional PRTS, comprising administering to a patient in need of such treatment the composition.

The invention further provides a method of screening for a compound that specifically binds to a polypeptide selected from the group consisting of a) a polypeptide comprising an amino acid sequence selected from the group consisting of SEQ ID NO:1-21, b) a polypeptide comprising a
15 naturally occurring amino acid sequence at least 90% identical to an amino acid sequence selected from the group consisting of SEQ ID NO:1-21, c) a biologically active fragment of a polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO:1-21, and d) an immunogenic fragment of a polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO:1-21. The method comprises a) combining the polypeptide with at least one test compound under
20 suitable conditions, and b) detecting binding of the polypeptide to the test compound, thereby identifying a compound that specifically binds to the polypeptide.

The invention further provides a method of screening for a compound that modulates the activity of a polypeptide selected from the group consisting of a) a polypeptide comprising an amino acid sequence selected from the group consisting of SEQ ID NO:1-21, b) a polypeptide comprising a
25 naturally occurring amino acid sequence at least 90% identical to an amino acid sequence selected from the group consisting of SEQ ID NO:1-21, c) a biologically active fragment of a polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO:1-21, and d) an immunogenic fragment of a polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO:1-21. The method comprises a) combining the polypeptide with at least one test compound under
30 conditions permissive for the activity of the polypeptide, b) assessing the activity of the polypeptide in the presence of the test compound, and c) comparing the activity of the polypeptide in the presence of the test compound with the activity of the polypeptide in the absence of the test compound, wherein a change in the activity of the polypeptide in the presence of the test compound is indicative of a compound that modulates the activity of the polypeptide.

The invention further provides a method for screening a compound for effectiveness in altering expression of a target polynucleotide, wherein said target polynucleotide comprises a sequence selected from the group consisting of SEQ ID NO:22-42, the method comprising a) exposing a sample comprising the target polynucleotide to a compound, and b) detecting altered
5 expression of the target polynucleotide.

The invention further provides a method for assessing toxicity of a test compound, said method comprising a) treating a biological sample containing nucleic acids with the test compound; b) hybridizing the nucleic acids of the treated biological sample with a probe comprising at least 20 contiguous nucleotides of a polynucleotide selected from the group consisting of i) a polynucleotide
10 comprising a polynucleotide sequence selected from the group consisting of SEQ ID NO:22-42, ii) a polynucleotide comprising a naturally occurring polynucleotide sequence at least 90% identical to a polynucleotide sequence selected from the group consisting of SEQ ID NO:22-42, iii) a polynucleotide having a sequence complementary to i), iv) a polynucleotide complementary to the polynucleotide of ii), and v) an RNA equivalent of i)-iv). Hybridization occurs under conditions
15 whereby a specific hybridization complex is formed between said probe and a target polynucleotide in the biological sample, said target polynucleotide selected from the group consisting of i) a polynucleotide comprising a polynucleotide sequence selected from the group consisting of SEQ ID NO:22-42, ii) a polynucleotide comprising a naturally occurring polynucleotide sequence at least 90% identical to a polynucleotide sequence selected from the group consisting of SEQ ID NO:22-42, iii) a
20 polynucleotide complementary to the polynucleotide of i), iv) a polynucleotide complementary to the polynucleotide of ii), and v) an RNA equivalent of i)-iv). Alternatively, the target polynucleotide comprises a fragment of a polynucleotide sequence selected from the group consisting of i)-v) above; c) quantifying the amount of hybridization complex; and d) comparing the amount of hybridization complex in the treated biological sample with the amount of hybridization complex in an untreated
25 biological sample, wherein a difference in the amount of hybridization complex in the treated biological sample is indicative of toxicity of the test compound.

BRIEF DESCRIPTION OF THE TABLES

Table 1 summarizes the nomenclature for the full length polynucleotide and polypeptide
30 sequences of the present invention.

Table 2 shows the GenBank identification number and annotation of the nearest GenBank homolog for polypeptides of the invention. The probability score for the match between each polypeptide and its GenBank homolog is also shown.

Table 3 shows structural features of polypeptide sequences of the invention, including predicted motifs and domains, along with the methods, algorithms, and searchable databases used for analysis of the polypeptides.

Table 4 lists the cDNA and/or genomic DNA fragments which were used to assemble
5 polynucleotide sequences of the invention, along with selected fragments of the polynucleotide sequences.

Table 5 shows the representative cDNA library for polynucleotides of the invention.

Table 6 provides an appendix which describes the tissues and vectors used for construction of the cDNA libraries shown in Table 5.

10 Table 7 shows the tools, programs, and algorithms used to analyze the polynucleotides and polypeptides of the invention, along with applicable descriptions, references, and threshold parameters.

DESCRIPTION OF THE INVENTION

Before the present proteins, nucleotide sequences, and methods are described, it is understood
15 that this invention is not limited to the particular machines, materials and methods described, as these may vary. It is also to be understood that the terminology used herein is for the purpose of describing particular embodiments only, and is not intended to limit the scope of the present invention which will be limited only by the appended claims.

It must be noted that as used herein and in the appended claims, the singular forms "a," "an,"
20 and "the" include plural reference unless the context clearly dictates otherwise. Thus, for example, a reference to "a host cell" includes a plurality of such host cells, and a reference to "an antibody" is a reference to one or more antibodies and equivalents thereof known to those skilled in the art, and so forth.

Unless defined otherwise, all technical and scientific terms used herein have the same meanings
25 as commonly understood by one of ordinary skill in the art to which this invention belongs. Although any machines, materials, and methods similar or equivalent to those described herein can be used to practice or test the present invention, the preferred machines, materials and methods are now described. All publications mentioned herein are cited for the purpose of describing and disclosing the cell lines, protocols, reagents and vectors which are reported in the publications and which might be used in
30 connection with the invention. Nothing herein is to be construed as an admission that the invention is not entitled to antedate such disclosure by virtue of prior invention.

DEFINITIONS

“PRTS” refers to the amino acid sequences of substantially purified PRTS obtained from any species, particularly a mammalian species, including bovine, ovine, porcine, murine, equine, and human, and from any source, whether natural, synthetic, semi-synthetic, or recombinant.

The term “agonist” refers to a molecule which intensifies or mimics the biological activity of PRTS. Agonists may include proteins, nucleic acids, carbohydrates, small molecules, or any other compound or composition which modulates the activity of PRTS either by directly interacting with PRTS or by acting on components of the biological pathway in which PRTS participates.

An “allelic variant” is an alternative form of the gene encoding PRTS. Allelic variants may result from at least one mutation in the nucleic acid sequence and may result in altered mRNAs or in polypeptides whose structure or function may or may not be altered. A gene may have none, one, or many allelic variants of its naturally occurring form. Common mutational changes which give rise to allelic variants are generally ascribed to natural deletions, additions, or substitutions of nucleotides. Each of these types of changes may occur alone, or in combination with the others, one or more times in a given sequence.

“Altered” nucleic acid sequences encoding PRTS include those sequences with deletions, insertions, or substitutions of different nucleotides, resulting in a polypeptide the same as PRTS or a polypeptide with at least one functional characteristic of PRTS. Included within this definition are polymorphisms which may or may not be readily detectable using a particular oligonucleotide probe of the polynucleotide encoding PRTS, and improper or unexpected hybridization to allelic variants, with a locus other than the normal chromosomal locus for the polynucleotide sequence encoding PRTS. The encoded protein may also be “altered,” and may contain deletions, insertions, or substitutions of amino acid residues which produce a silent change and result in a functionally equivalent PRTS. Deliberate amino acid substitutions may be made on the basis of similarity in polarity, charge, solubility, hydrophobicity, hydrophilicity, and/or the amphipathic nature of the residues, as long as the biological or immunological activity of PRTS is retained. For example, negatively charged amino acids may include aspartic acid and glutamic acid, and positively charged amino acids may include lysine and arginine. Amino acids with uncharged polar side chains having similar hydrophilicity values may include: asparagine and glutamine; and serine and threonine. Amino acids with uncharged side chains having similar hydrophilicity values may include: leucine, isoleucine, and valine; glycine and alanine; and phenylalanine and tyrosine.

The terms “amino acid” and “amino acid sequence” refer to an oligopeptide, peptide, polypeptide, or protein sequence, or a fragment of any of these, and to naturally occurring or synthetic

molecules. Where "amino acid sequence" is recited to refer to a sequence of a naturally occurring protein molecule, "amino acid sequence" and like terms are not meant to limit the amino acid sequence to the complete native amino acid sequence associated with the recited protein molecule.

"Amplification" relates to the production of additional copies of a nucleic acid sequence.

- 5 Amplification is generally carried out using polymerase chain reaction (PCR) technologies well known in the art.

The term "antagonist" refers to a molecule which inhibits or attenuates the biological activity of PRTS. Antagonists may include proteins such as antibodies, nucleic acids, carbohydrates, small molecules, or any other compound or composition which modulates the activity of PRTS either by
10 directly interacting with PRTS or by acting on components of the biological pathway in which PRTS participates.

The term "antibody" refers to intact immunoglobulin molecules as well as to fragments thereof, such as Fab, F(ab')₂, and Fv fragments, which are capable of binding an epitopic determinant. Antibodies that bind PRTS polypeptides can be prepared using intact polypeptides or using fragments
15 containing small peptides of interest as the immunizing antigen. The polypeptide or oligopeptide used to immunize an animal (e.g., a mouse, a rat, or a rabbit) can be derived from the translation of RNA, or synthesized chemically, and can be conjugated to a carrier protein if desired. Commonly used carriers that are chemically coupled to peptides include bovine serum albumin, thyroglobulin, and keyhole limpet hemocyanin (KLH). The coupled peptide is then used to immunize the animal.

20 The term "antigenic determinant" refers to that region of a molecule (i.e., an epitope) that makes contact with a particular antibody. When a protein or a fragment of a protein is used to immunize a host animal, numerous regions of the protein may induce the production of antibodies which bind specifically to antigenic determinants (particular regions or three-dimensional structures on the protein). An antigenic determinant may compete with the intact antigen (i.e., the immunogen used to
25 elicit the immune response) for binding to an antibody.

The term "antisense" refers to any composition capable of base-pairing with the "sense" (coding) strand of a specific nucleic acid sequence. Antisense compositions may include DNA; RNA; peptide nucleic acid (PNA); oligonucleotides having modified backbone linkages such as phosphorothioates, methylphosphonates, or benzylphosphonates; oligonucleotides having modified
30 sugar groups such as 2'-methoxyethyl sugars or 2'-methoxyethoxy sugars; or oligonucleotides having modified bases such as 5-methyl cytosine, 2'-deoxyuracil, or 7-deaza-2'-deoxyguanosine. Antisense molecules may be produced by any method including chemical synthesis or transcription. Once introduced into a cell, the complementary antisense molecule base-pairs with a naturally occurring nucleic acid sequence produced by the cell to form duplexes which block either transcription or

translation. The designation "negative" or "minus" can refer to the antisense strand, and the designation "positive" or "plus" can refer to the sense strand of a reference DNA molecule.

The term "biologically active" refers to a protein having structural, regulatory, or biochemical functions of a naturally occurring molecule. Likewise, "immunologically active" or "immunogenic" refers to the capability of the natural, recombinant, or synthetic PRTS, or of any oligopeptide thereof, to induce a specific immune response in appropriate animals or cells and to bind with specific antibodies.

"Complementary" describes the relationship between two single-stranded nucleic acid sequences that anneal by base-pairing. For example, 5'-AGT-3' pairs with its complement, 3'-TCA-5'.

A "composition comprising a given polynucleotide sequence" and a "composition comprising a given amino acid sequence" refer broadly to any composition containing the given polynucleotide or amino acid sequence. The composition may comprise a dry formulation or an aqueous solution. Compositions comprising polynucleotide sequences encoding PRTS or fragments of PRTS may be employed as hybridization probes. The probes may be stored in freeze-dried form and may be associated with a stabilizing agent such as a carbohydrate. In hybridizations, the probe may be deployed in an aqueous solution containing salts (e.g., NaCl), detergents (e.g., sodium dodecyl sulfate; SDS), and other components (e.g., Denhardt's solution, dry milk, salmon sperm DNA, etc.).

"Consensus sequence" refers to a nucleic acid sequence which has been subjected to repeated DNA sequence analysis to resolve uncalled bases, extended using the XL-PCR kit (Applied Biosystems, Foster City CA) in the 5' and/or the 3' direction, and resequenced, or which has been assembled from one or more overlapping cDNA, EST, or genomic DNA fragments using a computer program for fragment assembly, such as the GELVIEW fragment assembly system (GCG, Madison WI) or Phrap (University of Washington, Seattle WA). Some sequences have been both extended and assembled to produce the consensus sequence.

"Conservative amino acid substitutions" are those substitutions that are predicted to least interfere with the properties of the original protein, i.e., the structure and especially the function of the protein is conserved and not significantly changed by such substitutions. The table below shows amino acids which may be substituted for an original amino acid in a protein and which are regarded as conservative amino acid substitutions.

Original Residue	Conservative Substitution
Ala	Gly, Ser
Arg	His, Lys
Asn	Asp, Gln, His
Asp	Asn, Glu

	Cys	Ala, Ser
	Gln	Asn, Glu, His
	Glu	Asp, Gln, His
	Gly	Ala
5	His	Asn, Arg, Gln, Glu
	Ile	Leu, Val
	Leu	Ile, Val
	Lys	Arg, Gln, Glu
	Met	Leu, Ile
10	Phe	His, Met, Leu, Trp, Tyr
	Ser	Cys, Thr
	Thr	Ser, Val
	Trp	Phe, Tyr
	Tyr	His, Phe, Trp
15	Val	Ile, Leu, Thr

Conservative amino acid substitutions generally maintain (a) the structure of the polypeptide backbone in the area of the substitution, for example, as a beta sheet or alpha helical conformation, (b) the charge or hydrophobicity of the molecule at the site of the substitution, and/or (c) the bulk of the side chain.

A "deletion" refers to a change in the amino acid or nucleotide sequence that results in the absence of one or more amino acid residues or nucleotides.

The term "derivative" refers to a chemically modified polynucleotide or polypeptide. Chemical modifications of a polynucleotide can include, for example, replacement of hydrogen by an alkyl, acyl, hydroxyl, or amino group. A derivative polynucleotide encodes a polypeptide which retains at least one biological or immunological function of the natural molecule. A derivative polypeptide is one modified by glycosylation, pegylation, or any similar process that retains at least one biological or immunological function of the polypeptide from which it was derived.

A "detectable label" refers to a reporter molecule or enzyme that is capable of generating a measurable signal and is covalently or noncovalently joined to a polynucleotide or polypeptide.

"Differential expression" refers to increased or upregulated; or decreased, downregulated, or absent gene or protein expression, determined by comparing at least two different samples. Such comparisons may be carried out between, for example, a treated and an untreated sample, or a diseased and a normal sample.

A "fragment" is a unique portion of PRTS or the polynucleotide encoding PRTS which is identical in sequence to but shorter in length than the parent sequence. A fragment may comprise up to the entire length of the defined sequence, minus one nucleotide/amino acid residue. For example, a fragment may comprise from 5 to 1000 contiguous nucleotides or amino acid residues. A fragment used as a probe, primer, antigen, therapeutic molecule, or for other purposes, may be at least 5, 10,

15, 16, 20, 25, 30, 40, 50, 60, 75, 100, 150, 250 or at least 500 contiguous nucleotides or amino acid residues in length. Fragments may be preferentially selected from certain regions of a molecule. For example, a polypeptide fragment may comprise a certain length of contiguous amino acids selected from the first 250 or 500 amino acids (or first 25% or 50%) of a polypeptide as shown in a certain
5 defined sequence. Clearly these lengths are exemplary, and any length that is supported by the specification, including the Sequence Listing, tables, and figures, may be encompassed by the present embodiments.

A fragment of SEQ ID NO:22-42 comprises a region of unique polynucleotide sequence that specifically identifies SEQ ID NO:22-42, for example, as distinct from any other sequence in the
10 genome from which the fragment was obtained. A fragment of SEQ ID NO:22-42 is useful, for example, in hybridization and amplification technologies and in analogous methods that distinguish SEQ ID NO:22-42 from related polynucleotide sequences. The precise length of a fragment of SEQ ID NO:22-42 and the region of SEQ ID NO:22-42 to which the fragment corresponds are routinely determinable by one of ordinary skill in the art based on the intended purpose for the fragment.

15 A fragment of SEQ ID NO:1-21 is encoded by a fragment of SEQ ID NO:22-42. A fragment of SEQ ID NO:1-21 comprises a region of unique amino acid sequence that specifically identifies SEQ ID NO:1-21. For example, a fragment of SEQ ID NO:1-21 is useful as an immunogenic peptide for the development of antibodies that specifically recognize SEQ ID NO:1-21. The precise length of a fragment of SEQ ID NO:1-21 and the region of SEQ ID NO:1-21 to which the fragment
20 corresponds are routinely determinable by one of ordinary skill in the art based on the intended purpose for the fragment.

A "full length" polynucleotide sequence is one containing at least a translation initiation codon (e.g., methionine) followed by an open reading frame and a translation termination codon. A "full length" polynucleotide sequence encodes a "full length" polypeptide sequence.

25 "Homology" refers to sequence similarity or, interchangeably, sequence identity, between two or more polynucleotide sequences or two or more polypeptide sequences.

The terms "percent identity" and "% identity," as applied to polynucleotide sequences, refer to the percentage of residue matches between at least two polynucleotide sequences aligned using a standardized algorithm. Such an algorithm may insert, in a standardized and reproducible way, gaps in
30 the sequences being compared in order to optimize alignment between two sequences, and therefore achieve a more meaningful comparison of the two sequences.

Percent identity between polynucleotide sequences may be determined using the default parameters of the CLUSTAL V algorithm as incorporated into the MEGALIGN version 3.12e sequence alignment program. This program is part of the LASERGENE software package, a suite of molecular

biological analysis programs (DNASTAR, Madison WI). CLUSTAL V is described in Higgins, D.G. and P.M. Sharp (1989) CABIOS 5:151-153 and in Higgins, D.G. et al. (1992) CABIOS 8:189-191.

For pairwise alignments of polynucleotide sequences, the default parameters are set as follows:

5 Ktuple=2, gap penalty=5, window=4, and "diagonals saved"=4. The "weighted" residue weight table is selected as the default. Percent identity is reported by CLUSTAL V as the "percent similarity" between aligned polynucleotide sequences.

Alternatively, a suite of commonly used and freely available sequence comparison algorithms is provided by the National Center for Biotechnology Information (NCBI) Basic Local Alignment Search Tool (BLAST) (Altschul, S.F. et al. (1990) J. Mol. Biol. 215:403-410), which is available from several
10 sources, including the NCBI, Bethesda, MD, and on the Internet at <http://www.ncbi.nlm.nih.gov/BLAST/>. The BLAST software suite includes various sequence analysis programs including "blastn," that is used to align a known polynucleotide sequence with other polynucleotide sequences from a variety of databases. Also available is a tool called "BLAST 2 Sequences" that is used for direct pairwise comparison of two nucleotide sequences. "BLAST 2
15 Sequences" can be accessed and used interactively at <http://www.ncbi.nlm.nih.gov/gorf/bl2.html>. The "BLAST 2 Sequences" tool can be used for both blastn and blastp (discussed below). BLAST programs are commonly used with gap and other parameters set to default settings. For example, to compare two nucleotide sequences, one may use blastn with the "BLAST 2 Sequences" tool Version 2.0.12 (April-21-2000) set at default parameters. Such default parameters may be, for example:

20 *Matrix: BLOSUM62*
Reward for match: 1
Penalty for mismatch: -2
Open Gap: 5 and Extension Gap: 2 penalties
Gap x drop-off: 50
25 *Expect: 10*
Word Size: 11
Filter: on

Percent identity may be measured over the length of an entire defined sequence, for example, as defined by a particular SEQ ID number, or may be measured over a shorter length, for example, over
30 the length of a fragment taken from a larger, defined sequence, for instance, a fragment of at least 20, at least 30, at least 40, at least 50, at least 70, at least 100, or at least 200 contiguous nucleotides. Such lengths are exemplary only, and it is understood that any fragment length supported by the sequences shown herein, in the tables, figures, or Sequence Listing, may be used to describe a length over which percentage identity may be measured.

Nucleic acid sequences that do not show a high degree of identity may nevertheless encode similar amino acid sequences due to the degeneracy of the genetic code. It is understood that changes in a nucleic acid sequence can be made using this degeneracy to produce multiple nucleic acid sequences that all encode substantially the same protein.

5 The phrases "percent identity" and "% identity," as applied to polypeptide sequences, refer to the percentage of residue matches between at least two polypeptide sequences aligned using a standardized algorithm. Methods of polypeptide sequence alignment are well-known. Some alignment methods take into account conservative amino acid substitutions. Such conservative substitutions, explained in more detail above, generally preserve the charge and hydrophobicity at the site of
10 substitution, thus preserving the structure (and therefore function) of the polypeptide.

Percent identity between polypeptide sequences may be determined using the default parameters of the CLUSTAL V algorithm as incorporated into the MEGALIGN version 3.12e sequence alignment program (described and referenced above). For pairwise alignments of polypeptide sequences using CLUSTAL V, the default parameters are set as follows: Ktuple=1, gap penalty=3, window=5, and
15 "diagonals saved"=5. The PAM250 matrix is selected as the default residue weight table. As with polynucleotide alignments, the percent identity is reported by CLUSTAL V as the "percent similarity" between aligned polypeptide sequence pairs.

Alternatively the NCBI BLAST software suite may be used. For example, for a pairwise comparison of two polypeptide sequences, one may use the "BLAST 2 Sequences" tool Version 2.0.12
20 (April-21-2000) with blastp set at default parameters. Such default parameters may be, for example:

Matrix: BLOSUM62

Open Gap: 11 and Extension Gap: 1 penalties

Gap x drop-off: 50

Expect: 10

25 *Word Size: 3*

Filter: on

Percent identity may be measured over the length of an entire defined polypeptide sequence, for example, as defined by a particular SEQ ID number, or may be measured over a shorter length, for example, over the length of a fragment taken from a larger, defined polypeptide sequence, for instance,
30 a fragment of at least 15, at least 20, at least 30, at least 40, at least 50, at least 70 or at least 150 contiguous residues. Such lengths are exemplary only, and it is understood that any fragment length supported by the sequences shown herein, in the tables, figures or Sequence Listing, may be used to describe a length over which percentage identity may be measured.

"Human artificial chromosomes" (HACs) are linear microchromosomes which may contain DNA sequences of about 6 kb to 10 Mb in size and which contain all of the elements required for chromosome replication, segregation and maintenance.

The term "humanized antibody" refers to an antibody molecule in which the amino acid
5 sequence in the non-antigen binding regions has been altered so that the antibody more closely resembles a human antibody, and still retains its original binding ability.

"Hybridization" refers to the process by which a polynucleotide strand anneals with a complementary strand through base pairing under defined hybridization conditions. Specific hybridization is an indication that two nucleic acid sequences share a high degree of complementarity.
10 Specific hybridization complexes form under permissive annealing conditions and remain hybridized after the "washing" step(s). The washing step(s) is particularly important in determining the stringency of the hybridization process, with more stringent conditions allowing less non-specific binding, i.e., binding between pairs of nucleic acid strands that are not perfectly matched. Permissive conditions for annealing of nucleic acid sequences are routinely determinable by one of ordinary skill in the art and
15 may be consistent among hybridization experiments, whereas wash conditions may be varied among experiments to achieve the desired stringency, and therefore hybridization specificity. Permissive annealing conditions occur, for example, at 68°C in the presence of about 6 x SSC, about 1% (w/v) SDS, and about 100 µg/ml sheared, denatured salmon sperm DNA.

Generally, stringency of hybridization is expressed, in part, with reference to the temperature
20 under which the wash step is carried out. Such wash temperatures are typically selected to be about 5°C to 20°C lower than the thermal melting point (T_m) for the specific sequence at a defined ionic strength and pH. The T_m is the temperature (under defined ionic strength and pH) at which 50% of the target sequence hybridizes to a perfectly matched probe. An equation for calculating T_m and conditions for nucleic acid hybridization are well known and can be found in Sambrook, J. et al. (1989) Molecular
25 Cloning: A Laboratory Manual, 2nd ed., vol. 1-3, Cold Spring Harbor Press, Plainview NY; specifically see volume 2, chapter 9.

High stringency conditions for hybridization between polynucleotides of the present invention include wash conditions of 68°C in the presence of about 0.2 x SSC and about 0.1% SDS, for 1 hour. Alternatively, temperatures of about 65°C, 60°C, 55°C, or 42°C may be used. SSC concentration may
30 be varied from about 0.1 to 2 x SSC, with SDS being present at about 0.1%. Typically, blocking reagents are used to block non-specific hybridization. Such blocking reagents include, for instance, sheared and denatured salmon sperm DNA at about 100-200 µg/ml. Organic solvent, such as formamide at a concentration of about 35-50% v/v, may also be used under particular circumstances, such as for RNA:DNA hybridizations. Useful variations on these wash conditions will be readily

apparent to those of ordinary skill in the art. Hybridization, particularly under high stringency conditions, may be suggestive of evolutionary similarity between the nucleotides. Such similarity is strongly indicative of a similar role for the nucleotides and their encoded polypeptides.

The term "hybridization complex" refers to a complex formed between two nucleic acid sequences by virtue of the formation of hydrogen bonds between complementary bases. A hybridization complex may be formed in solution (e.g., C_0t or R_0t analysis) or formed between one nucleic acid sequence present in solution and another nucleic acid sequence immobilized on a solid support (e.g., paper, membranes, filters, chips, pins or glass slides, or any other appropriate substrate to which cells or their nucleic acids have been fixed).

The words "insertion" and "addition" refer to changes in an amino acid or nucleotide sequence resulting in the addition of one or more amino acid residues or nucleotides, respectively.

"Immune response" can refer to conditions associated with inflammation, trauma, immune disorders, or infectious or genetic disease, etc. These conditions can be characterized by expression of various factors, e.g., cytokines, chemokines, and other signaling molecules, which may affect cellular and systemic defense systems.

An "immunogenic fragment" is a polypeptide or oligopeptide fragment of PRTS which is capable of eliciting an immune response when introduced into a living organism, for example, a mammal. The term "immunogenic fragment" also includes any polypeptide or oligopeptide fragment of PRTS which is useful in any of the antibody production methods disclosed herein or known in the art.

The term "microarray" refers to an arrangement of a plurality of polynucleotides, polypeptides, or other chemical compounds on a substrate.

The terms "element" and "array element" refer to a polynucleotide, polypeptide, or other chemical compound having a unique and defined position on a microarray.

The term "modulate" refers to a change in the activity of PRTS. For example, modulation may cause an increase or a decrease in protein activity, binding characteristics, or any other biological, functional, or immunological properties of PRTS.

The phrases "nucleic acid" and "nucleic acid sequence" refer to a nucleotide, oligonucleotide, polynucleotide, or any fragment thereof. These phrases also refer to DNA or RNA of genomic or synthetic origin which may be single-stranded or double-stranded and may represent the sense or the antisense strand, to peptide nucleic acid (PNA), or to any DNA-like or RNA-like material.

"Operably linked" refers to the situation in which a first nucleic acid sequence is placed in a functional relationship with a second nucleic acid sequence. For instance, a promoter is operably linked to a coding sequence if the promoter affects the transcription or expression of the coding

sequence. Operably linked DNA sequences may be in close proximity or contiguous and, where necessary to join two protein coding regions, in the same reading frame.

5 "Peptide nucleic acid" (PNA) refers to an antisense molecule or anti-gene agent which comprises an oligonucleotide of at least about 5 nucleotides in length linked to a peptide backbone of amino acid residues ending in lysine. The terminal lysine confers solubility to the composition. PNAs preferentially bind complementary single stranded DNA or RNA and stop transcript elongation, and may be pegylated to extend their lifespan in the cell.

10 "Post-translational modification" of an PRTS may involve lipidation, glycosylation, phosphorylation, acetylation, racemization, proteolytic cleavage, and other modifications known in the art. These processes may occur synthetically or biochemically. Biochemical modifications will vary by cell type depending on the enzymatic milieu of PRTS.

"Probe" refers to nucleic acid sequences encoding PRTS, their complements, or fragments thereof, which are used to detect identical, allelic or related nucleic acid sequences. Probes are isolated oligonucleotides or polynucleotides attached to a detectable label or reporter molecule. Typical 15 labels include radioactive isotopes, ligands, chemiluminescent agents, and enzymes. "Primers" are short nucleic acids, usually DNA oligonucleotides, which may be annealed to a target polynucleotide by complementary base-pairing. The primer may then be extended along the target DNA strand by a DNA polymerase enzyme. Primer pairs can be used for amplification (and identification) of a nucleic acid sequence, e.g., by the polymerase chain reaction (PCR).

20 Probes and primers as used in the present invention typically comprise at least 15 contiguous nucleotides of a known sequence. In order to enhance specificity, longer probes and primers may also be employed, such as probes and primers that comprise at least 20, 25, 30, 40, 50, 60, 70, 80, 90, 100, or at least 150 consecutive nucleotides of the disclosed nucleic acid sequences. Probes and primers may be considerably longer than these examples, and it is understood that any length supported by the 25 specification, including the tables, figures, and Sequence Listing, may be used.

Methods for preparing and using probes and primers are described in the references, for example Sambrook, J. et al. (1989) Molecular Cloning: A Laboratory Manual, 2nd ed., vol. 1-3, Cold Spring Harbor Press, Plainview NY; Ausubel, F.M. et al. (1987) Current Protocols in Molecular Biology, Greene Publ. Assoc. & Wiley-Intersciences, New York NY; Innis, M. et al. (1990) PCR 30 Protocols, A Guide to Methods and Applications, Academic Press, San Diego CA. PCR primer pairs can be derived from a known sequence, for example, by using computer programs intended for that purpose such as Primer (Version 0.5, 1991, Whitehead Institute for Biomedical Research, Cambridge MA).

Oligonucleotides for use as primers are selected using software known in the art for such purpose. For example, OLIGO 4.06 software is useful for the selection of PCR primer pairs of up to 100 nucleotides each, and for the analysis of oligonucleotides and larger polynucleotides of up to 5,000 nucleotides from an input polynucleotide sequence of up to 32 kilobases. Similar primer selection programs have incorporated additional features for expanded capabilities. For example, the PrimOU primer selection program (available to the public from the Genome Center at University of Texas South West Medical Center, Dallas TX) is capable of choosing specific primers from megabase sequences and is thus useful for designing primers on a genome-wide scope. The Primer3 primer selection program (available to the public from the Whitehead Institute/MIT Center for Genome Research, Cambridge MA) allows the user to input a "mispriming library," in which sequences to avoid as primer binding sites are user-specified. Primer3 is useful, in particular, for the selection of oligonucleotides for microarrays. (The source code for the latter two primer selection programs may also be obtained from their respective sources and modified to meet the user's specific needs.) The PrimeGen program (available to the public from the UK Human Genome Mapping Project Resource Centre, Cambridge UK) designs primers based on multiple sequence alignments, thereby allowing selection of primers that hybridize to either the most conserved or least conserved regions of aligned nucleic acid sequences. Hence, this program is useful for identification of both unique and conserved oligonucleotides and polynucleotide fragments. The oligonucleotides and polynucleotide fragments identified by any of the above selection methods are useful in hybridization technologies, for example, as PCR or sequencing primers, microarray elements, or specific probes to identify fully or partially complementary polynucleotides in a sample of nucleic acids. Methods of oligonucleotide selection are not limited to those described above.

A "recombinant nucleic acid" is a sequence that is not naturally occurring or has a sequence that is made by an artificial combination of two or more otherwise separated segments of sequence. This artificial combination is often accomplished by chemical synthesis or, more commonly, by the artificial manipulation of isolated segments of nucleic acids, e.g., by genetic engineering techniques such as those described in Sambrook, *supra*. The term recombinant includes nucleic acids that have been altered solely by addition, substitution, or deletion of a portion of the nucleic acid. Frequently, a recombinant nucleic acid may include a nucleic acid sequence operably linked to a promoter sequence. Such a recombinant nucleic acid may be part of a vector that is used, for example, to transform a cell.

Alternatively, such recombinant nucleic acids may be part of a viral vector, e.g., based on a vaccinia virus, that could be used to vaccinate a mammal wherein the recombinant nucleic acid is expressed, inducing a protective immunological response in the mammal.

A "regulatory element" refers to a nucleic acid sequence usually derived from untranslated regions of a gene and includes enhancers, promoters, introns, and 5' and 3' untranslated regions (UTRs). Regulatory elements interact with host or viral proteins which control transcription, translation, or RNA stability.

5 "Reporter molecules" are chemical or biochemical moieties used for labeling a nucleic acid, amino acid, or antibody. Reporter molecules include radionuclides; enzymes; fluorescent, chemiluminescent, or chromogenic agents; substrates; cofactors; inhibitors; magnetic particles; and other moieties known in the art.

10 An "RNA equivalent," in reference to a DNA sequence, is composed of the same linear sequence of nucleotides as the reference DNA sequence with the exception that all occurrences of the nitrogenous base thymine are replaced with uracil, and the sugar backbone is composed of ribose instead of deoxyribose.

15 The term "sample" is used in its broadest sense. A sample suspected of containing PRTS, nucleic acids encoding PRTS, or fragments thereof may comprise a bodily fluid; an extract from a cell, chromosome, organelle, or membrane isolated from a cell; a cell; genomic DNA, RNA, or cDNA, in solution or bound to a substrate; a tissue; a tissue print; etc.

20 The terms "specific binding" and "specifically binding" refer to that interaction between a protein or peptide and an agonist, an antibody, an antagonist, a small molecule, or any natural or synthetic binding composition. The interaction is dependent upon the presence of a particular structure of the protein, e.g., the antigenic determinant or epitope, recognized by the binding molecule. For example, if an antibody is specific for epitope "A," the presence of a polypeptide comprising the epitope A, or the presence of free unlabeled A, in a reaction containing free labeled A and the antibody will reduce the amount of labeled A that binds to the antibody.

25 The term "substantially purified" refers to nucleic acid or amino acid sequences that are removed from their natural environment and are isolated or separated, and are at least 60% free, preferably at least 75% free, and most preferably at least 90% free from other components with which they are naturally associated.

A "substitution" refers to the replacement of one or more amino acid residues or nucleotides by different amino acid residues or nucleotides, respectively.

30 "Substrate" refers to any suitable rigid or semi-rigid support including membranes, filters, chips, slides, wafers, fibers, magnetic or nonmagnetic beads, gels, tubing, plates, polymers, microparticles and capillaries. The substrate can have a variety of surface forms, such as wells, trenches, pins, channels and pores, to which polynucleotides or polypeptides are bound.

A "transcript image" refers to the collective pattern of gene expression by a particular cell type or tissue under given conditions at a given time.

"Transformation" describes a process by which exogenous DNA is introduced into a recipient cell. Transformation may occur under natural or artificial conditions according to various methods well known in the art, and may rely on any known method for the insertion of foreign nucleic acid sequences into a prokaryotic or eukaryotic host cell. The method for transformation is selected based on the type of host cell being transformed and may include, but is not limited to, bacteriophage or viral infection, electroporation, heat shock, lipofection, and particle bombardment. The term "transformed cells" includes stably transformed cells in which the inserted DNA is capable of replication either as an autonomously replicating plasmid or as part of the host chromosome, as well as transiently transformed cells which express the inserted DNA or RNA for limited periods of time.

A "transgenic organism," as used herein, is any organism, including but not limited to animals and plants, in which one or more of the cells of the organism contains heterologous nucleic acid introduced by way of human intervention, such as by transgenic techniques well known in the art. The nucleic acid is introduced into the cell, directly or indirectly by introduction into a precursor of the cell, by way of deliberate genetic manipulation, such as by microinjection or by infection with a recombinant virus. The term genetic manipulation does not include classical cross-breeding, or in vitro fertilization, but rather is directed to the introduction of a recombinant DNA molecule. The transgenic organisms contemplated in accordance with the present invention include bacteria, cyanobacteria, fungi, plants and animals. The isolated DNA of the present invention can be introduced into the host by methods known in the art, for example infection, transfection, transformation or transconjugation. Techniques for transferring the DNA of the present invention into such organisms are widely known and provided in references such as Sambrook et al. (1989), supra.

A "variant" of a particular nucleic acid sequence is defined as a nucleic acid sequence having at least 40% sequence identity to the particular nucleic acid sequence over a certain length of one of the nucleic acid sequences using blastn with the "BLAST 2 Sequences" tool Version 2.0.9 (May-07-1999) set at default parameters. Such a pair of nucleic acids may show, for example, at least 50%, at least 60%, at least 70%, at least 80%, at least 85%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, or at least 99% or greater sequence identity over a certain defined length. A variant may be described as, for example, an "allelic" (as defined above), "splice," "species," or "polymorphic" variant. A splice variant may have significant identity to a reference molecule, but will generally have a greater or lesser number of polynucleotides due to alternative splicing of exons during mRNA processing. The corresponding polypeptide may

possess additional functional domains or lack domains that are present in the reference molecule. Species variants are polynucleotide sequences that vary from one species to another. The resulting polypeptides will generally have significant amino acid identity relative to each other. A polymorphic variant is a variation in the polynucleotide sequence of a particular gene between individuals of a given species. Polymorphic variants also may encompass "single nucleotide polymorphisms" (SNPs) in which the polynucleotide sequence varies by one nucleotide base. The presence of SNPs may be indicative of, for example, a certain population, a disease state, or a propensity for a disease state.

A "variant" of a particular polypeptide sequence is defined as a polypeptide sequence having at least 40% sequence identity to the particular polypeptide sequence over a certain length of one of the polypeptide sequences using blastp with the "BLAST 2 Sequences" tool Version 2.0.9 (May-07-1999) set at default parameters. Such a pair of polypeptides may show, for example, at least 50%, at least 60%, at least 70%, at least 80%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, or at least 99% or greater sequence identity over a certain defined length of one of the polypeptides.

THE INVENTION

The invention is based on the discovery of new human proteases (PRTS), the polynucleotides encoding PRTS, and the use of these compositions for the diagnosis, treatment, or prevention of gastrointestinal, cardiovascular, autoimmune/inflammatory, cell proliferative, developmental, epithelial, neurological, and reproductive disorders.

Table 1 summarizes the nomenclature for the full length polynucleotide and polypeptide sequences of the invention. Each polynucleotide and its corresponding polypeptide are correlated to a single Incyte project identification number (Incyte Project ID). Each polypeptide sequence is denoted by both a polypeptide sequence identification number (Polypeptide SEQ ID NO:) and an Incyte polypeptide sequence number (Incyte Polypeptide ID) as shown. Each polynucleotide sequence is denoted by both a polynucleotide sequence identification number (Polynucleotide SEQ ID NO:) and an Incyte polynucleotide consensus sequence number (Incyte Polynucleotide ID) as shown.

Table 2 shows sequences with homology to the polypeptides of the invention as identified by BLAST analysis against the GenBank protein (genpept) database. Columns 1 and 2 show the polypeptide sequence identification number (Polypeptide SEQ ID NO:) and the corresponding Incyte polypeptide sequence number (Incyte Polypeptide ID) for polypeptides of the invention. Column 3 shows the GenBank identification number (Genbank ID NO:) of the nearest GenBank homolog. Column 4 shows the probability score for the match between each polypeptide and its GenBank

homolog. Column 5 shows the annotation of the GenBank homolog along with relevant citations where applicable, all of which are expressly incorporated by reference herein.

Table 3 shows various structural features of the polypeptides of the invention. Columns 1 and 2 show the polypeptide sequence identification number (SEQ ID NO:) and the corresponding Incyte polypeptide sequence number (Incyte Polypeptide ID) for each polypeptide of the invention. Column 3 shows the number of amino acid residues in each polypeptide. Column 4 shows potential phosphorylation sites, and column 5 shows potential glycosylation sites, as determined by the MOTIFS program of the GCG sequence analysis software package (Genetics Computer Group, Madison WI). Column 6 shows amino acid residues comprising signature sequences, domains, and motifs. Column 7 shows analytical methods for protein structure/function analysis and in some cases, searchable databases to which the analytical methods were applied.

Together, Tables 2 and 3 summarize the properties of polypeptides of the invention, and these properties establish that the claimed polypeptides are proteases. For example, SEQ ID NO:1 is a ubiquitin carboxyl terminal hydrolase. SEQ ID NO:1 is 48% identical, from residue M1 to residue G225, to human ubiquitin-specific processing protease (GenBank ID g9971757) as determined by the Basic Local Alignment Search Tool (BLAST). (See Table 2.) The BLAST probability score is 1.00e-49, which indicates the probability of obtaining the observed polypeptide sequence alignment by chance. SEQ ID NO:1 contains a ubiquitin carboxyl terminal hydrolase catalytic site domain as determined by searching for statistically significant matches in the hidden Markov model (HMM)-based PFAM database of conserved protein family domains. The score is 53.4 bits and the E-value is 4.9e-12, which indicates the probability of obtaining the observed structural motif by chance. The presence of this motif was corroborated by BLIMPS (probability score=2.6e-4) and MOTIFS analyses. This provides further evidence that SEQ ID NO:1 is a ubiquitin carboxyl-terminal hydrolase. In an alternative example, SEQ ID NO:2 is 45% identical to amino acids 15-235 of human prostaticin, a serine protease (GenBank ID g1143194) as determined by the Basic Local Alignment Search Tool (BLAST). (See Table 2.) The BLAST probability score is 1.3e-46, which indicates the probability of obtaining the observed polypeptide sequence alignment by chance. SEQ ID NO:2 also contains a trypsin family serine protease active site domain as determined by searching for statistically significant matches in the hidden Markov model (HMM)-based PFAM database of conserved protein family domains. This match has a probability score of 2.7e-58. BLIMPS, MOTIFS, and PROFILESCAN analyses confirm the presence of this domain. (See Table 3.) BLIMPS analysis also reveals a kringle domain, providing further corroborative evidence that SEQ ID NO:2 is a serine protease of the trypsin family. In an alternative example, SEQ ID NO:7 is a dipeptidase which hydrolyses a variety of peptides (Kozak, E. and S. Tate (1982) J. Biol. Chem. 257:6322-6327), and is responsible for the hydrolysis of the beta

lactam rings of antibiotics such as penem and carbapenem (Campbell et al., (1984) J. Biol. Chem. 259:14586-14590). SEQ ID NO:7 shows 48% amino acid sequence identity over 377 amino acids (total length equals 411 amino acids) to human dipeptidase precursor (GenBank ID g219600) as determined by Basic Local Alignment Search Tool (BLAST). The BLAST probability score is 1.1×10^{-88} , which indicates the probability of obtaining the observed polypeptide sequence alignment by chance. Additionally, the protease of the invention demonstrates a renal dipeptidase domain as determined by searching for statistically significant matches in the hidden Markov model (HMM)-based PFAM database of conserved protein family domains. The HMM score for the renal dipeptidase PFAM hit is 412.7. Data from BLIMPS, MOTIFS, BLAST-DOMO, and BLAST-PRODOM analyses provide further corroborative evidence that SEQ ID NO:7 is a renal dipeptidase. The BLIMPS-BLOCKS hit scores for localized regions range from 1040-1537. The BLAST-DOMO hit probability score is 5.2×10^{-85} . The BLAST-PRODOM hit probability score is 4.7×10^{-73} . In an alternative example, SEQ ID NO:8 is 86% identical to human transmembrane trypsin (GenBank ID g6103629) as determined by the Basic Local Alignment Search Tool (BLAST). (See Table 2.) The BLAST probability score is 3.9×10^{-166} , which indicates the probability of obtaining the observed polypeptide sequence alignment by chance. SEQ ID NO:8 contains a trypsin family protease active site domain with a probability score of 5.3×10^{-89} as determined by searching for matches in the hidden Markov model (HMM)-based PFAM database of conserved protein family domains. BLIMPS, MOTIFS, and PROFILESCAN analyses confirm the presence of this motif. BLIMPS analysis also shows that SEQ ID NO:8 contains a kringle domain and a type I fibronectin domain. HMMER-based analysis reveals the presence of a transmembrane domain (See Table 3.). Taken together, these analyses show that SEQ ID NO:8 is a transmembrane member of the trypsin family of serine proteases. In an alternative example, SEQ ID NO:17 shares 44% local identity with human membrane-type serine protease 1 (MT-SP1, GenBank ID g6002714) as determined by the Basic Local Alignment Search Tool (BLAST). (See Table 2.) The BLAST probability score is 5.1×10^{-94} , which indicates the probability of obtaining the observed polypeptide sequence alignment by chance. SEQ ID NO:17 contains a trypsin family serine protease active site domain as determined by searching for statistically significant matches in the hidden Markov model (HMM)-based PFAM database of conserved protein family domains. (See Table 3.) HMM-based analysis also reveals a transmembrane domain near the N-terminus of SEQ ID NO:17. A domain found in the low-density lipoprotein receptor and other proteins, including MT-SP1 (PDOC00929) was also identified in this way. The presence of the trypsin active site motif is confirmed by PROFILESCAN, BLIMPS, and MOTIFS analyses. BLIMPS analysis revealed the presence of kringle and type I fibronectin domains. Taken together, these data provide further corroborative evidence that SEQ ID NO:17 is a transmembrane member of the trypsin family of serine proteases. SEQ ID NO:3-6, SEQ ID NO:9-16,

and SEQ ID NO:18-21 were analyzed and annotated in a similar manner. The algorithms and parameters for the analysis of SEQ ID NO:1-21 are described in Table 7.

As shown in Table 4, the full length polynucleotide sequences of the present invention were assembled using cDNA sequences or coding (exon) sequences derived from genomic DNA, or any combination of these two types of sequences. Columns 1 and 2 list the polynucleotide sequence identification number (Polynucleotide SEQ ID NO:) and the corresponding Incyte polynucleotide consensus sequence number (Incyte Polynucleotide ID) for each polynucleotide of the invention. Column 3 shows the length of each polynucleotide sequence in basepairs. Column 4 lists fragments of the polynucleotide sequences which are useful, for example, in hybridization or amplification technologies that identify SEQ ID NO:22-42 or that distinguish between SEQ ID NO:22-42 and related polynucleotide sequences. Column 5 shows identification numbers corresponding to cDNA sequences, coding sequences (exons) predicted from genomic DNA, and/or sequence assemblages comprised of both cDNA and genomic DNA. These sequences were used to assemble the full length polynucleotide sequences of the invention. Columns 6 and 7 of Table 4 show the nucleotide start (5') and stop (3') positions of the cDNA and/or genomic sequences in column 5 relative to their respective full length sequences.

The identification numbers in Column 5 of Table 4 may refer specifically, for example, to Incyte cDNAs along with their corresponding cDNA libraries. For example, 7246467T8 is the identification number of an Incyte cDNA sequence, and PROSTMY01 is the cDNA library from which it is derived. Incyte cDNAs for which cDNA libraries are not indicated were derived from pooled cDNA libraries (e.g., 71041539V1). Alternatively, the identification numbers in column 5 may refer to GenBank cDNAs or ESTs (e.g., g5745066) which contributed to the assembly of the full length polynucleotide sequences. Alternatively, the identification numbers in column 5 may refer to coding regions predicted by Genscan analysis of genomic DNA. For example, GNN.g7208751_000002_002.edit is the identification number of a Genscan-predicted coding sequence, with g7208751 being the GenBank identification number of the sequence to which Genscan was applied. The Genscan-predicted coding sequences may have been edited prior to assembly. (See Example IV.) Alternatively, the identification numbers in column 5 may refer to assemblages of both cDNA and Genscan-predicted exons brought together by an "exon stitching" algorithm. For example, FL1389845_00001 represents a "stitched" sequence in which 1389845 is the identification number of the cluster of sequences to which the algorithm was applied, and 00001 is the number of the prediction generated by the algorithm. (See Example V.) Alternatively, the identification numbers in column 5 may refer to assemblages of both cDNA and Genscan-predicted exons brought together by an "exon-stretching" algorithm. For example, FL2256251_g7708357_000002_g6103629 is the identification

number of a "stretched" sequence, with 2256251 being the Incyte project identification number, g7708357 being the GenBank identification number of the human genomic sequence to which the "exon-stretching" algorithm was applied, and g6103629 being the GenBank identification number of the nearest GenBank protein homolog. (See Example V.) In some cases, Incyte cDNA coverage
5 redundant with the sequence coverage shown in column 5 was obtained to confirm the final consensus polynucleotide sequence, but the relevant Incyte cDNA identification numbers are not shown.

Table 5 shows the representative cDNA libraries for those full length polynucleotide sequences which were assembled using Incyte cDNA sequences. The representative cDNA library is the Incyte cDNA library which is most frequently represented by the Incyte cDNA sequences which were used to
10 assemble and confirm the above polynucleotide sequences. The tissues and vectors which were used to construct the cDNA libraries shown in Table 5 are described in Table 6.

The invention also encompasses PRTS variants. A preferred PRTS variant is one which has at least about 80%, or alternatively at least about 90%, or even at least about 95% amino acid sequence identity to the PRTS amino acid sequence, and which contains at least one functional or structural
15 characteristic of PRTS.

The invention also encompasses polynucleotides which encode PRTS. In a particular embodiment, the invention encompasses a polynucleotide sequence comprising a sequence selected from the group consisting of SEQ ID NO:22-42, which encodes PRTS. The polynucleotide sequences of SEQ ID NO:22-42, as presented in the Sequence Listing, embrace the equivalent RNA sequences,
20 wherein occurrences of the nitrogenous base thymine are replaced with uracil, and the sugar backbone is composed of ribose instead of deoxyribose.

The invention also encompasses a variant of a polynucleotide sequence encoding PRTS. In particular, such a variant polynucleotide sequence will have at least about 70%, or alternatively at least about 85%, or even at least about 95% polynucleotide sequence identity to the polynucleotide sequence encoding PRTS. A particular aspect of the invention encompasses a variant of a polynucleotide
25 sequence comprising a sequence selected from the group consisting of SEQ ID NO:22-42 which has at least about 70%, or alternatively at least about 85%, or even at least about 95% polynucleotide sequence identity to a nucleic acid sequence selected from the group consisting of SEQ ID NO:22-42. Any one of the polynucleotide variants described above can encode an amino acid sequence which
30 contains at least one functional or structural characteristic of PRTS.

It will be appreciated by those skilled in the art that as a result of the degeneracy of the genetic code, a multitude of polynucleotide sequences encoding PRTS, some bearing minimal similarity to the polynucleotide sequences of any known and naturally occurring gene, may be produced. Thus, the invention contemplates each and every possible variation of polynucleotide sequence that could be made

by selecting combinations based on possible codon choices. These combinations are made in accordance with the standard triplet genetic code as applied to the polynucleotide sequence of naturally occurring PRTS, and all such variations are to be considered as being specifically disclosed.

Although nucleotide sequences which encode PRTS and its variants are generally capable of
5 hybridizing to the nucleotide sequence of the naturally occurring PRTS under appropriately selected conditions of stringency, it may be advantageous to produce nucleotide sequences encoding PRTS or its derivatives possessing a substantially different codon usage, e.g., inclusion of non-naturally occurring codons. Codons may be selected to increase the rate at which expression of the peptide occurs in a particular prokaryotic or eukaryotic host in accordance with the frequency with which particular codons
10 are utilized by the host. Other reasons for substantially altering the nucleotide sequence encoding PRTS and its derivatives without altering the encoded amino acid sequences include the production of RNA transcripts having more desirable properties, such as a greater half-life, than transcripts produced from the naturally occurring sequence.

The invention also encompasses production of DNA sequences which encode PRTS and PRTS
15 derivatives, or fragments thereof, entirely by synthetic chemistry. After production, the synthetic sequence may be inserted into any of the many available expression vectors and cell systems using reagents well known in the art. Moreover, synthetic chemistry may be used to introduce mutations into a sequence encoding PRTS or any fragment thereof.

Also encompassed by the invention are polynucleotide sequences that are capable of
20 hybridizing to the claimed polynucleotide sequences, and, in particular, to those shown in SEQ ID NO:22-42 and fragments thereof under various conditions of stringency. (See, e.g., Wahl, G.M. and S.L. Berger (1987) *Methods Enzymol.* 152:399-407; Kimmel, A.R. (1987) *Methods Enzymol.* 152:507-511.) Hybridization conditions, including annealing and wash conditions, are described in "Definitions."

25 Methods for DNA sequencing are well known in the art and may be used to practice any of the embodiments of the invention. The methods may employ such enzymes as the Klenow fragment of DNA polymerase I, SEQUENASE (US Biochemical, Cleveland OH), Taq polymerase (Applied Biosystems), thermostable T7 polymerase (Amersham Pharmacia Biotech, Piscataway NJ), or combinations of polymerases and proofreading exonucleases such as those found in the ELONGASE
30 amplification system (Life Technologies, Gaithersburg MD). Preferably, sequence preparation is automated with machines such as the MICROLAB 2200 liquid transfer system (Hamilton, Reno NV), PTC200 thermal cycler (MJ Research, Watertown MA) and ABI CATALYST 800 thermal cycler (Applied Biosystems). Sequencing is then carried out using either the ABI 373 or 377 DNA sequencing system (Applied Biosystems), the MEGABACE 1000 DNA sequencing system (Molecular Dynamics,

Sunnyvale CA), or other systems known in the art. The resulting sequences are analyzed using a variety of algorithms which are well known in the art. (See, e.g., Ausubel, F.M. (1997) Short Protocols in Molecular Biology, John Wiley & Sons, New York NY, unit 7.7; Meyers, R.A. (1995) Molecular Biology and Biotechnology, Wiley VCH, New York NY, pp. 856-853.)

- 5 The nucleic acid sequences encoding PRTS may be extended utilizing a partial nucleotide sequence and employing various PCR-based methods known in the art to detect upstream sequences, such as promoters and regulatory elements. For example, one method which may be employed, restriction-site PCR, uses universal and nested primers to amplify unknown sequence from genomic DNA within a cloning vector. (See, e.g., Sarkar, G. (1993) PCR Methods Applic. 2:318-322.)
- 10 Another method, inverse PCR, uses primers that extend in divergent directions to amplify unknown sequence from a circularized template. The template is derived from restriction fragments comprising a known genomic locus and surrounding sequences. (See, e.g., Triglia, T. et al. (1988) Nucleic Acids Res. 16:8186.) A third method, capture PCR, involves PCR amplification of DNA fragments adjacent to known sequences in human and yeast artificial chromosome DNA. (See, e.g., Lagerstrom, M. et al.
- 15 (1991) PCR Methods Applic. 1:111-119.) In this method, multiple restriction enzyme digestions and ligations may be used to insert an engineered double-stranded sequence into a region of unknown sequence before performing PCR. Other methods which may be used to retrieve unknown sequences are known in the art. (See, e.g., Parker, J.D. et al. (1991) Nucleic Acids Res. 19:3055-3060). Additionally, one may use PCR, nested primers, and PROMOTERFINDER libraries (Clontech, Palo
- 20 Alto CA) to walk genomic DNA. This procedure avoids the need to screen libraries and is useful in finding intron/exon junctions. For all PCR-based methods, primers may be designed using commercially available software, such as OLIGO 4.06 primer analysis software (National Biosciences, Plymouth MN) or another appropriate program, to be about 22 to 30 nucleotides in length, to have a GC content of about 50% or more, and to anneal to the template at temperatures of about 68°C to
- 25 72°C.

- When screening for full length cDNAs, it is preferable to use libraries that have been size-selected to include larger cDNAs. In addition, random-primed libraries, which often include sequences containing the 5' regions of genes, are preferable for situations in which an oligo d(T) library does not yield a full-length cDNA. Genomic libraries may be useful for extension of sequence into 5'
- 30 non-transcribed regulatory regions.

 Capillary electrophoresis systems which are commercially available may be used to analyze the size or confirm the nucleotide sequence of sequencing or PCR products. In particular, capillary sequencing may employ flowable polymers for electrophoretic separation, four different nucleotide-specific, laser-stimulated fluorescent dyes, and a charge coupled device camera for detection of the

emitted wavelengths. Output/light intensity may be converted to electrical signal using appropriate software (e.g., GENOTYPER and SEQUENCE NAVIGATOR, Applied Biosystems), and the entire process from loading of samples to computer analysis and electronic data display may be computer controlled. Capillary electrophoresis is especially preferable for sequencing small DNA fragments
5 which may be present in limited amounts in a particular sample.

In another embodiment of the invention, polynucleotide sequences or fragments thereof which encode PRTS may be cloned in recombinant DNA molecules that direct expression of PRTS; or fragments or functional equivalents thereof, in appropriate host cells. Due to the inherent degeneracy of the genetic code, other DNA sequences which encode substantially the same or a functionally equivalent
10 amino acid sequence may be produced and used to express PRTS.

The nucleotide sequences of the present invention can be engineered using methods generally known in the art in order to alter PRTS-encoding sequences for a variety of purposes including, but not limited to, modification of the cloning, processing, and/or expression of the gene product. DNA shuffling by random fragmentation and PCR reassembly of gene fragments and synthetic
15 oligonucleotides may be used to engineer the nucleotide sequences. For example, oligonucleotide-mediated site-directed mutagenesis may be used to introduce mutations that create new restriction sites, alter glycosylation patterns, change codon preference, produce splice variants, and so forth.

The nucleotides of the present invention may be subjected to DNA shuffling techniques such as MOLECULARBREEDING (Maxygen Inc., Santa Clara CA; described in U.S. Patent Number
20 5,837,458; Chang, C.-C. et al. (1999) Nat. Biotechnol. 17:793-797; Christians, F.C. et al. (1999) Nat. Biotechnol. 17:259-264; and Cramer, A. et al. (1996) Nat. Biotechnol. 14:315-319) to alter or improve the biological properties of PRTS, such as its biological or enzymatic activity or its ability to bind to other molecules or compounds. DNA shuffling is a process by which a library of gene variants is produced using PCR-mediated recombination of gene fragments. The library is then
25 subjected to selection or screening procedures that identify those gene variants with the desired properties. These preferred variants may then be pooled and further subjected to recursive rounds of DNA shuffling and selection/screening. Thus, genetic diversity is created through "artificial" breeding and rapid molecular evolution. For example, fragments of a single gene containing random point mutations may be recombined, screened, and then reshuffled until the desired properties are
30 optimized. Alternatively, fragments of a given gene may be recombined with fragments of homologous genes in the same gene family, either from the same or different species, thereby maximizing the genetic diversity of multiple naturally occurring genes in a directed and controllable manner.

In another embodiment, sequences encoding PRTS may be synthesized, in whole or in part, using chemical methods well known in the art. (See, e.g., Caruthers, M.H. et al. (1980) *Nucleic Acids Symp. Ser.* 7:215-223; and Horn, T. et al. (1980) *Nucleic Acids Symp. Ser.* 7:225-232.) Alternatively, PRTS itself or a fragment thereof may be synthesized using chemical methods. For example, peptide
5 synthesis can be performed using various solution-phase or solid-phase techniques. (See, e.g., Creighton, T. (1984) *Proteins, Structures and Molecular Properties*, WH Freeman, New York NY, pp. 55-60; and Roberge, J.Y. et al. (1995) *Science* 269:202-204.) Automated synthesis may be achieved using the ABI 431A peptide synthesizer (Applied Biosystems). Additionally, the amino acid sequence of PRTS, or any part thereof, may be altered during direct synthesis and/or combined with sequences
10 from other proteins, or any part thereof, to produce a variant polypeptide or a polypeptide having a sequence of a naturally occurring polypeptide.

The peptide may be substantially purified by preparative high performance liquid chromatography. (See, e.g., Chiez, R.M. and F.Z. Regnier (1990) *Methods Enzymol.* 182:392-421.) The composition of the synthetic peptides may be confirmed by amino acid analysis or by sequencing.
15 (See, e.g., Creighton, *supra*, pp. 28-53.)

In order to express a biologically active PRTS, the nucleotide sequences encoding PRTS or derivatives thereof may be inserted into an appropriate expression vector, i.e., a vector which contains the necessary elements for transcriptional and translational control of the inserted coding sequence in a suitable host. These elements include regulatory sequences, such as enhancers, constitutive and
20 inducible promoters, and 5' and 3' untranslated regions in the vector and in polynucleotide sequences encoding PRTS. Such elements may vary in their strength and specificity. Specific initiation signals may also be used to achieve more efficient translation of sequences encoding PRTS. Such signals include the ATG initiation codon and adjacent sequences, e.g. the Kozak sequence. In cases where sequences encoding PRTS and its initiation codon and upstream regulatory sequences are inserted into
25 the appropriate expression vector, no additional transcriptional or translational control signals may be needed. However, in cases where only coding sequence, or a fragment thereof, is inserted, exogenous translational control signals including an in-frame ATG initiation codon should be provided by the vector. Exogenous translational elements and initiation codons may be of various origins, both natural and synthetic. The efficiency of expression may be enhanced by the inclusion of enhancers appropriate
30 for the particular host cell system used. (See, e.g., Scharf, D. et al. (1994) *Results Probl. Cell Differ.* 20:125-162.)

Methods which are well known to those skilled in the art may be used to construct expression vectors containing sequences encoding PRTS and appropriate transcriptional and translational control elements. These methods include *in vitro* recombinant DNA techniques, synthetic techniques, and *in*

vivo genetic recombination. (See, e.g., Sambrook, J. et al. (1989) Molecular Cloning, A Laboratory Manual, Cold Spring Harbor Press, Plainview NY, ch. 4, 8, and 16-17; Ausubel, F.M. et al. (1995) Current Protocols in Molecular Biology, John Wiley & Sons, New York NY, ch. 9, 13, and 16.)

A variety of expression vector/host systems may be utilized to contain and express sequences
5 encoding PRTS. These include, but are not limited to, microorganisms such as bacteria transformed
with recombinant bacteriophage, plasmid, or cosmid DNA expression vectors; yeast transformed with
yeast expression vectors; insect cell systems infected with viral expression vectors (e.g., baculovirus);
plant cell systems transformed with viral expression vectors (e.g., cauliflower mosaic virus, CaMV, or
tobacco mosaic virus, TMV) or with bacterial expression vectors (e.g., Ti or pBR322 plasmids); or
10 animal cell systems. (See, e.g., Sambrook, supra; Ausubel, supra; Van Heeke, G. and S.M. Schuster
(1989) J. Biol. Chem. 264:5503-5509; Engelhard, E.K. et al. (1994) Proc. Natl. Acad. Sci. USA
91:3224-3227; Sandig, V. et al. (1996) Hum. Gene Ther. 7:1937-1945; Takamatsu, N. (1987) EMBO
J. 6:307-311; The McGraw Hill Yearbook of Science and Technology (1992) McGraw Hill, New
York NY, pp. 191-196; Logan, J. and T. Shenk (1984) Proc. Natl. Acad. Sci. USA 81:3655-3659; and
15 Harrington, J.J. et al. (1997) Nat. Genet. 15:345-355.) Expression vectors derived from retroviruses,
adenoviruses, or herpes or vaccinia viruses, or from various bacterial plasmids, may be used for
delivery of nucleotide sequences to the targeted organ, tissue, or cell population. (See, e.g., Di
Nicola, M. et al. (1998) Cancer Gen. Ther. 5(6):350-356; Yu, M. et al. (1993) Proc. Natl. Acad. Sci.
USA 90(13):6340-6344; Buller, R.M. et al. (1985) Nature 317(6040):813-815; McGregor, D.P. et al.
20 (1994) Mol. Immunol. 31(3):219-226; and Verma, I.M. and N. Somia (1997) Nature 389:239-242.)
The invention is not limited by the host cell employed.

In bacterial systems, a number of cloning and expression vectors may be selected depending
upon the use intended for polynucleotide sequences encoding PRTS. For example, routine cloning,
subcloning, and propagation of polynucleotide sequences encoding PRTS can be achieved using a
25 multifunctional E. coli vector such as PBLUESCRIPT (Stratagene, La Jolla CA) or PSPORT1 plasmid
(Life Technologies). Ligation of sequences encoding PRTS into the vector's multiple cloning site
disrupts the *lacZ* gene, allowing a colorimetric screening procedure for identification of transformed
bacteria containing recombinant molecules. In addition, these vectors may be useful for in vitro
transcription, dideoxy sequencing, single strand rescue with helper phage, and creation of nested
30 deletions in the cloned sequence. (See, e.g., Van Heeke, G. and S.M. Schuster (1989) J. Biol. Chem.
264:5503-5509.) When large quantities of PRTS are needed, e.g. for the production of antibodies,
vectors which direct high level expression of PRTS may be used. For example, vectors containing the
strong, inducible SP6 or T7 bacteriophage promoter may be used.

Yeast expression systems may be used for production of PRTS. A number of vectors containing constitutive or inducible promoters, such as alpha factor, alcohol oxidase, and PGH promoters, may be used in the yeast *Saccharomyces cerevisiae* or *Pichia pastoris*. In addition, such vectors direct either the secretion or intracellular retention of expressed proteins and enable integration of foreign sequences into the host genome for stable propagation. (See, e.g., Ausubel, 1995, *supra*; Bitter, G.A. et al. (1987) *Methods Enzymol.* 153:516-544; and Scorer, C.A. et al. (1994) *Bio/Technology* 12:181-184.)

Plant systems may also be used for expression of PRTS. Transcription of sequences encoding PRTS may be driven by viral promoters, e.g., the 35S and 19S promoters of CaMV used alone or in combination with the omega leader sequence from TMV (Takamatsu, N. (1987) *EMBO J.* 6:307-311). Alternatively, plant promoters such as the small subunit of RUBISCO or heat shock promoters may be used. (See, e.g., Coruzzi, G. et al. (1984) *EMBO J.* 3:1671-1680; Broglie, R. et al. (1984) *Science* 224:838-843; and Winter, J. et al. (1991) *Results Probl. Cell Differ.* 17:85-105.) These constructs can be introduced into plant cells by direct DNA transformation or pathogen-mediated transfection. (See, e.g., *The McGraw Hill Yearbook of Science and Technology* (1992) McGraw Hill, New York NY, pp. 191-196.)

In mammalian cells, a number of viral-based expression systems may be utilized. In cases where an adenovirus is used as an expression vector, sequences encoding PRTS may be ligated into an adenovirus transcription/translation complex consisting of the late promoter and tripartite leader sequence. Insertion in a non-essential E1 or E3 region of the viral genome may be used to obtain infective virus which expresses PRTS in host cells. (See, e.g., Logan, J. and T. Shenk (1984) *Proc. Natl. Acad. Sci. USA* 81:3655-3659.) In addition, transcription enhancers, such as the Rous sarcoma virus (RSV) enhancer, may be used to increase expression in mammalian host cells. SV40 or EBV-based vectors may also be used for high-level protein expression.

Human artificial chromosomes (HACs) may also be employed to deliver larger fragments of DNA than can be contained in and expressed from a plasmid. HACs of about 6 kb to 10 Mb are constructed and delivered via conventional delivery methods (liposomes, polycationic amino polymers, or vesicles) for therapeutic purposes. (See, e.g., Harrington, J.J. et al. (1997) *Nat. Genet.* 15:345-355.)

For long term production of recombinant proteins in mammalian systems, stable expression of PRTS in cell lines is preferred. For example, sequences encoding PRTS can be transformed into cell lines using expression vectors which may contain viral origins of replication and/or endogenous expression elements and a selectable marker gene on the same or on a separate vector. Following the introduction of the vector, cells may be allowed to grow for about 1 to 2 days in enriched media before being switched to selective media. The purpose of the selectable marker is to confer resistance to a

selective agent, and its presence allows growth and recovery of cells which successfully express the introduced sequences. Resistant clones of stably transformed cells may be propagated using tissue culture techniques appropriate to the cell type.

Any number of selection systems may be used to recover transformed cell lines. These include, but are not limited to, the herpes simplex virus thymidine kinase and adenine phosphoribosyltransferase genes, for use in *tk* and *apr* cells, respectively. (See, e.g., Wigler, M. et al. (1977) Cell 11:223-232; Lowy, I. et al. (1980) Cell 22:817-823.) Also, antimetabolite, antibiotic, or herbicide resistance can be used as the basis for selection. For example, *dhfr* confers resistance to methotrexate; *neo* confers resistance to the aminoglycosides neomycin and G-418; and *als* and *pat* confer resistance to chlorsulfuron and phosphinotricin acetyltransferase, respectively. (See, e.g., Wigler, M. et al. (1980) Proc. Natl. Acad. Sci. USA 77:3567-3570; Colbere-Garapin, F. et al. (1981) J. Mol. Biol. 150:1-14.) Additional selectable genes have been described, e.g., *trpB* and *hisD*, which alter cellular requirements for metabolites. (See, e.g., Hartman, S.C. and R.C. Mulligan (1988) Proc. Natl. Acad. Sci. USA 85:8047-8051.) Visible markers, e.g., anthocyanins, green fluorescent proteins (GFP; Clontech), β glucuronidase and its substrate β -glucuronide, or luciferase and its substrate luciferin may be used. These markers can be used not only to identify transformants, but also to quantify the amount of transient or stable protein expression attributable to a specific vector system. (See, e.g., Rhodes, C.A. (1995) Methods Mol. Biol. 55:121-131.)

Although the presence/absence of marker gene expression suggests that the gene of interest is also present, the presence and expression of the gene may need to be confirmed. For example, if the sequence encoding PRTS is inserted within a marker gene sequence, transformed cells containing sequences encoding PRTS can be identified by the absence of marker gene function. Alternatively, a marker gene can be placed in tandem with a sequence encoding PRTS under the control of a single promoter. Expression of the marker gene in response to induction or selection usually indicates expression of the tandem gene as well.

In general, host cells that contain the nucleic acid sequence encoding PRTS and that express PRTS may be identified by a variety of procedures known to those of skill in the art. These procedures include, but are not limited to, DNA-DNA or DNA-RNA hybridizations, PCR amplification, and protein bioassay or immunoassay techniques which include membrane, solution, or chip based technologies for the detection and/or quantification of nucleic acid or protein sequences.

Immunological methods for detecting and measuring the expression of PRTS using either specific polyclonal or monoclonal antibodies are known in the art. Examples of such techniques include enzyme-linked immunosorbent assays (ELISAs), radioimmunoassays (RIAs), and fluorescence activated cell sorting (FACS). A two-site, monoclonal-based immunoassay utilizing monoclonal

antibodies reactive to two non-interfering epitopes on PRTS is preferred, but a competitive binding assay may be employed. These and other assays are well known in the art. (See, e.g., Hampton, R. et al. (1990) Serological Methods, a Laboratory Manual, APS Press, St. Paul MN, Sect. IV; Coligan, J.E. et al. (1997) Current Protocols in Immunology, Greene Pub. Associates and Wiley-Interscience, New York NY; and Pound, J.D. (1998) Immunochemical Protocols, Humana Press, Totowa NJ.)

A wide variety of labels and conjugation techniques are known by those skilled in the art and may be used in various nucleic acid and amino acid assays. Means for producing labeled hybridization or PCR probes for detecting sequences related to polynucleotides encoding PRTS include oligolabeling, nick translation, end-labeling, or PCR amplification using a labeled nucleotide. Alternatively, the sequences encoding PRTS, or any fragments thereof, may be cloned into a vector for the production of an mRNA probe. Such vectors are known in the art, are commercially available, and may be used to synthesize RNA probes in vitro by addition of an appropriate RNA polymerase such as T7, T3, or SP6 and labeled nucleotides. These procedures may be conducted using a variety of commercially available kits, such as those provided by Amersham Pharmacia Biotech, Promega (Madison WI), and US Biochemical. Suitable reporter molecules or labels which may be used for ease of detection include radionuclides, enzymes, fluorescent, chemiluminescent, or chromogenic agents, as well as substrates, cofactors, inhibitors, magnetic particles, and the like.

Host cells transformed with nucleotide sequences encoding PRTS may be cultured under conditions suitable for the expression and recovery of the protein from cell culture. The protein produced by a transformed cell may be secreted or retained intracellularly depending on the sequence and/or the vector used. As will be understood by those of skill in the art, expression vectors containing polynucleotides which encode PRTS may be designed to contain signal sequences which direct secretion of PRTS through a prokaryotic or eukaryotic cell membrane.

In addition, a host cell strain may be chosen for its ability to modulate expression of the inserted sequences or to process the expressed protein in the desired fashion. Such modifications of the polypeptide include, but are not limited to, acetylation, carboxylation, glycosylation, phosphorylation, lipidation, and acylation. Post-translational processing which cleaves a "prepro" or "pro" form of the protein may also be used to specify protein targeting, folding, and/or activity. Different host cells which have specific cellular machinery and characteristic mechanisms for post-translational activities (e.g., CHO, HeLa, MDCK, HEK293, and WI38) are available from the American Type Culture Collection (ATCC, Manassas VA) and may be chosen to ensure the correct modification and processing of the foreign protein.

In another embodiment of the invention, natural, modified, or recombinant nucleic acid sequences encoding PRTS may be ligated to a heterologous sequence resulting in translation of a fusion

protein in any of the aforementioned host systems. For example, a chimeric PRTS protein containing a heterologous moiety that can be recognized by a commercially available antibody may facilitate the screening of peptide libraries for inhibitors of PRTS activity. Heterologous protein and peptide moieties may also facilitate purification of fusion proteins using commercially available affinity matrices. Such moieties include, but are not limited to, glutathione S-transferase (GST), maltose binding protein (MBP), thioredoxin (Trx), calmodulin binding peptide (CBP), 6-His, FLAG, *c-myc*, and hemagglutinin (HA). GST, MBP, Trx, CBP, and 6-His enable purification of their cognate fusion proteins on immobilized glutathione, maltose, phenylarsine oxide, calmodulin, and metal-chelate resins, respectively. FLAG, *c-myc*, and hemagglutinin (HA) enable immunoaffinity purification of fusion proteins using commercially available monoclonal and polyclonal antibodies that specifically recognize these epitope tags. A fusion protein may also be engineered to contain a proteolytic cleavage site located between the PRTS encoding sequence and the heterologous protein sequence, so that PRTS may be cleaved away from the heterologous moiety following purification. Methods for fusion protein expression and purification are discussed in Ausubel (1995, supra, ch. 10). A variety of commercially available kits may also be used to facilitate expression and purification of fusion proteins.

In a further embodiment of the invention, synthesis of radiolabeled PRTS may be achieved in vitro using the TNT rabbit reticulocyte lysate or wheat germ extract system (Promega). These systems couple transcription and translation of protein-coding sequences operably associated with the T7, T3, or SP6 promoters. Translation takes place in the presence of a radiolabeled amino acid precursor, for example, ³⁵S-methionine.

PRTS of the present invention or fragments thereof may be used to screen for compounds that specifically bind to PRTS. At least one and up to a plurality of test compounds may be screened for specific binding to PRTS. Examples of test compounds include antibodies, oligonucleotides, proteins (e.g., receptors), or small molecules.

In one embodiment, the compound thus identified is closely related to the natural ligand of PRTS, e.g., a ligand or fragment thereof, a natural substrate, a structural or functional mimetic, or a natural binding partner. (See, e.g., Coligan, J.E. et al. (1991) Current Protocols in Immunology 1(2): Chapter 5.) Similarly, the compound can be closely related to the natural receptor to which PRTS binds, or to at least a fragment of the receptor, e.g., the ligand binding site. In either case, the compound can be rationally designed using known techniques. In one embodiment, screening for these compounds involves producing appropriate cells which express PRTS, either as a secreted protein or on the cell membrane. Preferred cells include cells from mammals, yeast, Drosophila, or E. coli. Cells expressing PRTS or cell membrane fractions which contain PRTS are then contacted with

a test compound and binding, stimulation, or inhibition of activity of either PRTS or the compound is analyzed.

An assay may simply test binding of a test compound to the polypeptide, wherein binding is detected by a fluorophore, radioisotope, enzyme conjugate, or other detectable label. For example, the assay may comprise the steps of combining at least one test compound with PRTS, either in solution or affixed to a solid support, and detecting the binding of PRTS to the compound. Alternatively, the assay may detect or measure binding of a test compound in the presence of a labeled competitor. Additionally, the assay may be carried out using cell-free preparations, chemical libraries, or natural product mixtures, and the test compound(s) may be free in solution or affixed to a solid support.

PRTS of the present invention or fragments thereof may be used to screen for compounds that modulate the activity of PRTS. Such compounds may include agonists, antagonists, or partial or inverse agonists. In one embodiment, an assay is performed under conditions permissive for PRTS activity, wherein PRTS is combined with at least one test compound, and the activity of PRTS in the presence of a test compound is compared with the activity of PRTS in the absence of the test compound. A change in the activity of PRTS in the presence of the test compound is indicative of a compound that modulates the activity of PRTS. Alternatively, a test compound is combined with an *in vitro* or cell-free system comprising PRTS under conditions suitable for PRTS activity, and the assay is performed. In either of these assays, a test compound which modulates the activity of PRTS may do so indirectly and need not come in direct contact with the test compound. At least one and up to a plurality of test compounds may be screened.

In another embodiment, polynucleotides encoding PRTS or their mammalian homologs may be "knocked out" in an animal model system using homologous recombination in embryonic stem (ES) cells. Such techniques are well known in the art and are useful for the generation of animal models of human disease. (See, e.g., U.S. Patent Number 5,175,383 and U.S. Patent Number 5,767,337.) For example, mouse ES cells, such as the mouse 129/SvJ cell line, are derived from the early mouse embryo and grown in culture. The ES cells are transformed with a vector containing the gene of interest disrupted by a marker gene, e.g., the neomycin phosphotransferase gene (neo; Capecchi, M.R. (1989) Science 244:1288-1292). The vector integrates into the corresponding region of the host genome by homologous recombination. Alternatively, homologous recombination takes place using the Cre-loxP system to knockout a gene of interest in a tissue- or developmental stage-specific manner (Marth, J.D. (1996) Clin. Invest. 97:1999-2002; Wagner, K.U. et al. (1997) Nucleic Acids Res. 25:4323-4330). Transformed ES cells are identified and microinjected into mouse cell blastocysts such as those from the C57BL/6 mouse strain. The blastocysts are surgically transferred to pseudopregnant dams, and the

resulting chimeric progeny are genotyped and bred to produce heterozygous or homozygous strains. Transgenic animals thus generated may be tested with potential therapeutic or toxic agents.

Polynucleotides encoding PRTS may also be manipulated in vitro in ES cells derived from human blastocysts. Human ES cells have the potential to differentiate into at least eight separate cell lineages including endoderm, mesoderm, and ectodermal cell types. These cell lineages differentiate into, for example, neural cells, hematopoietic lineages, and cardiomyocytes (Thomson, J.A. et al. (1998) Science 282:1145-1147).

Polynucleotides encoding PRTS can also be used to create "knockin" humanized animals (pigs) or transgenic animals (mice or rats) to model human disease. With knockin technology, a region of a polynucleotide encoding PRTS is injected into animal ES cells, and the injected sequence integrates into the animal cell genome. Transformed cells are injected into blastulae, and the blastulae are implanted as described above. Transgenic progeny or inbred lines are studied and treated with potential pharmaceutical agents to obtain information on treatment of a human disease. Alternatively, a mammal inbred to overexpress PRTS, e.g., by secreting PRTS in its milk, may also serve as a convenient source of that protein (Janne, J. et al. (1998) Biotechnol. Annu. Rev. 4:55-74).

THERAPEUTICS

PRTS are useful for hydrolyzing peptide bonds. Chemical and structural similarity, e.g., in the context of sequences and motifs, exists between regions of PRTS and proteases. In addition, the expression of PRTS is closely associated with hemic, neurological, reproductive, endocrine, urogenital, diseased, teratocarcinoma, and tumorous tissues. Therefore, PRTS appears to play a role in gastrointestinal, cardiovascular, autoimmune/inflammatory, cell proliferative, developmental, epithelial, neurological, and reproductive disorders. In the treatment of disorders associated with increased PRTS expression or activity, it is desirable to decrease the expression or activity of PRTS. In the treatment of disorders associated with decreased PRTS expression or activity, it is desirable to increase the expression or activity of PRTS.

Therefore, in one embodiment, PRTS or a fragment or derivative thereof may be administered to a subject to treat or prevent a disorder associated with decreased expression or activity of PRTS. Examples of such disorders include, but are not limited to, a gastrointestinal disorder, such as dysphagia, peptic esophagitis, esophageal spasm, esophageal stricture, esophageal carcinoma, dyspepsia, indigestion, gastritis, gastric carcinoma, anorexia, nausea, emesis, gastroparesis, antral or pyloric edema, abdominal angina, pyrosis, gastroenteritis, intestinal obstruction, infections of the intestinal tract, peptic ulcer, cholelithiasis, cholecystitis, cholestasis, pancreatitis, pancreatic carcinoma, biliary tract disease, hepatitis, hyperbilirubinemia, cirrhosis, passive congestion of the liver, hepatoma, infectious colitis, ulcerative colitis, ulcerative proctitis, Crohn's disease, Whipple's disease, Mallory-

Weiss syndrome, colonic carcinoma, colonic obstruction, irritable bowel syndrome, short bowel syndrome, diarrhea, constipation, gastrointestinal hemorrhage, acquired immunodeficiency syndrome (AIDS) enteropathy, jaundice, hepatic encephalopathy, hepatorenal syndrome, hepatic steatosis, hemochromatosis, Wilson's disease, alpha₁-antitrypsin deficiency, Reye's syndrome, primary sclerosing

5 cholangitis, liver infarction, portal vein obstruction and thrombosis, centrilobular necrosis, peliosis hepatis, hepatic vein thrombosis, veno-occlusive disease, preeclampsia, eclampsia, acute fatty liver of pregnancy, intrahepatic cholestasis of pregnancy, and hepatic tumors including nodular hyperplasias, adenomas, and carcinomas; a cardiovascular disorder, such as arteriovenous fistula, atherosclerosis, hypertension, vasculitis, Raynaud's disease, aneurysms, arterial dissections, varicose veins,

10 thrombophlebitis and phlebothrombosis, vascular tumors, and complications of thrombolysis, balloon angioplasty, vascular replacement, and coronary artery bypass graft surgery, congestive heart failure, ischemic heart disease, angina pectoris, myocardial infarction, hypertensive heart disease, degenerative valvular heart disease, calcific aortic valve stenosis, congenitally bicuspid aortic valve, mitral annular calcification, mitral valve prolapse, rheumatic fever and rheumatic heart disease, infective endocarditis,

15 nonbacterial thrombotic endocarditis, endocarditis of systemic lupus erythematosus, carcinoid heart disease, cardiomyopathy, myocarditis, pericarditis, neoplastic heart disease, congenital heart disease, and complications of cardiac transplantation; an autoimmune/inflammatory disorder, such as acquired immunodeficiency syndrome (AIDS), Addison's disease, adult respiratory distress syndrome, allergies, ankylosing spondylitis, amyloidosis, anemia, asthma, atherosclerosis, atherosclerotic plaque rupture,

20 autoimmune hemolytic anemia, autoimmune thyroiditis, autoimmune polyendocrinopathy-candidiasis-ectodermal dystrophy (APECED), bronchitis, cholecystitis, contact dermatitis, Crohn's disease, atopic dermatitis, dermatomyositis, diabetes mellitus, emphysema, episodic lymphopenia with lymphocytotoxins, erythroblastosis fetalis, erythema nodosum, atrophic gastritis, glomerulonephritis, Goodpasture's syndrome, gout, Graves' disease, Hashimoto's thyroiditis, hypereosinophilia, irritable

25 bowel syndrome, multiple sclerosis, myasthenia gravis, myocardial or pericardial inflammation, osteoarthritis, degradation of articular cartilage, osteoporosis, pancreatitis, polymyositis, psoriasis, Reiter's syndrome, rheumatoid arthritis, scleroderma, Sjögren's syndrome, systemic anaphylaxis, systemic lupus erythematosus, systemic sclerosis, thrombocytopenic purpura, ulcerative colitis, uveitis, Werner syndrome, complications of cancer, hemodialysis, and extracorporeal circulation, viral,

30 bacterial, fungal, parasitic, protozoal, and helminthic infections, and trauma; a cell proliferative disorder such as actinic keratosis, arteriosclerosis, atherosclerosis, bursitis, cirrhosis, hepatitis, mixed connective tissue disease (MCTD), myelofibrosis, paroxysmal nocturnal hemoglobinuria, polycythemia vera, psoriasis, primary thrombocythemia, and cancers including adenocarcinoma, leukemia, lymphoma, melanoma, myeloma, sarcoma, teratocarcinoma, and, in particular, cancers of the adrenal

- gland, bladder, bone, bone marrow, brain, breast, cervix, gall bladder, ganglia, gastrointestinal tract, heart, kidney, liver, lung, muscle, ovary, pancreas, parathyroid, penis, prostate, salivary glands, skin, spleen, testis, thymus, thyroid, and uterus; a developmental disorder, such as renal tubular acidosis, anemia, Cushing's syndrome, achondroplastic dwarfism, Duchenne and Becker muscular dystrophy,
- 5 bone resorption, epilepsy, gonadal dysgenesis, WAGR syndrome (Wilms' tumor, aniridia, genitourinary abnormalities, and mental retardation), Smith-Magenis syndrome, myelodysplastic syndrome, hereditary mucoepithelial dysplasia, hereditary keratodermas, hereditary neuropathies such as Charcot-Marie-Tooth disease and neurofibromatosis, hypothyroidism, hydrocephalus, seizure disorders such as Sydenham's chorea and cerebral palsy, spina bifida, anencephaly, craniorachischisis, congenital
- 10 glaucoma, cataract, age-related macular degeneration, and sensorineural hearing loss; an epithelial disorder, such as dyshidrotic eczema, allergic contact dermatitis, keratosis pilaris, melasma, vitiligo, actinic keratosis, basal cell carcinoma, squamous cell carcinoma, seborrheic keratosis, folliculitis, herpes simplex, herpes zoster, varicella, candidiasis, dermatophytosis, scabies, insect bites, cherry angioma, keloid, dermatofibroma, acrochordons, urticaria, transient acantholytic dermatosis, xerosis,
- 15 eczema, atopic dermatitis, contact dermatitis, hand eczema, nummular eczema, lichen simplex chronicus, asteatotic eczema, stasis dermatitis and stasis ulceration, seborrheic dermatitis, psoriasis, lichen planus, pityriasis rosea, impetigo, ecthyma, dermatophytosis, tinea versicolor, warts, acne vulgaris, acne rosacea, pemphigus vulgaris, pemphigus foliaceus, paraneoplastic pemphigus, bullous pemphigoid, herpes gestationis, dermatitis herpetiformis, linear IgA disease, epidermolysis bullosa
- 20 acquisita, dermatomyositis, lupus erythematosus, scleroderma and morphea, erythroderma, alopecia, figurate skin lesions, telangiectasias, hypopigmentation, hyperpigmentation, vesicles/bullae, exanthems, cutaneous drug reactions, papulonodular skin lesions, chronic non-healing wounds, photosensitivity diseases, epidermolysis bullosa simplex, epidermolytic hyperkeratosis, epidermolytic and nonepidermolytic palmoplantar keratoderma, ichthyosis bullosa of Siemens, ichthyosis exfoliativa,
- 25 keratosis palmaris et plantaris, keratosis palmoplantaris, palmoplantar keratoderma, keratosis punctata, Meesmann's corneal dystrophy, pachyonychia congenita, white sponge nevus, steatocystoma multiplex, epidermal nevi/epidermolytic hyperkeratosis type, monilethrix, trichothiodystrophy, chronic hepatitis/cryptogenic cirrhosis, and colorectal hyperplasia; a neurological disorder, such as epilepsy, ischemic cerebrovascular disease, stroke, cerebral neoplasms, Alzheimer's disease, Pick's disease,
- 30 Huntington's disease, dementia, Parkinson's disease and other extrapyramidal disorders, amyotrophic lateral sclerosis and other motor neuron disorders, progressive neural muscular atrophy, retinitis pigmentosa, hereditary ataxias, multiple sclerosis and other demyelinating diseases, bacterial and viral meningitis, brain abscess, subdural empyema, epidural abscess, suppurative intracranial thrombophlebitis, myelitis and radiculitis, viral central nervous system disease, prion diseases including

kuru, Creutzfeldt-Jakob disease, and Gerstmann-Straussler-Scheinker syndrome, fatal familial insomnia, nutritional and metabolic diseases of the nervous system, neurofibromatosis, tuberous sclerosis, cerebelloretinal hemangioblastomatosis, encephalotrigeminal syndrome, mental retardation and other developmental disorders of the central nervous system including Down syndrome, cerebral palsy, neuroskeletal disorders, autonomic nervous system disorders, cranial nerve disorders, spinal cord diseases, muscular dystrophy and other neuromuscular disorders, peripheral nervous system disorders, dermatomyositis and polymyositis, inherited, metabolic, endocrine, and toxic myopathies; myasthenia gravis, periodic paralysis, mental disorders including mood, anxiety, and schizophrenic disorders, seasonal affective disorder (SAD), akathisia, amnesia, catatonia, diabetic neuropathy, tardive dyskinesia, dystonias, paranoid psychoses, postherpetic neuralgia, Tourette's disorder, progressive supranuclear palsy, corticobasal degeneration, and familial frontotemporal dementia; and a reproductive disorder, such as infertility, including tubal disease, ovulatory defects, and endometriosis, a disorder of prolactin production, a disruption of the estrous cycle, a disruption of the menstrual cycle, polycystic ovary syndrome, ovarian hyperstimulation syndrome, an endometrial or ovarian tumor, a uterine fibroid, autoimmune disorders, an ectopic pregnancy, and teratogenesis; cancer of the breast, fibrocystic breast disease, and galactorrhea; a disruption of spermatogenesis, abnormal sperm physiology, cancer of the testis, cancer of the prostate, benign prostatic hyperplasia, prostatitis, Peyronie's disease, impotence, carcinoma of the male breast, and gynecomastia.

In another embodiment, a vector capable of expressing PRTS or a fragment or derivative thereof may be administered to a subject to treat or prevent a disorder associated with decreased expression or activity of PRTS including, but not limited to, those described above.

In a further embodiment, a composition comprising a substantially purified PRTS in conjunction with a suitable pharmaceutical carrier may be administered to a subject to treat or prevent a disorder associated with decreased expression or activity of PRTS including, but not limited to, those provided above.

In still another embodiment, an agonist which modulates the activity of PRTS may be administered to a subject to treat or prevent a disorder associated with decreased expression or activity of PRTS including, but not limited to, those listed above.

In a further embodiment, an antagonist of PRTS may be administered to a subject to treat or prevent a disorder associated with increased expression or activity of PRTS. Examples of such disorders include, but are not limited to, those gastrointestinal, cardiovascular, autoimmune/inflammatory, cell proliferative, developmental, epithelial, neurological, and reproductive disorders described above. In one aspect, an antibody which specifically binds PRTS may be used

directly as an antagonist or indirectly as a targeting or delivery mechanism for bringing a pharmaceutical agent to cells or tissues which express PRTS.

In an additional embodiment, a vector expressing the complement of the polynucleotide encoding PRTS may be administered to a subject to treat or prevent a disorder associated with
5 increased expression or activity of PRTS including, but not limited to, those described above.

In other embodiments, any of the proteins, antagonists, antibodies, agonists, complementary sequences, or vectors of the invention may be administered in combination with other appropriate therapeutic agents. Selection of the appropriate agents for use in combination therapy may be made by one of ordinary skill in the art, according to conventional pharmaceutical principles. The combination
10 of therapeutic agents may act synergistically to effect the treatment or prevention of the various disorders described above. Using this approach, one may be able to achieve therapeutic efficacy with lower dosages of each agent, thus reducing the potential for adverse side effects.

An antagonist of PRTS may be produced using methods which are generally known in the art. In particular, purified PRTS may be used to produce antibodies or to screen libraries of pharmaceutical
15 agents to identify those which specifically bind PRTS. Antibodies to PRTS may also be generated using methods that are well known in the art. Such antibodies may include, but are not limited to, polyclonal, monoclonal, chimeric, and single chain antibodies, Fab fragments, and fragments produced by a Fab expression library. Neutralizing antibodies (i.e., those which inhibit dimer formation) are generally preferred for therapeutic use.

20 For the production of antibodies, various hosts including goats, rabbits, rats, mice, humans, and others may be immunized by injection with PRTS or with any fragment or oligopeptide thereof which has immunogenic properties. Depending on the host species, various adjuvants may be used to increase immunological response. Such adjuvants include, but are not limited to, Freund's, mineral gels such as aluminum hydroxide, and surface active substances such as lysolecithin, pluronic polyols,
25 polyanions, peptides, oil emulsions, KLH, and dinitrophenol. Among adjuvants used in humans, BCG (bacilli Calmette-Guerin) and Corynebacterium parvum are especially preferable.

It is preferred that the oligopeptides, peptides, or fragments used to induce antibodies to PRTS have an amino acid sequence consisting of at least about 5 amino acids, and generally will consist of at least about 10 amino acids. It is also preferable that these oligopeptides, peptides, or fragments are
30 identical to a portion of the amino acid sequence of the natural protein. Short stretches of PRTS amino acids may be fused with those of another protein, such as KLH, and antibodies to the chimeric molecule may be produced.

Monoclonal antibodies to PRTS may be prepared using any technique which provides for the production of antibody molecules by continuous cell lines in culture. These include, but are not limited

to, the hybridoma technique, the human B-cell hybridoma technique, and the EBV-hybridoma technique. (See, e.g., Kohler, G. et al. (1975) *Nature* 256:495-497; Kozbor, D. et al. (1985) *J. Immunol. Methods* 81:31-42; Cote, R.J. et al. (1983) *Proc. Natl. Acad. Sci. USA* 80:2026-2030; and Cole, S.P. et al. (1984) *Mol. Cell Biol.* 62:109-120.)

5 In addition, techniques developed for the production of "chimeric antibodies," such as the splicing of mouse antibody genes to human antibody genes to obtain a molecule with appropriate antigen specificity and biological activity, can be used. (See, e.g., Morrison, S.L. et al. (1984) *Proc. Natl. Acad. Sci. USA* 81:6851-6855; Neuberger, M.S. et al. (1984) *Nature* 312:604-608; and Takeda, S. et al. (1985) *Nature* 314:452-454.) Alternatively, techniques described for the production of single
10 chain antibodies may be adapted, using methods known in the art, to produce PRTS-specific single chain antibodies. Antibodies with related specificity, but of distinct idiotypic composition, may be generated by chain shuffling from random combinatorial immunoglobulin libraries. (See, e.g., Burton, D.R. (1991) *Proc. Natl. Acad. Sci. USA* 88:10134-10137.)

Antibodies may also be produced by inducing in vivo production in the lymphocyte population
15 or by screening immunoglobulin libraries or panels of highly specific binding reagents as disclosed in the literature. (See, e.g., Orlandi, R. et al. (1989) *Proc. Natl. Acad. Sci. USA* 86:3833-3837; Winter, G. et al. (1991) *Nature* 349:293-299.)

Antibody fragments which contain specific binding sites for PRTS may also be generated. For example, such fragments include, but are not limited to, $F(ab')_2$ fragments produced by pepsin digestion
20 of the antibody molecule and Fab fragments generated by reducing the disulfide bridges of the $F(ab')_2$ fragments. Alternatively, Fab expression libraries may be constructed to allow rapid and easy identification of monoclonal Fab fragments with the desired specificity. (See, e.g., Huse, W.D. et al. (1989) *Science* 246:1275-1281.)

Various immunoassays may be used for screening to identify antibodies having the desired
25 specificity. Numerous protocols for competitive binding or immunoradiometric assays using either polyclonal or monoclonal antibodies with established specificities are well known in the art. Such immunoassays typically involve the measurement of complex formation between PRTS and its specific antibody. A two-site, monoclonal-based immunoassay utilizing monoclonal antibodies reactive to two non-interfering PRTS epitopes is generally used, but a competitive binding assay may also be employed
30 (Pound, supra).

Various methods such as Scatchard analysis in conjunction with radioimmunoassay techniques may be used to assess the affinity of antibodies for PRTS. Affinity is expressed as an association constant, K_a , which is defined as the molar concentration of PRTS-antibody complex divided by the molar concentrations of free antigen and free antibody under equilibrium conditions. The K_a determined

for a preparation of polyclonal antibodies, which are heterogeneous in their affinities for multiple PRTS epitopes, represents the average affinity, or avidity, of the antibodies for PRTS. The K_a determined for a preparation of monoclonal antibodies, which are monospecific for a particular PRTS epitope, represents a true measure of affinity. High-affinity antibody preparations with K_a ranging from about 10^9 to 10^{12} L/mole are preferred for use in immunoassays in which the PRTS-antibody complex must withstand rigorous manipulations. Low-affinity antibody preparations with K_a ranging from about 10^6 to 10^7 L/mole are preferred for use in immunopurification and similar procedures which ultimately require dissociation of PRTS, preferably in active form, from the antibody (Catty, D. (1988) Antibodies, Volume I: A Practical Approach, IRL Press, Washington DC; Liddell, J.E. and A. Cryer (1991) A Practical Guide to Monoclonal Antibodies, John Wiley & Sons, New York NY).

The titer and avidity of polyclonal antibody preparations may be further evaluated to determine the quality and suitability of such preparations for certain downstream applications. For example, a polyclonal antibody preparation containing at least 1-2 mg specific antibody/ml, preferably 5-10 mg specific antibody/ml, is generally employed in procedures requiring precipitation of PRTS-antibody complexes. Procedures for evaluating antibody specificity, titer, and avidity, and guidelines for antibody quality and usage in various applications, are generally available. (See, e.g., Catty, *supra*, and Coligan et al. *supra*.)

In another embodiment of the invention, the polynucleotides encoding PRTS, or any fragment or complement thereof, may be used for therapeutic purposes. In one aspect, modifications of gene expression can be achieved by designing complementary sequences or antisense molecules (DNA, RNA, PNA, or modified oligonucleotides) to the coding or regulatory regions of the gene encoding PRTS. Such technology is well known in the art, and antisense oligonucleotides or larger fragments can be designed from various locations along the coding or control regions of sequences encoding PRTS. (See, e.g., Agrawal, S., ed. (1996) Antisense Therapeutics, Humana Press Inc., Totawa NJ.)

In therapeutic use, any gene delivery system suitable for introduction of the antisense sequences into appropriate target cells can be used. Antisense sequences can be delivered intracellularly in the form of an expression plasmid which, upon transcription, produces a sequence complementary to at least a portion of the cellular sequence encoding the target protein. (See, e.g., Slater, J.E. et al. (1998) *J. Allergy Clin. Immunol.* 102(3):469-475; and Scanlon, K.J. et al. (1995) 9(13):1288-1296.) Antisense sequences can also be introduced intracellularly through the use of viral vectors, such as retrovirus and adeno-associated virus vectors. (See, e.g., Miller, A.D. (1990) *Blood* 76:271; Ausubel, *supra*; Uckert, W. and W. Walther (1994) *Pharmacol. Ther.* 63(3):323-347.) Other gene delivery mechanisms include liposome-derived systems, artificial viral envelopes, and other systems known in the art. (See, e.g., Rossi, J.J. (1995) *Br. Med. Bull.* 51(1):217-225; Boado, R.J. et

al. (1998) J. Pharm. Sci. 87(11):1308-1315; and Morris, M.C. et al. (1997) Nucleic Acids Res. 25(14):2730-2736.)

In another embodiment of the invention, polynucleotides encoding PRTS may be used for somatic or germline gene therapy. Gene therapy may be performed to (i) correct a genetic deficiency (e.g., in the cases of severe combined immunodeficiency (SCID)-X1 disease characterized by X-linked inheritance (Cavazzana-Calvo, M. et al. (2000) Science 288:669-672), severe combined immunodeficiency syndrome associated with an inherited adenosine deaminase (ADA) deficiency (Blaese, R.M. et al. (1995) Science 270:475-480; Bordignon, C. et al. (1995) Science 270:470-475), cystic fibrosis (Zabner, J. et al. (1993) Cell 75:207-216; Crystal, R.G. et al. (1995) Hum. Gene Therapy 6:643-666; Crystal, R.G. et al. (1995) Hum. Gene Therapy 6:667-703), thalassemias, familial hypercholesterolemia, and hemophilia resulting from Factor VIII or Factor IX deficiencies (Crystal, R.G. (1995) Science 270:404-410; Verma, I.M. and N. Somia (1997) Nature 389:239-242)), (ii) express a conditionally lethal gene product (e.g., in the case of cancers which result from unregulated cell proliferation), or (iii) express a protein which affords protection against intracellular parasites (e.g., against human retroviruses, such as human immunodeficiency virus (HIV) (Baltimore, D. (1988) Nature 335:395-396; Poeschla, E. et al. (1996) Proc. Natl. Acad. Sci. USA. 93:11395-11399), hepatitis B or C virus (HBV, HCV); fungal parasites, such as Candida albicans and Paracoccidioides brasiliensis; and protozoan parasites such as Plasmodium falciparum and Trypanosoma cruzi). In the case where a genetic deficiency in PRTS expression or regulation causes disease, the expression of PRTS from an appropriate population of transduced cells may alleviate the clinical manifestations caused by the genetic deficiency.

In a further embodiment of the invention, diseases or disorders caused by deficiencies in PRTS are treated by constructing mammalian expression vectors encoding PRTS and introducing these vectors by mechanical means into PRTS-deficient cells. Mechanical transfer technologies for use with cells in vivo or ex vitro include (i) direct DNA microinjection into individual cells, (ii) ballistic gold particle delivery, (iii) liposome-mediated transfection, (iv) receptor-mediated gene transfer, and (v) the use of DNA transposons (Morgan, R.A. and W.F. Anderson (1993) Annu. Rev. Biochem. 62:191-217; Ivics, Z. (1997) Cell 91:501-510; Boulay, J-L. and H. Récipon (1998) Curr. Opin. Biotechnol. 9:445-450).

Expression vectors that may be effective for the expression of PRTS include, but are not limited to, the PCDNA 3.1, EPITAG, PRCCMV2, PREP, PVAX vectors (Invitrogen, Carlsbad CA), PCMV-SCRIPT, PCMV-TAG, PEGSH/PERV (Stratagene, La Jolla CA), and PTET-OFF, PTET-ON, PTRE2, PTRE2-LUC, PTK-HYG (Clontech, Palo Alto CA). PRTS may be expressed using (i) a constitutively active promoter, (e.g., from cytomegalovirus (CMV), Rous sarcoma virus

(RSV), SV40 virus, thymidine kinase (TK), or β -actin genes), (ii) an inducible promoter (e.g., the tetracycline-regulated promoter (Gossen, M. and H. Bujard (1992) *Proc. Natl. Acad. Sci. USA* 89:5547-5551; Gossen, M. et al. (1995) *Science* 268:1766-1769; Rossi, F.M.V. and H.M. Blau (1998) *Curr. Opin. Biotechnol.* 9:451-456), commercially available in the T-REX plasmid (Invitrogen)); the
5 ecdysone-inducible promoter (available in the plasmids PVGRXR and PIND; Invitrogen); the FK506/rapamycin inducible promoter; or the RU486/mifepristone inducible promoter (Rossi, F.M.V. and Blau, H.M. *supra*)), or (iii) a tissue-specific promoter or the native promoter of the endogenous gene encoding PRTS from a normal individual.

Commercially available liposome transformation kits (e.g., the PERFECT LIPID
10 TRANSFECTION KIT, available from Invitrogen) allow one with ordinary skill in the art to deliver polynucleotides to target cells in culture and require minimal effort to optimize experimental parameters. In the alternative, transformation is performed using the calcium phosphate method (Graham, F.L. and A.J. Eb (1973) *Virology* 52:456-467), or by electroporation (Neumann, E. et al. (1982) *EMBO J.* 1:841-845). The introduction of DNA to primary cells requires modification of these
15 standardized mammalian transfection protocols.

In another embodiment of the invention, diseases or disorders caused by genetic defects with respect to PRTS expression are treated by constructing a retrovirus vector consisting of (i) the polynucleotide encoding PRTS under the control of an independent promoter or the retrovirus long terminal repeat (LTR) promoter, (ii) appropriate RNA packaging signals, and (iii) a Rev-responsive
20 element (RRE) along with additional retrovirus *cis*-acting RNA sequences and coding sequences required for efficient vector propagation. Retrovirus vectors (e.g., PFB and PFBNEO) are commercially available (Stratagene) and are based on published data (Riviere, I. et al. (1995) *Proc. Natl. Acad. Sci. USA* 92:6733-6737), incorporated by reference herein. The vector is propagated in an appropriate vector producing cell line (VPCL) that expresses an envelope gene with a tropism for
25 receptors on the target cells or a promiscuous envelope protein such as VSVg (Armentano, D. et al. (1987) *J. Virol.* 61:1647-1650; Bender, M.A. et al. (1987) *J. Virol.* 61:1639-1646; Adam, M.A. and A.D. Miller (1988) *J. Virol.* 62:3802-3806; Dull, T. et al. (1998) *J. Virol.* 72:8463-8471; Zufferey, R. et al. (1998) *J. Virol.* 72:9873-9880). U.S. Patent Number 5,910,434 to Rigg ("Method for obtaining retrovirus packaging cell lines producing high transducing efficiency retroviral supernatant") discloses a
30 method for obtaining retrovirus packaging cell lines and is hereby incorporated by reference. Propagation of retrovirus vectors, transduction of a population of cells (e.g., CD4⁺ T-cells), and the return of transduced cells to a patient are procedures well known to persons skilled in the art of gene therapy and have been well documented (Ranga, U. et al. (1997) *J. Virol.* 71:7020-7029; Bauer, G. et

al. (1997) Blood 89:2259-2267; Bonyhadi, M.L. (1997) J. Virol. 71:4707-4716; Ranga, U. et al. (1998) Proc. Natl. Acad. Sci. USA 95:1201-1206; Su, L. (1997) Blood 89:2283-2290).

In the alternative, an adenovirus-based gene therapy delivery system is used to deliver polynucleotides encoding PRTS to cells which have one or more genetic abnormalities with respect to the expression of PRTS. The construction and packaging of adenovirus-based vectors are well known to those with ordinary skill in the art. Replication defective adenovirus vectors have proven to be versatile for importing genes encoding immunoregulatory proteins into intact islets in the pancreas (Csete, M.E. et al. (1995) Transplantation 27:263-268). Potentially useful adenoviral vectors are described in U.S. Patent Number 5,707,618 to Armentano ("Adenovirus vectors for gene therapy"), hereby incorporated by reference. For adenoviral vectors, see also Antinozzi, P.A. et al. (1999) Annu. Rev. Nutr. 19:511-544 and Verma, I.M. and N. Somia (1997) Nature 18:389:239-242, both incorporated by reference herein.

In another alternative, a herpes-based, gene therapy delivery system is used to deliver polynucleotides encoding PRTS to target cells which have one or more genetic abnormalities with respect to the expression of PRTS. The use of herpes simplex virus (HSV)-based vectors may be especially valuable for introducing PRTS to cells of the central nervous system, for which HSV has a tropism. The construction and packaging of herpes-based vectors are well known to those with ordinary skill in the art. A replication-competent herpes simplex virus (HSV) type 1-based vector has been used to deliver a reporter gene to the eyes of primates (Liu, X. et al. (1999) Exp. Eye Res. 169:385-395). The construction of a HSV-1 virus vector has also been disclosed in detail in U.S. Patent Number 5,804,413 to DeLuca ("Herpes simplex virus strains for gene transfer"), which is hereby incorporated by reference. U.S. Patent Number 5,804,413 teaches the use of recombinant HSV d92 which consists of a genome containing at least one exogenous gene to be transferred to a cell under the control of the appropriate promoter for purposes including human gene therapy. Also taught by this patent are the construction and use of recombinant HSV strains deleted for ICP4, ICP27 and ICP22. For HSV vectors, see also Goins, W.F. et al. (1999) J. Virol. 73:519-532 and Xu, H. et al. (1994) Dev. Biol. 163:152-161, hereby incorporated by reference. The manipulation of cloned herpesvirus sequences, the generation of recombinant virus following the transfection of multiple plasmids containing different segments of the large herpesvirus genomes, the growth and propagation of herpesvirus, and the infection of cells with herpesvirus are techniques well known to those of ordinary skill in the art.

In another alternative, an alphavirus (positive, single-stranded RNA virus) vector is used to deliver polynucleotides encoding PRTS to target cells. The biology of the prototypic alphavirus, Semliki Forest Virus (SFV), has been studied extensively and gene transfer vectors have been based on

the SFV genome (Garoff, H. and K.-J. Li (1998) *Curr. Opin. Biotechnol.* 9:464-469). During alphavirus RNA replication, a subgenomic RNA is generated that normally encodes the viral capsid proteins. This subgenomic RNA replicates to higher levels than the full length genomic RNA, resulting in the overproduction of capsid proteins relative to the viral proteins with enzymatic activity (e.g.,
5 protease and polymerase). Similarly, inserting the coding sequence for PRTS into the alphavirus genome in place of the capsid-coding region results in the production of a large number of PRTS-coding RNAs and the synthesis of high levels of PRTS in vector transduced cells. While alphavirus infection is typically associated with cell lysis within a few days, the ability to establish a persistent infection in hamster normal kidney cells (BHK-21) with a variant of Sindbis virus (SIN) indicates that the lytic
10 replication of alphaviruses can be altered to suit the needs of the gene therapy application (Dryga, S.A. et al. (1997) *Virology* 228:74-83). The wide host range of alphaviruses will allow the introduction of PRTS into a variety of cell types. The specific transduction of a subset of cells in a population may require the sorting of cells prior to transduction. The methods of manipulating infectious cDNA clones of alphaviruses, performing alphavirus cDNA and RNA transfections, and performing alphavirus
15 infections, are well known to those with ordinary skill in the art.

Oligonucleotides derived from the transcription initiation site, e.g., between about positions -10 and +10 from the start site, may also be employed to inhibit gene expression. Similarly, inhibition can be achieved using triple helix base-pairing methodology. Triple helix pairing is useful because it causes inhibition of the ability of the double helix to open sufficiently for the binding of polymerases,
20 transcription factors, or regulatory molecules. Recent therapeutic advances using triplex DNA have been described in the literature. (See, e.g., Gee, J.E. et al. (1994) in Huber, B.E. and B.I. Carr, Molecular and Immunologic Approaches, Futura Publishing, Mt. Kisco NY, pp. 163-177.) A complementary sequence or antisense molecule may also be designed to block translation of mRNA by preventing the transcript from binding to ribosomes.

25 Ribozymes, enzymatic RNA molecules, may also be used to catalyze the specific cleavage of RNA. The mechanism of ribozyme action involves sequence-specific hybridization of the ribozyme molecule to complementary target RNA, followed by endonucleolytic cleavage. For example, engineered hammerhead motif ribozyme molecules may specifically and efficiently catalyze endonucleolytic cleavage of sequences encoding PRTS.

30 Specific ribozyme cleavage sites within any potential RNA target are initially identified by scanning the target molecule for ribozyme cleavage sites, including the following sequences: GUA, GUU, and GUC. Once identified, short RNA sequences of between 15 and 20 ribonucleotides, corresponding to the region of the target gene containing the cleavage site, may be evaluated for secondary structural features which may render the oligonucleotide inoperable. The suitability of

candidate targets may also be evaluated by testing accessibility to hybridization with complementary oligonucleotides using ribonuclease protection assays.

Complementary ribonucleic acid molecules and ribozymes of the invention may be prepared by any method known in the art for the synthesis of nucleic acid molecules. These include techniques for chemically synthesizing oligonucleotides such as solid phase phosphoramidite chemical synthesis. Alternatively, RNA molecules may be generated by in vitro and in vivo transcription of DNA sequences encoding PRTS. Such DNA sequences may be incorporated into a wide variety of vectors with suitable RNA polymerase promoters such as T7 or SP6. Alternatively, these cDNA constructs that synthesize complementary RNA, constitutively or inducibly, can be introduced into cell lines, cells, or tissues.

RNA molecules may be modified to increase intracellular stability and half-life. Possible modifications include, but are not limited to, the addition of flanking sequences at the 5' and/or 3' ends of the molecule, or the use of phosphorothioate or 2' O-methyl rather than phosphodiesterase linkages within the backbone of the molecule. This concept is inherent in the production of PNAs and can be extended in all of these molecules by the inclusion of nontraditional bases such as inosine, queosine, and wybutosine, as well as acetyl-, methyl-, thio-, and similarly modified forms of adenine, cytidine, guanine, thymine, and uridine which are not as easily recognized by endogenous endonucleases.

An additional embodiment of the invention encompasses a method for screening for a compound which is effective in altering expression of a polynucleotide encoding PRTS. Compounds which may be effective in altering expression of a specific polynucleotide may include, but are not limited to, oligonucleotides, antisense oligonucleotides, triple helix-forming oligonucleotides, transcription factors and other polypeptide transcriptional regulators, and non-macromolecular chemical entities which are capable of interacting with specific polynucleotide sequences. Effective compounds may alter polynucleotide expression by acting as either inhibitors or promoters of polynucleotide expression. Thus, in the treatment of disorders associated with increased PRTS expression or activity, a compound which specifically inhibits expression of the polynucleotide encoding PRTS may be therapeutically useful, and in the treatment of disorders associated with decreased PRTS expression or activity, a compound which specifically promotes expression of the polynucleotide encoding PRTS may be therapeutically useful.

At least one, and up to a plurality, of test compounds may be screened for effectiveness in altering expression of a specific polynucleotide. A test compound may be obtained by any method commonly known in the art, including chemical modification of a compound known to be effective in altering polynucleotide expression; selection from an existing, commercially-available or proprietary library of naturally-occurring or non-natural chemical compounds; rational design of a compound based on chemical and/or structural properties of the target polynucleotide; and selection from a

library of chemical compounds created combinatorially or randomly. A sample comprising a polynucleotide encoding PRTS is exposed to at least one test compound thus obtained. The sample may comprise, for example, an intact or permeabilized cell, or an in vitro cell-free or reconstituted biochemical system. Alterations in the expression of a polynucleotide encoding PRTS are assayed by
5 any method commonly known in the art. Typically, the expression of a specific nucleotide is detected by hybridization with a probe having a nucleotide sequence complementary to the sequence of the polynucleotide encoding PRTS. The amount of hybridization may be quantified, thus forming the basis for a comparison of the expression of the polynucleotide both with and without exposure to one or more test compounds. Detection of a change in the expression of a polynucleotide exposed to a
10 test compound indicates that the test compound is effective in altering the expression of the polynucleotide. A screen for a compound effective in altering expression of a specific polynucleotide can be carried out, for example, using a Schizosaccharomyces pombe gene expression system (Atkins, D. et al. (1999) U.S. Patent No. 5,932,435; Arndt, G.M. et al. (2000) Nucleic Acids Res. 28:E15) or a human cell line such as HeLa cell (Clarke, M.L. et al. (2000) Biochem. Biophys. Res. Commun.
15 268:8-13). A particular embodiment of the present invention involves screening a combinatorial library of oligonucleotides (such as deoxyribonucleotides, ribonucleotides, peptide nucleic acids, and modified oligonucleotides) for antisense activity against a specific polynucleotide sequence (Bruce, T.W. et al. (1997) U.S. Patent No. 5,686,242; Bruce, T.W. et al. (2000) U.S. Patent No. 6,022,691).

Many methods for introducing vectors into cells or tissues are available and equally suitable for
20 use in vivo, in vitro, and ex vivo. For ex vivo therapy, vectors may be introduced into stem cells taken from the patient and clonally propagated for autologous transplant back into that same patient. Delivery by transfection, by liposome injections, or by polycationic amino polymers may be achieved using methods which are well known in the art. (See, e.g., Goldman, C.K. et al. (1997) Nat. Biotechnol. 15:462-466.)

25 Any of the therapeutic methods described above may be applied to any subject in need of such therapy, including, for example, mammals such as humans, dogs, cats, cows, horses, rabbits, and monkeys.

An additional embodiment of the invention relates to the administration of a composition which generally comprises an active ingredient formulated with a pharmaceutically acceptable excipient.
30 Excipients may include, for example, sugars, starches, celluloses, gums, and proteins. Various formulations are commonly known and are thoroughly discussed in the latest edition of Remington's Pharmaceutical Sciences (Maack Publishing, Easton PA). Such compositions may consist of PRTS, antibodies to PRTS, and mimetics, agonists, antagonists, or inhibitors of PRTS.

The compositions utilized in this invention may be administered by any number of routes including, but not limited to, oral, intravenous, intramuscular, intra-arterial, intramedullary, intrathecal, intraventricular, pulmonary, transdermal, subcutaneous, intraperitoneal, intranasal, enteral, topical, sublingual, or rectal means.

5 Compositions for pulmonary administration may be prepared in liquid or dry powder form. These compositions are generally aerosolized immediately prior to inhalation by the patient. In the case of small molecules (e.g. traditional low molecular weight organic drugs), aerosol delivery of fast-acting formulations is well-known in the art. In the case of macromolecules (e.g. larger peptides and proteins), recent developments in the field of pulmonary delivery via the alveolar region of the lung have enabled
10 the practical delivery of drugs such as insulin to blood circulation (see, e.g., Patton, J.S. et al., U.S. Patent No. 5,997,848). Pulmonary delivery has the advantage of administration without needle injection, and obviates the need for potentially toxic penetration enhancers.

 Compositions suitable for use in the invention include compositions wherein the active ingredients are contained in an effective amount to achieve the intended purpose. The determination of
15 an effective dose is well within the capability of those skilled in the art.

 Specialized forms of compositions may be prepared for direct intracellular delivery of macromolecules comprising PRTS or fragments thereof. For example, liposome preparations containing a cell-impermeable macromolecule may promote cell fusion and intracellular delivery of the macromolecule. Alternatively, PRTS or a fragment thereof may be joined to a short cationic N-
20 terminal portion from the HIV Tat-1 protein. Fusion proteins thus generated have been found to transduce into the cells of all tissues, including the brain, in a mouse model system (Schwarze, S.R. et al. (1999) Science 285:1569-1572).

 For any compound, the therapeutically effective dose can be estimated initially either in cell culture assays, e.g., of neoplastic cells, or in animal models such as mice, rats, rabbits, dogs, monkeys,
25 or pigs. An animal model may also be used to determine the appropriate concentration range and route of administration. Such information can then be used to determine useful doses and routes for administration in humans.

 A therapeutically effective dose refers to that amount of active ingredient, for example PRTS or fragments thereof, antibodies of PRTS, and agonists, antagonists or inhibitors of PRTS, which
30 ameliorates the symptoms or condition. Therapeutic efficacy and toxicity may be determined by standard pharmaceutical procedures in cell cultures or with experimental animals, such as by calculating the ED₅₀ (the dose therapeutically effective in 50% of the population) or LD₅₀ (the dose lethal to 50% of the population) statistics. The dose ratio of toxic to therapeutic effects is the therapeutic index, which can be expressed as the LD₅₀/ED₅₀ ratio. Compositions which exhibit large

therapeutic indices are preferred. The data obtained from cell culture assays and animal studies are used to formulate a range of dosage for human use. The dosage contained in such compositions is preferably within a range of circulating concentrations that includes the ED_{50} with little or no toxicity. The dosage varies within this range depending upon the dosage form employed, the sensitivity of the patient, and the route of administration.

The exact dosage will be determined by the practitioner, in light of factors related to the subject requiring treatment. Dosage and administration are adjusted to provide sufficient levels of the active moiety or to maintain the desired effect. Factors which may be taken into account include the severity of the disease state, the general health of the subject, the age, weight, and gender of the subject, time and frequency of administration, drug combination(s), reaction sensitivities, and response to therapy. Long-acting compositions may be administered every 3 to 4 days, every week, or biweekly depending on the half-life and clearance rate of the particular formulation.

Normal dosage amounts may vary from about 0.1 μ g to 100,000 μ g, up to a total dose of about 1 gram, depending upon the route of administration. Guidance as to particular dosages and methods of delivery is provided in the literature and generally available to practitioners in the art. Those skilled in the art will employ different formulations for nucleotides than for proteins or their inhibitors. Similarly, delivery of polynucleotides or polypeptides will be specific to particular cells, conditions, locations, etc.

DIAGNOSTICS

In another embodiment, antibodies which specifically bind PRTS may be used for the diagnosis of disorders characterized by expression of PRTS, or in assays to monitor patients being treated with PRTS or agonists, antagonists, or inhibitors of PRTS. Antibodies useful for diagnostic purposes may be prepared in the same manner as described above for therapeutics. Diagnostic assays for PRTS include methods which utilize the antibody and a label to detect PRTS in human body fluids or in extracts of cells or tissues. The antibodies may be used with or without modification, and may be labeled by covalent or non-covalent attachment of a reporter molecule. A wide variety of reporter molecules, several of which are described above, are known in the art and may be used.

A variety of protocols for measuring PRTS, including ELISAs, RIAs, and FACS, are known in the art and provide a basis for diagnosing altered or abnormal levels of PRTS expression. Normal or standard values for PRTS expression are established by combining body fluids or cell extracts taken from normal mammalian subjects, for example, human subjects, with antibodies to PRTS under conditions suitable for complex formation. The amount of standard complex formation may be quantitated by various methods, such as photometric means. Quantities of PRTS expressed in subject,

control, and disease samples from biopsied tissues are compared with the standard values. Deviation between standard and subject values establishes the parameters for diagnosing disease.

In another embodiment of the invention, the polynucleotides encoding PRTS may be used for diagnostic purposes. The polynucleotides which may be used include oligonucleotide sequences, complementary RNA and DNA molecules, and PNAs. The polynucleotides may be used to detect and quantify gene expression in biopsied tissues in which expression of PRTS may be correlated with disease. The diagnostic assay may be used to determine absence, presence, and excess expression of PRTS, and to monitor regulation of PRTS levels during therapeutic intervention.

In one aspect, hybridization with PCR probes which are capable of detecting polynucleotide sequences, including genomic sequences, encoding PRTS or closely related molecules may be used to identify nucleic acid sequences which encode PRTS. The specificity of the probe, whether it is made from a highly specific region, e.g., the 5' regulatory region, or from a less specific region, e.g., a conserved motif, and the stringency of the hybridization or amplification will determine whether the probe identifies only naturally occurring sequences encoding PRTS, allelic variants, or related sequences.

Probes may also be used for the detection of related sequences, and may have at least 50% sequence identity to any of the PRTS encoding sequences. The hybridization probes of the subject invention may be DNA or RNA and may be derived from the sequence of SEQ ID NO:22-42 or from genomic sequences including promoters, enhancers, and introns of the PRTS gene.

Means for producing specific hybridization probes for DNAs encoding PRTS include the cloning of polynucleotide sequences encoding PRTS or PRTS derivatives into vectors for the production of mRNA probes. Such vectors are known in the art, are commercially available, and may be used to synthesize RNA probes *in vitro* by means of the addition of the appropriate RNA polymerases and the appropriate labeled nucleotides. Hybridization probes may be labeled by a variety of reporter groups, for example, by radionuclides such as ^{32}P or ^{35}S , or by enzymatic labels, such as alkaline phosphatase coupled to the probe via avidin/biotin coupling systems, and the like.

Polynucleotide sequences encoding PRTS may be used for the diagnosis of disorders associated with expression of PRTS. Examples of such disorders include, but are not limited to, a gastrointestinal disorder, such as dysphagia, peptic esophagitis, esophageal spasm, esophageal stricture, esophageal carcinoma, dyspepsia, indigestion, gastritis, gastric carcinoma, anorexia, nausea, emesis, gastroparesis, antral or pyloric edema, abdominal angina, pyrosis, gastroenteritis, intestinal obstruction, infections of the intestinal tract, peptic ulcer, cholelithiasis, cholecystitis, cholestasis, pancreatitis, pancreatic carcinoma, biliary tract disease, hepatitis, hyperbilirubinemia, cirrhosis, passive congestion of the liver, hepatoma, infectious colitis, ulcerative colitis, ulcerative proctitis, Crohn's disease, Whipple's disease,

Mallory-Weiss syndrome, colonic carcinoma, colonic obstruction, irritable bowel syndrome, short bowel syndrome, diarrhea, constipation, gastrointestinal hemorrhage, acquired immunodeficiency syndrome (AIDS) enteropathy, jaundice, hepatic encephalopathy, hepatorenal syndrome, hepatic steatosis, hemochromatosis, Wilson's disease, alpha₁-antitrypsin deficiency, Reye's syndrome, primary sclerosing cholangitis, liver infarction, portal vein obstruction and thrombosis, centrilobular necrosis, peliosis hepatis, hepatic vein thrombosis, veno-occlusive disease, preeclampsia, eclampsia, acute fatty liver of pregnancy, intrahepatic cholestasis of pregnancy, and hepatic tumors including nodular hyperplasias, adenomas, and carcinomas; a cardiovascular disorder, such as arteriovenous fistula, atherosclerosis, hypertension, vasculitis, Raynaud's disease, aneurysms, arterial dissections, varicose veins, thrombophlebitis and phlebothrombosis, vascular tumors, and complications of thrombolysis, balloon angioplasty, vascular replacement, and coronary artery bypass graft surgery, congestive heart failure, ischemic heart disease, angina pectoris, myocardial infarction, hypertensive heart disease, degenerative valvular heart disease, calcific aortic valve stenosis, congenitally bicuspid aortic valve, mitral annular calcification, mitral valve prolapse, rheumatic fever and rheumatic heart disease, infective endocarditis, nonbacterial thrombotic endocarditis, endocarditis of systemic lupus erythematosus, carcinoid heart disease, cardiomyopathy, myocarditis, pericarditis, neoplastic heart disease, congenital heart disease, and complications of cardiac transplantation; an autoimmune/inflammatory disorder, such as acquired immunodeficiency syndrome (AIDS), Addison's disease, adult respiratory distress syndrome, allergies, ankylosing spondylitis, amyloidosis, anemia, asthma, atherosclerosis, atherosclerotic plaque rupture, autoimmune hemolytic anemia, autoimmune thyroiditis, autoimmune polyendocrinopathy-candidiasis-ectodermal dystrophy (APECED), bronchitis, cholecystitis, contact dermatitis, Crohn's disease, atopic dermatitis, dermatomyositis, diabetes mellitus, emphysema, episodic lymphopenia with lymphocytotoxins, erythroblastosis fetalis, erythema nodosum, atrophic gastritis, glomerulonephritis, Goodpasture's syndrome, gout, Graves' disease, Hashimoto's thyroiditis, hypereosinophilia, irritable bowel syndrome, multiple sclerosis, myasthenia gravis, myocardial or pericardial inflammation, osteoarthritis, degradation of articular cartilage, osteoporosis, pancreatitis, polymyositis, psoriasis, Reiter's syndrome, rheumatoid arthritis, scleroderma, Sjögren's syndrome, systemic anaphylaxis, systemic lupus erythematosus, systemic sclerosis, thrombocytopenic purpura, ulcerative colitis, uveitis, Werner syndrome, complications of cancer, hemodialysis, and extracorporeal circulation, viral, bacterial, fungal, parasitic, protozoal, and helminthic infections, and trauma; a cell proliferative disorder such as actinic keratosis, arteriosclerosis, atherosclerosis, bursitis, cirrhosis, hepatitis, mixed connective tissue disease (MCTD), myelofibrosis, paroxysmal nocturnal hemoglobinuria, polycythemia vera, psoriasis, primary thrombocythemia, and cancers including adenocarcinoma, leukemia, lymphoma, melanoma, myeloma, sarcoma, teratocarcinoma, and, in

- particular, cancers of the adrenal gland, bladder, bone, bone marrow, brain, breast, cervix, gall bladder, ganglia, gastrointestinal tract, heart, kidney, liver, lung, muscle, ovary, pancreas, parathyroid, penis, prostate, salivary glands, skin, spleen, testis, thymus, thyroid, and uterus; a developmental disorder, such as renal tubular acidosis, anemia, Cushing's syndrome, achondroplastic dwarfism, Duchenne and
- 5 Becker muscular dystrophy, bone resorption, epilepsy, gonadal dysgenesis, WAGR syndrome (Wilms' tumor, aniridia, genitourinary abnormalities, and mental retardation), Smith-Magenis syndrome, myelodysplastic syndrome, hereditary mucoepithelial dysplasia, hereditary keratodermas, hereditary neuropathies such as Charcot-Marie-Tooth disease and neurofibromatosis, hypothyroidism, hydrocephalus, seizure disorders such as Sydenham's chorea and cerebral palsy, spina bifida,
 - 10 anencephaly, craniorachischisis, congenital glaucoma, cataract, age-related macular degeneration, and sensorineural hearing loss; an epithelial disorder, such as dyshidrotic eczema, allergic contact dermatitis, keratosis pilaris, melasma, vitiligo, actinic keratosis, basal cell carcinoma, squamous cell carcinoma, seborrheic keratosis, folliculitis, herpes simplex, herpes zoster, varicella, candidiasis, dermatophytosis, scabies, insect bites, cherry angioma, keloid, dermatofibroma, acrochordons,
 - 15 urticaria, transient acantholytic dermatosis, xerosis, eczema, atopic dermatitis, contact dermatitis, hand eczema, nummular eczema, lichen simplex chronicus, asteatotic eczema, stasis dermatitis and stasis ulceration, seborrheic dermatitis, psoriasis, lichen planus, pityriasis rosea, impetigo, ecthyma, dermatophytosis, tinea versicolor, warts, acne vulgaris, acne rosacea, pemphigus vulgaris, pemphigus foliaceus, paraneoplastic pemphigus, bullous pemphigoid, herpes gestationis, dermatitis herpetiformis,
 - 20 linear IgA disease, epidermolysis bullosa acquisita, dermatomyositis, lupus erythematosus, scleroderma and morphea, erythroderma, alopecia, figurate skin lesions, telangiectasias, hypopigmentation, hyperpigmentation, vesicles/bullae, exanthems, cutaneous drug reactions, papulonodular skin lesions, chronic non-healing wounds, photosensitivity diseases, epidermolysis bullosa simplex, epidermolytic hyperkeratosis, epidermolytic and nonepidermolytic palmoplantar keratoderma, ichthyosis bullosa of
 - 25 Siemens, ichthyosis exfoliativa, keratosis palmaris et plantaris, keratosis palmoplantaris, palmoplantar keratoderma, keratosis punctata, Meesmann's corneal dystrophy, pachyonychia congenita, white sponge nevus, steatocystoma multiplex, epidermal nevi/epidermolytic hyperkeratosis type, monilethrix, trichothiodystrophy, chronic hepatitis/cryptogenic cirrhosis, and colorectal hyperplasia; a neurological
 - 30 disease, Pick's disease, Huntington's disease, dementia, Parkinson's disease and other extrapyramidal disorders, amyotrophic lateral sclerosis and other motor neuron disorders, progressive neural muscular atrophy, retinitis pigmentosa, hereditary ataxias, multiple sclerosis and other demyelinating diseases, bacterial and viral meningitis, brain abscess, subdural empyema, epidural abscess, suppurative intracranial thrombophlebitis, myelitis and radiculitis, viral central nervous system disease, prion

diseases including kuru, Creutzfeldt-Jakob disease, and Gerstmann-Straussler-Scheinker syndrome, fatal familial insomnia, nutritional and metabolic diseases of the nervous system, neurofibromatosis, tuberous sclerosis, cerebelloretinal hemangioblastomatosis, encephalotrigeminal syndrome, mental retardation and other developmental disorders of the central nervous system including Down syndrome, 5 cerebral palsy, neuroskeletal disorders, autonomic nervous system disorders, cranial nerve disorders, spinal cord diseases, muscular dystrophy and other neuromuscular disorders, peripheral nervous system disorders, dermatomyositis and polymyositis, inherited, metabolic, endocrine, and toxic myopathies, myasthenia gravis, periodic paralysis, mental disorders including mood, anxiety, and schizophrenic disorders, seasonal affective disorder (SAD), akathisia, amnesia, catatonia, diabetic neuropathy, tardive 10 dyskinesia, dystonias, paranoid psychoses, postherpetic neuralgia, Tourette's disorder, progressive supranuclear palsy, corticobasal degeneration, and familial frontotemporal dementia; and a reproductive disorder, such as infertility, including tubal disease, ovulatory defects, and endometriosis, a disorder of prolactin production, a disruption of the estrous cycle, a disruption of the menstrual cycle, polycystic ovary syndrome, ovarian hyperstimulation syndrome, an endometrial or ovarian tumor, a uterine 15 fibroid, autoimmune disorders, an ectopic pregnancy, and teratogenesis; cancer of the breast, fibrocystic breast disease, and galactorrhea; a disruption of spermatogenesis, abnormal sperm physiology, cancer of the testis, cancer of the prostate, benign prostatic hyperplasia, prostatitis, Peyronie's disease, impotence, carcinoma of the male breast, and gynecomastia. The polynucleotide sequences encoding PRTS may be used in Southern or northern analysis, dot blot, or other membrane-based technologies; in 20 PCR technologies; in dipstick, pin, and multiformat ELISA-like assays; and in microarrays utilizing fluids or tissues from patients to detect altered PRTS expression. Such qualitative or quantitative methods are well known in the art.

In a particular aspect, the nucleotide sequences encoding PRTS may be useful in assays that detect the presence of associated disorders, particularly those mentioned above. The nucleotide 25 sequences encoding PRTS may be labeled by standard methods and added to a fluid or tissue sample from a patient under conditions suitable for the formation of hybridization complexes. After a suitable incubation period, the sample is washed and the signal is quantified and compared with a standard value. If the amount of signal in the patient sample is significantly altered in comparison to a control sample then the presence of altered levels of nucleotide sequences encoding PRTS in the sample 30 indicates the presence of the associated disorder. Such assays may also be used to evaluate the efficacy of a particular therapeutic treatment regimen in animal studies, in clinical trials, or to monitor the treatment of an individual patient.

In order to provide a basis for the diagnosis of a disorder associated with expression of PRTS, a normal or standard profile for expression is established. This may be accomplished by combining

body fluids or cell extracts taken from normal subjects, either animal or human, with a sequence, or a fragment thereof, encoding PRTS, under conditions suitable for hybridization or amplification.

Standard hybridization may be quantified by comparing the values obtained from normal subjects with values from an experiment in which a known amount of a substantially purified polynucleotide is used.

- 5 Standard values obtained in this manner may be compared with values obtained from samples from patients who are symptomatic for a disorder. Deviation from standard values is used to establish the presence of a disorder.

- Once the presence of a disorder is established and a treatment protocol is initiated, hybridization assays may be repeated on a regular basis to determine if the level of expression in the
10 patient begins to approximate that which is observed in the normal subject. The results obtained from successive assays may be used to show the efficacy of treatment over a period ranging from several days to months.

- With respect to cancer, the presence of an abnormal amount of transcript (either under- or overexpressed) in biopsied tissue from an individual may indicate a predisposition for the development
15 of the disease, or may provide a means for detecting the disease prior to the appearance of actual clinical symptoms. A more definitive diagnosis of this type may allow health professionals to employ preventative measures or aggressive treatment earlier thereby preventing the development or further progression of the cancer.

- Additional diagnostic uses for oligonucleotides designed from the sequences encoding PRTS
20 may involve the use of PCR. These oligomers may be chemically synthesized, generated enzymatically, or produced in vitro. Oligomers will preferably contain a fragment of a polynucleotide encoding PRTS, or a fragment of a polynucleotide complementary to the polynucleotide encoding PRTS, and will be employed under optimized conditions for identification of a specific gene or condition. Oligomers may also be employed under less stringent conditions for detection or quantification of closely related DNA
25 or RNA sequences.

- In a particular aspect, oligonucleotide primers derived from the polynucleotide sequences encoding PRTS may be used to detect single nucleotide polymorphisms (SNPs). SNPs are substitutions, insertions and deletions that are a frequent cause of inherited or acquired genetic disease in humans. Methods of SNP detection include, but are not limited to, single-stranded conformation
30 polymorphism (SSCP) and fluorescent SSCP (fSSCP) methods. In SSCP, oligonucleotide primers derived from the polynucleotide sequences encoding PRTS are used to amplify DNA using the polymerase chain reaction (PCR). The DNA may be derived, for example, from diseased or normal tissue, biopsy samples, bodily fluids, and the like. SNPs in the DNA cause differences in the secondary and tertiary structures of PCR products in single-stranded form, and these differences are detectable

using gel electrophoresis in non-denaturing gels. In fSCCP, the oligonucleotide primers are fluorescently labeled, which allows detection of the amplimers in high-throughput equipment such as DNA sequencing machines. Additionally, sequence database analysis methods, termed in silico SNP (isSNP), are capable of identifying polymorphisms by comparing the sequence of individual
5 overlapping DNA fragments which assemble into a common consensus sequence. These computer-based methods filter out sequence variations due to laboratory preparation of DNA and sequencing errors using statistical models and automated analyses of DNA sequence chromatograms. In the alternative, SNPs may be detected and characterized by mass spectrometry using, for example, the high throughput MASSARRAY system (Sequenom, Inc., San Diego CA).

10 Methods which may also be used to quantify the expression of PRTS include radiolabeling or biotinylating nucleotides, coamplification of a control nucleic acid, and interpolating results from standard curves. (See, e.g., Melby, P.C. et al. (1993) *J. Immunol. Methods* 159:235-244; Duplaa, C. et al. (1993) *Anal. Biochem.* 212:229-236.) The speed of quantitation of multiple samples may be accelerated by running the assay in a high-throughput format where the oligomer or polynucleotide of
15 interest is presented in various dilutions and a spectrophotometric or colorimetric response gives rapid quantitation.

In further embodiments, oligonucleotides or longer fragments derived from any of the polynucleotide sequences described herein may be used as elements on a microarray. The microarray can be used in transcript imaging techniques which monitor the relative expression levels of large
20 numbers of genes simultaneously as described below. The microarray may also be used to identify genetic variants, mutations, and polymorphisms. This information may be used to determine gene function, to understand the genetic basis of a disorder, to diagnose a disorder, to monitor progression/regression of disease as a function of gene expression, and to develop and monitor the activities of therapeutic agents in the treatment of disease. In particular, this information may be used
25 to develop a pharmacogenomic profile of a patient in order to select the most appropriate and effective treatment regimen for that patient. For example, therapeutic agents which are highly effective and display the fewest side effects may be selected for a patient based on his/her pharmacogenomic profile.

In another embodiment, PRTS, fragments of PRTS, or antibodies specific for PRTS may be used as elements on a microarray. The microarray may be used to monitor or measure protein-protein
30 interactions, drug-target interactions, and gene expression profiles, as described above.

A particular embodiment relates to the use of the polynucleotides of the present invention to generate a transcript image of a tissue or cell type. A transcript image represents the global pattern of gene expression by a particular tissue or cell type. Global gene expression patterns are analyzed by quantifying the number of expressed genes and their relative abundance under given conditions and at a

given time. (See Seilhamer et al., "Comparative Gene Transcript Analysis," U.S. Patent Number 5,840,484, expressly incorporated by reference herein.) Thus a transcript image may be generated by hybridizing the polynucleotides of the present invention or their complements to the totality of transcripts or reverse transcripts of a particular tissue or cell type. In one embodiment, the
5 hybridization takes place in high-throughput format, wherein the polynucleotides of the present invention or their complements comprise a subset of a plurality of elements on a microarray. The resultant transcript image would provide a profile of gene activity.

Transcript images may be generated using transcripts isolated from tissues, cell lines, biopsies, or other biological samples. The transcript image may thus reflect gene expression in vivo, as in the
10 case of a tissue or biopsy sample, or in vitro, as in the case of a cell line.

Transcript images which profile the expression of the polynucleotides of the present invention may also be used in conjunction with in vitro model systems and preclinical evaluation of pharmaceuticals, as well as toxicological testing of industrial and naturally-occurring environmental compounds. All compounds induce characteristic gene expression patterns, frequently termed
15 molecular fingerprints or toxicant signatures, which are indicative of mechanisms of action and toxicity (Nuwaysir, E.F. et al. (1999) Mol. Carcinog. 24:153-159; Steiner, S. and N.L. Anderson (2000) Toxicol. Lett. 112-113:467-471, expressly incorporated by reference herein). If a test compound has a signature similar to that of a compound with known toxicity, it is likely to share those toxic properties. These fingerprints or signatures are most useful and refined when they contain expression information
20 from a large number of genes and gene families. Ideally, a genome-wide measurement of expression provides the highest quality signature. Even genes whose expression is not altered by any tested compounds are important as well, as the levels of expression of these genes are used to normalize the rest of the expression data. The normalization procedure is useful for comparison of expression data after treatment with different compounds. While the assignment of gene function to elements of a
25 toxicant signature aids in interpretation of toxicity mechanisms, knowledge of gene function is not necessary for the statistical matching of signatures which leads to prediction of toxicity. (See, for example, Press Release 00-02 from the National Institute of Environmental Health Sciences, released February 29, 2000, available at <http://www.niehs.nih.gov/oc/news/toxchip.htm>.) Therefore, it is important and desirable in toxicological screening using toxicant signatures to include all expressed
30 gene sequences.

In one embodiment, the toxicity of a test compound is assessed by treating a biological sample containing nucleic acids with the test compound. Nucleic acids that are expressed in the treated biological sample are hybridized with one or more probes specific to the polynucleotides of the present invention, so that transcript levels corresponding to the polynucleotides of the present

invention may be quantified. The transcript levels in the treated biological sample are compared with levels in an untreated biological sample. Differences in the transcript levels between the two samples are indicative of a toxic response caused by the test compound in the treated sample.

Another particular embodiment relates to the use of the polypeptide sequences of the present invention to analyze the proteome of a tissue or cell type. The term proteome refers to the global pattern of protein expression in a particular tissue or cell type. Each protein component of a proteome can be subjected individually to further analysis. Proteome expression patterns, or profiles, are analyzed by quantifying the number of expressed proteins and their relative abundance under given conditions and at a given time. A profile of a cell's proteome may thus be generated by separating and analyzing the polypeptides of a particular tissue or cell type. In one embodiment, the separation is achieved using two-dimensional gel electrophoresis, in which proteins from a sample are separated by isoelectric focusing in the first dimension, and then according to molecular weight by sodium dodecyl sulfate slab gel electrophoresis in the second dimension (Steiner and Anderson, *supra*). The proteins are visualized in the gel as discrete and uniquely positioned spots, typically by staining the gel with an agent such as Coomassie Blue or silver or fluorescent stains. The optical density of each protein spot is generally proportional to the level of the protein in the sample. The optical densities of equivalently positioned protein spots from different samples, for example, from biological samples either treated or untreated with a test compound or therapeutic agent, are compared to identify any changes in protein spot density related to the treatment. The proteins in the spots are partially sequenced using, for example, standard methods employing chemical or enzymatic cleavage followed by mass spectrometry. The identity of the protein in a spot may be determined by comparing its partial sequence, preferably of at least 5 contiguous amino acid residues, to the polypeptide sequences of the present invention. In some cases, further sequence data may be obtained for definitive protein identification.

A proteomic profile may also be generated using antibodies specific for PRTS to quantify the levels of PRTS expression. In one embodiment, the antibodies are used as elements on a microarray, and protein expression levels are quantified by exposing the microarray to the sample and detecting the levels of protein bound to each array element (Lueking, A. et al. (1999) *Anal. Biochem.* 270:103-111; Mendoze, L.G. et al. (1999) *Biotechniques* 27:778-788). Detection may be performed by a variety of methods known in the art, for example, by reacting the proteins in the sample with a thiol- or amino-reactive fluorescent compound and detecting the amount of fluorescence bound at each array element.

Toxicant signatures at the proteome level are also useful for toxicological screening, and should be analyzed in parallel with toxicant signatures at the transcript level. There is a poor correlation between transcript and protein abundances for some proteins in some tissues (Anderson, N.L. and J. Seilhamer (1997) *Electrophoresis* 18:533-537), so proteome toxicant signatures may be useful in the

analysis of compounds which do not significantly affect the transcript image, but which alter the proteomic profile. In addition, the analysis of transcripts in body fluids is difficult, due to rapid degradation of mRNA, so proteomic profiling may be more reliable and informative in such cases.

In another embodiment, the toxicity of a test compound is assessed by treating a biological sample containing proteins with the test compound. Proteins that are expressed in the treated biological sample are separated so that the amount of each protein can be quantified. The amount of each protein is compared to the amount of the corresponding protein in an untreated biological sample. A difference in the amount of protein between the two samples is indicative of a toxic response to the test compound in the treated sample. Individual proteins are identified by sequencing the amino acid residues of the individual proteins and comparing these partial sequences to the polypeptides of the present invention.

In another embodiment, the toxicity of a test compound is assessed by treating a biological sample containing proteins with the test compound. Proteins from the biological sample are incubated with antibodies specific to the polypeptides of the present invention. The amount of protein recognized by the antibodies is quantified. The amount of protein in the treated biological sample is compared with the amount in an untreated biological sample. A difference in the amount of protein between the two samples is indicative of a toxic response to the test compound in the treated sample.

Microarrays may be prepared, used, and analyzed using methods known in the art. (See, e.g., Brennan, T.M. et al. (1995) U.S. Patent No. 5,474,796; Schena, M. et al. (1996) Proc. Natl. Acad. Sci. USA 93:10614-10619; Baldeschweiler et al. (1995) PCT application WO95/251116; Shalon, D. et al. (1995) PCT application WO95/35505; Heller, R.A. et al. (1997) Proc. Natl. Acad. Sci. USA 94:2150-2155; and Heller, M.J. et al. (1997) U.S. Patent No. 5,605,662.) Various types of microarrays are well known and thoroughly described in DNA Microarrays: A Practical Approach, M. Schena, ed. (1999) Oxford University Press, London, hereby expressly incorporated by reference.

In another embodiment of the invention, nucleic acid sequences encoding PRTS may be used to generate hybridization probes useful in mapping the naturally occurring genomic sequence. Either coding or noncoding sequences may be used, and in some instances, noncoding sequences may be preferable over coding sequences. For example, conservation of a coding sequence among members of a multi-gene family may potentially cause undesired cross hybridization during chromosomal mapping. The sequences may be mapped to a particular chromosome, to a specific region of a chromosome, or to artificial chromosome constructions, e.g., human artificial chromosomes (HACs), yeast artificial chromosomes (YACs), bacterial artificial chromosomes (BACs), bacterial P1 constructions, or single chromosome cDNA libraries. (See, e.g., Harrington, J.J. et al. (1997) Nat. Genet. 15:345-355; Price, C.M. (1993) Blood Rev. 7:127-134; and Trask, B.J. (1991) Trends Genet. 7:149-154.) Once mapped, the nucleic acid sequences of the invention may be used to develop genetic

linkage maps, for example, which correlate the inheritance of a disease state with the inheritance of a particular chromosome region or restriction fragment length polymorphism (RFLP). (See, for example, Lander, E.S. and D. Botstein (1986) Proc. Natl. Acad. Sci. USA 83:7353-7357.)

Fluorescent in situ hybridization (FISH) may be correlated with other physical and genetic map data. (See, e.g., Heinz-Ulrich, et al. (1995) in Meyers, supra, pp. 965-968.) Examples of genetic map data can be found in various scientific journals or at the Online Mendelian Inheritance in Man (OMIM) World Wide Web site. Correlation between the location of the gene encoding PRTS on a physical map and a specific disorder, or a predisposition to a specific disorder, may help define the region of DNA associated with that disorder and thus may further positional cloning efforts.

In situ hybridization of chromosomal preparations and physical mapping techniques, such as linkage analysis using established chromosomal markers, may be used for extending genetic maps. Often the placement of a gene on the chromosome of another mammalian species, such as mouse, may reveal associated markers even if the exact chromosomal locus is not known. This information is valuable to investigators searching for disease genes using positional cloning or other gene discovery techniques. Once the gene or genes responsible for a disease or syndrome have been crudely localized by genetic linkage to a particular genomic region, e.g., ataxia-telangiectasia to 11q22-23, any sequences mapping to that area may represent associated or regulatory genes for further investigation. (See, e.g., Gatti, R.A. et al. (1988) Nature 336:577-580.) The nucleotide sequence of the instant invention may also be used to detect differences in the chromosomal location due to translocation, inversion, etc., among normal, carrier, or affected individuals.

In another embodiment of the invention, PRTS, its catalytic or immunogenic fragments, or oligopeptides thereof can be used for screening libraries of compounds in any of a variety of drug screening techniques. The fragment employed in such screening may be free in solution, affixed to a solid support, borne on a cell surface, or located intracellularly. The formation of binding complexes between PRTS and the agent being tested may be measured.

Another technique for drug screening provides for high throughput screening of compounds having suitable binding affinity to the protein of interest. (See, e.g., Geysen, et al. (1984) PCT application WO84/03564.) In this method, large numbers of different small test compounds are synthesized on a solid substrate. The test compounds are reacted with PRTS, or fragments thereof, and washed. Bound PRTS is then detected by methods well known in the art. Purified PRTS can also be coated directly onto plates for use in the aforementioned drug screening techniques. Alternatively, non-neutralizing antibodies can be used to capture the peptide and immobilize it on a solid support.

In another embodiment, one may use competitive drug screening assays in which neutralizing antibodies capable of binding PRTS specifically compete with a test compound for binding PRTS. In

this manner, antibodies can be used to detect the presence of any peptide which shares one or more antigenic determinants with PRTS.

In additional embodiments, the nucleotide sequences which encode PRTS may be used in any molecular biology techniques that have yet to be developed, provided the new techniques rely on properties of nucleotide sequences that are currently known, including, but not limited to, such properties as the triplet genetic code and specific base pair interactions.

Without further elaboration, it is believed that one skilled in the art can, using the preceding description, utilize the present invention to its fullest extent. The following embodiments are, therefore, to be construed as merely illustrative, and not limitative of the remainder of the disclosure in any way whatsoever.

The disclosures of all patents, applications, and publications mentioned above and below, including U.S. Ser. No. 60/212,336, U.S. Ser. No. 60/213,995, U.S. Ser. No. 60/215,396, U.S. Ser. No. 60/216,821, and U.S. Ser. No. 60/218,946, are hereby expressly incorporated by reference.

15

EXAMPLES

I. Construction of cDNA Libraries

Incyte cDNAs were derived from cDNA libraries described in the LIFESEQ GOLD database (Incyte Genomics, Palo Alto CA) and shown in Table 4, column 5. Some tissues were homogenized and lysed in guanidinium isothiocyanate, while others were homogenized and lysed in phenol or in a suitable mixture of denaturants, such as TRIZOL (Life Technologies), a monophasic solution of phenol and guanidine isothiocyanate. The resulting lysates were centrifuged over CsCl cushions or extracted with chloroform. RNA was precipitated from the lysates with either isopropanol or sodium acetate and ethanol, or by other routine methods.

Phenol extraction and precipitation of RNA were repeated as necessary to increase RNA purity. In some cases, RNA was treated with DNase. For most libraries, poly(A)+ RNA was isolated using oligo d(T)-coupled paramagnetic particles (Promega), OLIGOTEX latex particles (QIAGEN, Chatsworth CA), or an OLIGOTEX mRNA purification kit (QIAGEN). Alternatively, RNA was isolated directly from tissue lysates using other RNA isolation kits, e.g., the POLY(A)PURE mRNA purification kit (Ambion, Austin TX).

In some cases, Stratagene was provided with RNA and constructed the corresponding cDNA libraries. Otherwise, cDNA was synthesized and cDNA libraries were constructed with the UNIZAP vector system (Stratagene) or SUPERScript plasmid system (Life Technologies), using the recommended procedures or similar methods known in the art. (See, e.g., Ausubel, 1997, *supra*, units 5.1-6.6.) Reverse transcription was initiated using oligo d(T) or random primers. Synthetic

oligonucleotide adapters were ligated to double stranded cDNA, and the cDNA was digested with the appropriate restriction enzyme or enzymes. For most libraries, the cDNA was size-selected (300-1000 bp) using SEPHACRYL S1000, SEPHAROSE CL2B, or SEPHAROSE CL4B column chromatography (Amersham Pharmacia Biotech) or preparative agarose gel electrophoresis. cDNAs were ligated into compatible restriction enzyme sites of the polylinker of a suitable plasmid, e.g., PBLUESCRIPT plasmid (Stratagene), PSPO1 plasmid (Life Technologies), PCDNA2.1 plasmid (Invitrogen, Carlsbad CA), PBK-CMV plasmid (Stratagene), or pINCY (Incyte Genomics, Palo Alto CA), or derivatives thereof. Recombinant plasmids were transformed into competent *E. coli* cells including XL1-Blue, XL1-BlueMRF, or SOLR from Stratagene or DH5 α , DH10B, or ElectroMAX DH10B from Life Technologies.

II. Isolation of cDNA Clones

Plasmids obtained as described in Example I were recovered from host cells by *in vivo* excision using the UNIZAP vector system (Stratagene) or by cell lysis. Plasmids were purified using at least one of the following: a Magic or WIZARD Minipreps DNA purification system (Promega); an AGTC Miniprep purification kit (Edge Biosystems, Gaithersburg MD); and QIAWELL 8 Plasmid, QIAWELL 8 Plus Plasmid, QIAWELL 8 Ultra Plasmid purification systems or the R.E.A.L. PREP 96 plasmid purification kit from QIAGEN. Following precipitation, plasmids were resuspended in 0.1 ml of distilled water and stored, with or without lyophilization, at 4°C.

Alternatively, plasmid DNA was amplified from host cell lysates using direct link PCR in a high-throughput format (Rao, V.B. (1994) Anal. Biochem. 216:1-14). Host cell lysis and thermal cycling steps were carried out in a single reaction mixture. Samples were processed and stored in 384-well plates, and the concentration of amplified plasmid DNA was quantified fluorometrically using PICOGREEN dye (Molecular Probes, Eugene OR) and a FLUOROSKAN II fluorescence scanner (Labsystems Oy, Helsinki, Finland).

III. Sequencing and Analysis

Incyte cDNA recovered in plasmids as described in Example II were sequenced as follows. Sequencing reactions were processed using standard methods or high-throughput instrumentation such as the ABI CATALYST 800 (Applied Biosystems) thermal cycler or the PTC-200 thermal cycler (MJ Research) in conjunction with the HYDRA microdispenser (Robbins Scientific) or the MICROLAB 2200 (Hamilton) liquid transfer system. cDNA sequencing reactions were prepared using reagents provided by Amersham Pharmacia Biotech or supplied in ABI sequencing kits such as the ABI PRISM BIGDYE Terminator cycle sequencing ready reaction kit (Applied Biosystems). Electrophoretic separation of cDNA sequencing reactions and detection of labeled polynucleotides were carried out using the MEGABACE 1000 DNA sequencing system (Molecular Dynamics); the ABI

PRISM 373 or 377 sequencing system (Applied Biosystems) in conjunction with standard ABI protocols and base calling software; or other sequence analysis systems known in the art. Reading frames within the cDNA sequences were identified using standard methods (reviewed in Ausubel, 1997, supra, unit 7.7). Some of the cDNA sequences were selected for extension using the techniques disclosed in Example VIII.

The polynucleotide sequences derived from Incyte cDNAs were validated by removing vector, linker, and poly(A) sequences and by masking ambiguous bases, using algorithms and programs based on BLAST, dynamic programming, and dinucleotide nearest neighbor analysis. The Incyte cDNA sequences or translations thereof were then queried against a selection of public databases such as the GenBank primate, rodent, mammalian, vertebrate, and eukaryote databases, and BLOCKS, PRINTS, DOMO, PRODOM, and hidden Markov model (HMM)-based protein family databases such as PFAM. (HMM is a probabilistic approach which analyzes consensus primary structures of gene families. See, for example, Eddy, S.R. (1996) Curr. Opin. Struct. Biol. 6:361-365.) The queries were performed using programs based on BLAST, FASTA, BLIMPS, and HMMER. The Incyte cDNA sequences were assembled to produce full length polynucleotide sequences. Alternatively, GenBank cDNAs, GenBank ESTs, stitched sequences, stretched sequences, or Genscan-predicted coding sequences (see Examples IV and V) were used to extend Incyte cDNA assemblages to full length. Assembly was performed using programs based on Phred, Phrap, and Consed, and cDNA assemblages were screened for open reading frames using programs based on GeneMark, BLAST, and FASTA. The full length polynucleotide sequences were translated to derive the corresponding full length polypeptide sequences. Alternatively, a polypeptide of the invention may begin at any of the methionine residues of the full length translated polypeptide. Full length polypeptide sequences were subsequently analyzed by querying against databases such as the GenBank protein databases (genpept), SwissProt, BLOCKS, PRINTS, DOMO, PRODOM, Prosite, and hidden Markov model (HMM)-based protein family databases such as PFAM. Full length polynucleotide sequences are also analyzed using MACDNASIS PRO software (Hitachi Software Engineering, South San Francisco CA) and LASERGENE software (DNASTAR). Polynucleotide and polypeptide sequence alignments are generated using default parameters specified by the CLUSTAL algorithm as incorporated into the MEGALIGN multisequence alignment program (DNASTAR), which also calculates the percent identity between aligned sequences.

Table 7 summarizes the tools, programs, and algorithms used for the analysis and assembly of Incyte cDNA and full length sequences and provides applicable descriptions, references, and threshold parameters. The first column of Table 7 shows the tools, programs, and algorithms used, the second column provides brief descriptions thereof, the third column presents appropriate references, all of

which are incorporated by reference herein in their entirety, and the fourth column presents, where applicable, the scores, probability values, and other parameters used to evaluate the strength of a match between two sequences (the higher the score or the lower the probability value, the greater the identity between two sequences).

5 The programs described above for the assembly and analysis of full length polynucleotide and polypeptide sequences were also used to identify polynucleotide sequence fragments from SEQ ID NO:22-42. Fragments from about 20 to about 4000 nucleotides which are useful in hybridization and amplification technologies are described in Table 4, column 4.

IV. Identification and Editing of Coding Sequences from Genomic DNA

10 Putative proteases were initially identified by running the Genscan gene identification program against public genomic sequence databases (e.g., gbpri and gbhtg). Genscan is a general-purpose gene identification program which analyzes genomic DNA sequences from a variety of organisms (See Burge, C. and S. Karlin (1997) J. Mol. Biol. 268:78-94, and Burge, C. and S. Karlin (1998) Curr. Opin. Struct. Biol. 8:346-354). The program concatenates predicted exons to form an assembled
15 cDNA sequence extending from a methionine to a stop codon. The output of Genscan is a FASTA database of polynucleotide and polypeptide sequences. The maximum range of sequence for Genscan to analyze at once was set to 30 kb. To determine which of these Genscan predicted cDNA sequences encode proteases, the encoded polypeptides were analyzed by querying against PFAM models for proteases. Potential proteases were also identified by homology to Incyte cDNA sequences that had
20 been annotated as proteases. These selected Genscan-predicted sequences were then compared by BLAST analysis to the genpept and gbpri public databases. Where necessary, the Genscan-predicted sequences were then edited by comparison to the top BLAST hit from genpept to correct errors in the sequence predicted by Genscan, such as extra or omitted exons. BLAST analysis was also used to find any Incyte cDNA or public cDNA coverage of the Genscan-predicted sequences, thus providing
25 evidence for transcription. When Incyte cDNA coverage was available, this information was used to correct or confirm the Genscan predicted sequence. Full length polynucleotide sequences were obtained by assembling Genscan-predicted coding sequences with Incyte cDNA sequences and/or public cDNA sequences using the assembly process described in Example III. Alternatively, full length polynucleotide sequences were derived entirely from edited or unedited Genscan-predicted coding
30 sequences.

V. Assembly of Genomic Sequence Data with cDNA Sequence Data

"Stitched" Sequences

Partial cDNA sequences were extended with exons predicted by the Genscan gene identification program described in Example IV. Partial cDNAs assembled as described in Example III were mapped

to genomic DNA and parsed into clusters containing related cDNAs and Genscan exon predictions from one or more genomic sequences. Each cluster was analyzed using an algorithm based on graph theory and dynamic programming to integrate cDNA and genomic information, generating possible splice variants that were subsequently confirmed, edited, or extended to create a full length sequence.

- 5 Sequence intervals in which the entire length of the interval was present on more than one sequence in the cluster were identified, and intervals thus identified were considered to be equivalent by transitivity. For example, if an interval was present on a cDNA and two genomic sequences, then all three intervals were considered to be equivalent. This process allows unrelated but consecutive genomic sequences to be brought together, bridged by cDNA sequence. Intervals thus identified were then "stitched" together
- 10 by the stitching algorithm in the order that they appear along their parent sequences to generate the longest possible sequence, as well as sequence variants. Linkages between intervals which proceed along one type of parent sequence (cDNA to cDNA or genomic sequence to genomic sequence) were given preference over linkages which change parent type (cDNA to genomic sequence). The resultant stitched sequences were translated and compared by BLAST analysis to the genpept and gbpr public
- 15 databases. Incorrect exons predicted by Genscan were corrected by comparison to the top BLAST hit from genpept. Sequences were further extended with additional cDNA sequences, or by inspection of genomic DNA, when necessary.

"Stretched" Sequences

- Partial DNA sequences were extended to full length with an algorithm based on BLAST
- 20 analysis. First, partial cDNAs assembled as described in Example III were queried against public databases such as the GenBank primate, rodent, mammalian, vertebrate, and eukaryote databases using the BLAST program. The nearest GenBank protein homolog was then compared by BLAST analysis to either Incyte cDNA sequences or GenScan exon predicted sequences described in Example IV. A chimeric protein was generated by using the resultant high-scoring segment pairs (HSPs) to map the
- 25 translated sequences onto the GenBank protein homolog. Insertions or deletions may occur in the chimeric protein with respect to the original GenBank protein homolog. The GenBank protein homolog, the chimeric protein, or both were used as probes to search for homologous genomic sequences from the public human genome databases. Partial DNA sequences were therefore "stretched" or extended by the addition of homologous genomic sequences. The resultant stretched sequences were examined to
- 30 determine whether it contained a complete gene.

VI. Chromosomal Mapping of PRTS Encoding Polynucleotides

The sequences which were used to assemble SEQ ID NO:22-42 were compared with sequences from the Incyte LIFESEQ database and public domain databases using BLAST and other implementations of the Smith-Waterman algorithm. Sequences from these databases that matched

SEQ ID NO:22-42 were assembled into clusters of contiguous and overlapping sequences using assembly algorithms such as Phrap (Table 7). Radiation hybrid and genetic mapping data available from public resources such as the Stanford Human Genome Center (SHGC), Whitehead Institute for Genome Research (WIGR), and Généthon were used to determine if any of the clustered sequences
 5 had been previously mapped. Inclusion of a mapped sequence in a cluster resulted in the assignment of all sequences of that cluster, including its particular SEQ ID NO., to that map location.

Map locations are represented by ranges, or intervals, of human chromosomes. The map position of an interval, in centiMorgans, is measured relative to the terminus of the chromosome's p-arm. (The centiMorgan (cM) is a unit of measurement based on recombination frequencies between
 10 chromosomal markers. On average, 1 cM is roughly equivalent to 1 megabase (Mb) of DNA in humans, although this can vary widely due to hot and cold spots of recombination.) The cM distances are based on genetic markers mapped by Généthon which provide boundaries for radiation hybrid markers whose sequences were included in each of the clusters. Human genome maps and other resources available to the public, such as the NCBI "GeneMap'99" World Wide Web site
 15 (<http://www.ncbi.nlm.nih.gov/genemap/>), can be employed to determine if previously identified disease genes map within or in proximity to the intervals indicated above.

In this manner, SEQ ID NO:25 was mapped to chromosome 5 within the interval from 69.60 to 76.50 centiMorgans. SEQ ID NO:28 was mapped to chromosome 16 within the interval from 81.80 to 84.40 centiMorgans.

20 VII. Analysis of Polynucleotide Expression

Northern analysis is a laboratory technique used to detect the presence of a transcript of a gene and involves the hybridization of a labeled nucleotide sequence to a membrane on which RNAs from a particular cell type or tissue have been bound. (See, e.g., Sambrook, *supra*, ch. 7; Ausubel (1995) *supra*, ch. 4 and 16.)

25 Analogous computer techniques applying BLAST were used to search for identical or related molecules in cDNA databases such as GenBank or LIFESEQ (Incyte Genomics). This analysis is much faster than multiple membrane-based hybridizations. In addition, the sensitivity of the computer search can be modified to determine whether any particular match is categorized as exact or similar. The basis of the search is the product score, which is defined as:

30

$$\frac{\text{BLAST Score} \times \text{Percent Identity}}{5 \times \text{minimum} \{ \text{length}(\text{Seq. 1}), \text{length}(\text{Seq. 2}) \}}$$

The product score takes into account both the degree of similarity between two sequences and the length
 35 of the sequence match. The product score is a normalized value between 0 and 100, and is calculated

as follows: the BLAST score is multiplied by the percent nucleotide identity and the product is divided by (5 times the length of the shorter of the two sequences). The BLAST score is calculated by assigning a score of +5 for every base that matches in a high-scoring segment pair (HSP), and -4 for every mismatch. Two sequences may share more than one HSP (separated by gaps). If there is more than one HSP, then the pair with the highest BLAST score is used to calculate the product score. The product score represents a balance between fractional overlap and quality in a BLAST alignment. For example, a product score of 100 is produced only for 100% identity over the entire length of the shorter of the two sequences being compared. A product score of 70 is produced either by 100% identity and 70% overlap at one end, or by 88% identity and 100% overlap at the other. A product score of 50 is produced either by 100% identity and 50% overlap at one end, or 79% identity and 100% overlap.

Alternatively, polynucleotide sequences encoding PRTS are analyzed with respect to the tissue sources from which they were derived. For example, some full length sequences are assembled, at least in part, with overlapping Incyte cDNA sequences (see Example III). Each cDNA sequence is derived from a cDNA library constructed from a human tissue. Each human tissue is classified into one of the following organ/tissue categories: cardiovascular system; connective tissue; digestive system; embryonic structures; endocrine system; exocrine glands; genitalia, female; genitalia, male; germ cells; hemic and immune system; liver; musculoskeletal system; nervous system; pancreas; respiratory system; sense organs; skin; stomatognathic system; unclassified/mixed; or urinary tract. The number of libraries in each category is counted and divided by the total number of libraries across all categories. Similarly, each human tissue is classified into one of the following disease/condition categories: cancer, cell line, developmental, inflammation, neurological, trauma, cardiovascular, pooled, and other, and the number of libraries in each category is counted and divided by the total number of libraries across all categories. The resulting percentages reflect the tissue- and disease-specific expression of cDNA encoding PRTS. cDNA sequences and cDNA library/tissue information are found in the LIFESEQ GOLD database (Incyte Genomics, Palo Alto CA).

VIII. Extension of PRTS Encoding Polynucleotides

Full length polynucleotide sequences were also produced by extension of an appropriate fragment of the full length molecule using oligonucleotide primers designed from this fragment. One primer was synthesized to initiate 5' extension of the known fragment, and the other primer was synthesized to initiate 3' extension of the known fragment. The initial primers were designed using OLIGO 4.06 software (National Biosciences), or another appropriate program, to be about 22 to 30 nucleotides in length, to have a GC content of about 50% or more, and to anneal to the target sequence at temperatures of about 68°C to about 72°C. Any stretch of nucleotides which would result in hairpin structures and primer-primer dimerizations was avoided.

Selected human cDNA libraries were used to extend the sequence. If more than one extension was necessary or desired, additional or nested sets of primers were designed.

High fidelity amplification was obtained by PCR using methods well known in the art. PCR was performed in 96-well plates using the PTC-200 thermal cycler (MJ Research, Inc.). The reaction mix contained DNA template, 200 nmol of each primer, reaction buffer containing Mg^{2+} , $(NH_4)_2SO_4$, and 2-mercaptoethanol, Taq DNA polymerase (Amersham Pharmacia Biotech), ELONGASE enzyme (Life Technologies), and Pfu DNA polymerase (Stratagene); with the following parameters for primer pair PCI A and PCI B: Step 1: 94°C, 3 min; Step 2: 94°C, 15 sec; Step 3: 60°C, 1 min; Step 4: 68°C, 2 min; Step 5: Steps 2, 3, and 4 repeated 20 times; Step 6: 68°C, 5 min; Step 7: storage at 4°C. In the alternative, the parameters for primer pair T7 and SK+ were as follows: Step 1: 94°C, 3 min; Step 2: 94°C, 15 sec; Step 3: 57°C, 1 min; Step 4: 68°C, 2 min; Step 5: Steps 2, 3, and 4 repeated 20 times; Step 6: 68°C, 5 min; Step 7: storage at 4°C.

The concentration of DNA in each well was determined by dispensing 100 μ l PICOGREEN quantitation reagent (0.25% (v/v) PICOGREEN; Molecular Probes, Eugene OR) dissolved in 1X TE and 0.5 μ l of undiluted PCR product into each well of an opaque fluorimeter plate (Corning Costar, Acton MA), allowing the DNA to bind to the reagent. The plate was scanned in a Fluoroskan II (Labsystems Oy, Helsinki, Finland) to measure the fluorescence of the sample and to quantify the concentration of DNA. A 5 μ l to 10 μ l aliquot of the reaction mixture was analyzed by electrophoresis on a 1 % agarose gel to determine which reactions were successful in extending the sequence.

The extended nucleotides were desalted and concentrated, transferred to 384-well plates, digested with CviJI cholera virus endonuclease (Molecular Biology Research, Madison WI), and sonicated or sheared prior to religation into pUC 18 vector (Amersham Pharmacia Biotech). For shotgun sequencing, the digested nucleotides were separated on low concentration (0.6 to 0.8%) agarose gels, fragments were excised, and agar digested with Agar ACE (Promega). Extended clones were religated using T4 ligase (New England Biolabs, Beverly MA) into pUC 18 vector (Amersham Pharmacia Biotech), treated with Pfu DNA polymerase (Stratagene) to fill-in restriction site overhangs, and transfected into competent *E. coli* cells. Transformed cells were selected on antibiotic-containing media, and individual colonies were picked and cultured overnight at 37°C in 384-well plates in LB/2x carb liquid media.

The cells were lysed, and DNA was amplified by PCR using Taq DNA polymerase (Amersham Pharmacia Biotech) and Pfu DNA polymerase (Stratagene) with the following parameters: Step 1: 94°C, 3 min; Step 2: 94°C, 15 sec; Step 3: 60°C, 1 min; Step 4: 72°C, 2 min; Step 5: steps 2, 3, and 4 repeated 29 times; Step 6: 72°C, 5 min; Step 7: storage at 4°C. DNA was quantified by PICOGREEN reagent (Molecular Probes) as described above. Samples with low DNA recoveries were reamplified

using the same conditions as described above. Samples were diluted with 20% dimethylsulfoxide (1:2, v/v), and sequenced using DYENAMIC energy transfer sequencing primers and the DYENAMIC DIRECT kit (Amersham Pharmacia Biotech) or the ABI PRISM BIGDYE Terminator cycle sequencing ready reaction kit (Applied Biosystems).

- 5 In like manner, full length polynucleotide sequences are verified using the above procedure or are used to obtain 5' regulatory sequences using the above procedure along with oligonucleotides designed for such extension, and an appropriate genomic library.

IX. Labeling and Use of Individual Hybridization Probes

- Hybridization probes derived from SEQ ID NO:22-42 are employed to screen cDNAs, genomic
10 DNAs, or mRNAs. Although the labeling of oligonucleotides, consisting of about 20 base pairs, is specifically described, essentially the same procedure is used with larger nucleotide fragments. Oligonucleotides are designed using state-of-the-art software such as OLIGO 4.06 software (National Biosciences) and labeled by combining 50 pmol of each oligomer, 250 μ Ci of [γ -³²P] adenosine triphosphate (Amersham Pharmacia Biotech), and T4 polynucleotide kinase (DuPont NEN, Boston
15 MA). The labeled oligonucleotides are substantially purified using a SEPHADEX G-25 superfine size exclusion dextran bead column (Amersham Pharmacia Biotech). An aliquot containing 10⁷ counts per minute of the labeled probe is used in a typical membrane-based hybridization analysis of human genomic DNA digested with one of the following endonucleases: Ase I, Bgl II, Eco RI, Pst I, Xba I, or Pvu II (DuPont NEN).

- 20 The DNA from each digest is fractionated on a 0.7% agarose gel and transferred to nylon membranes (Nytran Plus, Schleicher & Schuell, Durham NH). Hybridization is carried out for 16 hours at 40°C. To remove nonspecific signals, blots are sequentially washed at room temperature under conditions of up to, for example, 0.1 x saline sodium citrate and 0.5% sodium dodecyl sulfate. Hybridization patterns are visualized using autoradiography or an alternative imaging means and
25 compared.

X. Microarrays

- The linkage or synthesis of array elements upon a microarray can be achieved utilizing photolithography, piezoelectric printing (ink-jet printing, See, e.g., Baldeschweiler, *supra*), mechanical microspotting technologies, and derivatives thereof. The substrate in each of the aforementioned
30 technologies should be uniform and solid with a non-porous surface (Schena (1999), *supra*). Suggested substrates include silicon, silica, glass slides, glass chips, and silicon wafers. Alternatively, a procedure analogous to a dot or slot blot may also be used to arrange and link elements to the surface of a substrate using thermal, UV, chemical, or mechanical bonding procedures. A typical array may be produced using available methods and machines well known to those of ordinary skill in the art and may

contain any appropriate number of elements. (See, e.g., Schena, M. et al. (1995) Science 270:467-470; Shalon, D. et al. (1996) Genome Res. 6:639-645; Marshall, A. and J. Hodgson (1998) Nat. Biotechnol. 16:27-31.)

Full length cDNAs, Expressed Sequence Tags (ESTs), or fragments or oligomers thereof may
5 comprise the elements of the microarray. Fragments or oligomers suitable for hybridization can be selected using software well known in the art such as LASERGENE software (DNASTAR). The array elements are hybridized with polynucleotides in a biological sample. The polynucleotides in the biological sample are conjugated to a fluorescent label or other molecular tag for ease of detection. After hybridization, nonhybridized nucleotides from the biological sample are removed, and a
10 fluorescence scanner is used to detect hybridization at each array element. Alternatively, laser desorption and mass spectrometry may be used for detection of hybridization. The degree of complementarity and the relative abundance of each polynucleotide which hybridizes to an element on the microarray may be assessed. In one embodiment, microarray preparation and usage is described in detail below.

15 Tissue or Cell Sample Preparation

Total RNA is isolated from tissue samples using the guanidinium thiocyanate method and poly(A)⁺ RNA is purified using the oligo-(dT) cellulose method. Each poly(A)⁺ RNA sample is reverse transcribed using MMLV reverse-transcriptase, 0.05 pg/ μ l oligo-(dT) primer (21mer), 1X first strand buffer, 0.03 units/ μ l RNase inhibitor, 500 μ M dATP, 500 μ M dGTP, 500 μ M dTTP, 40 μ M
20 dCTP, 40 μ M dCTP-Cy3 (BDS) or dCTP-Cy5 (Amersham Pharmacia Biotech). The reverse transcription reaction is performed in a 25 ml volume containing 200 ng poly(A)⁺ RNA with GEMBRIGHT kits (Incyte). Specific control poly(A)⁺ RNAs are synthesized by in vitro transcription from non-coding yeast genomic DNA. After incubation at 37° C for 2 hr, each reaction sample (one with Cy3 and another with Cy5 labeling) is treated with 2.5 ml of 0.5M sodium hydroxide and
25 incubated for 20 minutes at 85° C to stop the reaction and degrade the RNA. Samples are purified using two successive CHROMA SPIN 30 gel filtration spin columns (CLONTECH Laboratories, Inc. (CLONTECH), Palo Alto CA) and after combining, both reaction samples are ethanol precipitated using 1 ml of glycogen (1 mg/ml), 60 ml sodium acetate, and 300 ml of 100% ethanol. The sample is then dried to completion using a SpeedVAC (Savant Instruments Inc., Holbrook NY) and
30 resuspended in 14 μ l 5X SSC/0.2% SDS.

Microarray Preparation

Sequences of the present invention are used to generate array elements. Each array element is amplified from bacterial cells containing vectors with cloned cDNA inserts. PCR amplification uses primers complementary to the vector sequences flanking the cDNA insert. Array elements are

amplified in thirty cycles of PCR from an initial quantity of 1-2 ng to a final quantity greater than 5 µg. Amplified array elements are then purified using SEPHACRYL-400 (Amersham Pharmacia Biotech).

Purified array elements are immobilized on polymer-coated glass slides. Glass microscope slides (Corning) are cleaned by ultrasound in 0.1% SDS and acetone, with extensive distilled water washes between and after treatments. Glass slides are etched in 4% hydrofluoric acid (VWR Scientific Products Corporation (VWR), West Chester PA), washed extensively in distilled water, and coated with 0.05% aminopropyl silane (Sigma) in 95% ethanol. Coated slides are cured in a 110°C oven.

Array elements are applied to the coated glass substrate using a procedure described in US Patent No. 5,807,522, incorporated herein by reference. 1 µl of the array element DNA, at an average concentration of 100 ng/µl, is loaded into the open capillary printing element by a high-speed robotic apparatus. The apparatus then deposits about 5 nl of array element sample per slide.

Microarrays are UV-crosslinked using a STRATALINKER UV-crosslinker (Stratagene).

Microarrays are washed at room temperature once in 0.2% SDS and three times in distilled water. Non-specific binding sites are blocked by incubation of microarrays in 0.2% casein in phosphate buffered saline (PBS) (Tropix, Inc., Bedford MA) for 30 minutes at 60°C followed by washes in 0.2% SDS and distilled water as before.

Hybridization

Hybridization reactions contain 9 µl of sample mixture consisting of 0.2 µg each of Cy3 and Cy5 labeled cDNA synthesis products in 5X SSC, 0.2% SDS hybridization buffer. The sample mixture is heated to 65°C for 5 minutes and is aliquoted onto the microarray surface and covered with an 1.8 cm² coverslip. The arrays are transferred to a waterproof chamber having a cavity just slightly larger than a microscope slide. The chamber is kept at 100% humidity internally by the addition of 140 µl of 5X SSC in a corner of the chamber. The chamber containing the arrays is incubated for about 6.5 hours at 60°C. The arrays are washed for 10 min at 45°C in a first wash buffer (1X SSC, 0.1% SDS), three times for 10 minutes each at 45°C in a second wash buffer (0.1X SSC), and dried.

Detection

Reporter-labeled hybridization complexes are detected with a microscope equipped with an Innova 70 mixed gas 10 W laser (Coherent, Inc., Santa Clara CA) capable of generating spectral lines at 488 nm for excitation of Cy3 and at 632 nm for excitation of Cy5. The excitation laser light is focused on the array using a 20X microscope objective (Nikon, Inc., Melville NY). The slide containing the array is placed on a computer-controlled X-Y stage on the microscope and raster-scanned past the objective. The 1.8 cm x 1.8 cm array used in the present example is scanned with a resolution of 20 micrometers.

In two separate scans, a mixed gas multiline laser excites the two fluorophores sequentially. Emitted light is split, based on wavelength, into two photomultiplier tube detectors (PMT R1477, Hamamatsu Photonics Systems, Bridgewater NJ) corresponding to the two fluorophores. Appropriate filters positioned between the array and the photomultiplier tubes are used to filter the signals. The
5 emission maxima of the fluorophores used are 565 nm for Cy3 and 650 nm for Cy5. Each array is typically scanned twice, one scan per fluorophore using the appropriate filters at the laser source, although the apparatus is capable of recording the spectra from both fluorophores simultaneously.

The sensitivity of the scans is typically calibrated using the signal intensity generated by a cDNA control species added to the sample mixture at a known concentration. A specific location on
10 the array contains a complementary DNA sequence, allowing the intensity of the signal at that location to be correlated with a weight ratio of hybridizing species of 1:100,000. When two samples from different sources (e.g., representing test and control cells), each labeled with a different fluorophore, are hybridized to a single array for the purpose of identifying genes that are differentially expressed, the calibration is done by labeling samples of the calibrating cDNA with the two
15 fluorophores and adding identical amounts of each to the hybridization mixture.

The output of the photomultiplier tube is digitized using a 12-bit RTI-835H analog-to-digital (A/D) conversion board (Analog Devices, Inc., Norwood MA) installed in an IBM-compatible PC computer. The digitized data are displayed as an image where the signal intensity is mapped using a linear 20-color transformation to a pseudocolor scale ranging from blue (low signal) to red (high
20 signal). The data is also analyzed quantitatively. Where two different fluorophores are excited and measured simultaneously, the data are first corrected for optical crosstalk (due to overlapping emission spectra) between the fluorophores using each fluorophore's emission spectrum.

A grid is superimposed over the fluorescence signal image such that the signal from each spot is centered in each element of the grid. The fluorescence signal within each element is then integrated
25 to obtain a numerical value corresponding to the average intensity of the signal. The software used for signal analysis is the GEMTOOLS gene expression analysis program (Incyte).

XI. Complementary Polynucleotides

Sequences complementary to the PRTS-encoding sequences, or any parts thereof, are used to detect, decrease, or inhibit expression of naturally occurring PRTS. Although use of oligonucleotides
30 comprising from about 15 to 30 base pairs is described, essentially the same procedure is used with smaller or with larger sequence fragments. Appropriate oligonucleotides are designed using OLIGO 4.06 software (National Biosciences) and the coding sequence of PRTS. To inhibit transcription, a complementary oligonucleotide is designed from the most unique 5' sequence and used to prevent promoter binding to the coding sequence. To inhibit translation, a complementary oligonucleotide is
35 designed to prevent ribosomal binding to the PRTS-encoding transcript.

XII. Expression of PRTS

Expression and purification of PRTS is achieved using bacterial or virus-based expression systems. For expression of PRTS in bacteria, cDNA is subcloned into an appropriate vector containing an antibiotic resistance gene and an inducible promoter that directs high levels of cDNA transcription.

- 5 Examples of such promoters include, but are not limited to, the *trp-lac (tac)* hybrid promoter and the T5 or T7 bacteriophage promoter in conjunction with the *lac* operator regulatory element.

Recombinant vectors are transformed into suitable bacterial hosts, e.g., BL21(DE3). Antibiotic resistant bacteria express PRTS upon induction with isopropyl beta-D-thiogalactopyranoside (IPTG).

- Expression of PRTS in eukaryotic cells is achieved by infecting insect or mammalian cell lines with
10 recombinant Autographica californica nuclear polyhedrosis virus (AcMNPV), commonly known as baculovirus. The nonessential polyhedrin gene of baculovirus is replaced with cDNA encoding PRTS by either homologous recombination or bacterial-mediated transposition involving transfer plasmid intermediates. Viral infectivity is maintained and the strong polyhedrin promoter drives high levels of cDNA transcription. Recombinant baculovirus is used to infect Spodoptera frugiperda (Sf9) insect
15 cells in most cases, or human hepatocytes, in some cases. Infection of the latter requires additional genetic modifications to baculovirus. (See Engelhard, E.K. et al. (1994) Proc. Natl. Acad. Sci. USA 91:3224-3227; Sandig, V. et al. (1996) Hum. Gene Ther. 7:1937-1945.)

- In most expression systems, PRTS is synthesized as a fusion protein with, e.g., glutathione S-transferase (GST) or a peptide epitope tag, such as FLAG or 6-His, permitting rapid, single-step,
20 affinity-based purification of recombinant fusion protein from crude cell lysates. GST, a 26-kilodalton enzyme from Schistosoma japonicum, enables the purification of fusion proteins on immobilized glutathione under conditions that maintain protein activity and antigenicity (Amersham Pharmacia Biotech). Following purification, the GST moiety can be proteolytically cleaved from PRTS at specifically engineered sites. FLAG, an 8-amino acid peptide, enables immunoaffinity purification
25 using commercially available monoclonal and polyclonal anti-FLAG antibodies (Eastman Kodak). 6-His, a stretch of six consecutive histidine residues, enables purification on metal-chelate resins (QIAGEN). Methods for protein expression and purification are discussed in Ausubel (1995, supra, ch. 10 and 16). Purified PRTS obtained by these methods can be used directly in the assays shown in Examples XVI, XVII, XVIII, and XIX, where applicable.

30 XIII. Functional Assays

PRTS function is assessed by expressing the sequences encoding PRTS at physiologically elevated levels in mammalian cell culture systems. cDNA is subcloned into a mammalian expression vector containing a strong promoter that drives high levels of cDNA expression. Vectors of choice include PCMV SPORT (Life Technologies) and PCR3.1 (Invitrogen, Carlsbad CA), both of which

contain the cytomegalovirus promoter. 5-10 μ g of recombinant vector are transiently transfected into a human cell line, for example, an endothelial or hematopoietic cell line, using either liposome formulations or electroporation. 1-2 μ g of an additional plasmid containing sequences encoding a marker protein are co-transfected. Expression of a marker protein provides a means to distinguish transfected cells from nontransfected cells and is a reliable predictor of cDNA expression from the recombinant vector. Marker proteins of choice include, e.g., Green Fluorescent Protein (GFP; Clontech), CD64, or a CD64-GFP fusion protein. Flow cytometry (FCM), an automated, laser optics-based technique, is used to identify transfected cells expressing GFP or CD64-GFP and to evaluate the apoptotic state of the cells and other cellular properties. FCM detects and quantifies the uptake of fluorescent molecules that diagnose events preceding or coincident with cell death. These events include changes in nuclear DNA content as measured by staining of DNA with propidium iodide; changes in cell size and granularity as measured by forward light scatter and 90 degree side light scatter; down-regulation of DNA synthesis as measured by decrease in bromodeoxyuridine uptake; alterations in expression of cell surface and intracellular proteins as measured by reactivity with specific antibodies; and alterations in plasma membrane composition as measured by the binding of fluorescein-conjugated Annexin V protein to the cell surface. Methods in flow cytometry are discussed in Ormerod, M.G. (1994) Flow Cytometry, Oxford, New York NY.

The influence of PRTS on gene expression can be assessed using highly purified populations of cells transfected with sequences encoding PRTS and either CD64 or CD64-GFP. CD64 and CD64-GFP are expressed on the surface of transfected cells and bind to conserved regions of human immunoglobulin G (IgG). Transfected cells are efficiently separated from nontransfected cells using magnetic beads coated with either human IgG or antibody against CD64 (DYNAL, Lake Success NY). mRNA can be purified from the cells using methods well known by those of skill in the art. Expression of mRNA encoding PRTS and other genes of interest can be analyzed by northern analysis or microarray techniques.

XIV. Production of PRTS Specific Antibodies

PRTS substantially purified using polyacrylamide gel electrophoresis (PAGE; see, e.g., Harrington, M.G. (1990) *Methods Enzymol.* 182:488-495), or other purification techniques, is used to immunize rabbits and to produce antibodies using standard protocols.

Alternatively, the PRTS amino acid sequence is analyzed using LASERGENE software (DNASTAR) to determine regions of high immunogenicity, and a corresponding oligopeptide is synthesized and used to raise antibodies by means known to those of skill in the art. Methods for selection of appropriate epitopes, such as those near the C-terminus or in hydrophilic regions are well described in the art. (See, e.g., Ausubel, 1995, supra, ch. 11.)

Typically, oligopeptides of about 15 residues in length are synthesized using an ABI 431A peptide synthesizer (Applied Biosystems) using FMOC chemistry and coupled to KLH (Sigma-Aldrich, St. Louis MO) by reaction with N-maleimidobenzoyl-N-hydroxysuccinimide ester (MBS) to increase immunogenicity. (See, e.g., Ausubel, 1995, *supra*.) Rabbits are immunized with the

- 5 oligopeptide-KLH complex in complete Freund's adjuvant. Resulting antisera are tested for antipeptide and anti-PRTS activity by, for example, binding the peptide or PRTS to a substrate, blocking with 1% BSA, reacting with rabbit antisera, washing, and reacting with radio-iodinated goat anti-rabbit IgG.

XV. Purification of Naturally Occurring PRTS Using Specific Antibodies

- Naturally occurring or recombinant PRTS is substantially purified by immunoaffinity
10 chromatography using antibodies specific for PRTS. An immunoaffinity column is constructed by covalently coupling anti-PRTS antibody to an activated chromatographic resin, such as CNBr-activated SEPHAROSE (Amersham Pharmacia Biotech). After the coupling, the resin is blocked and washed according to the manufacturer's instructions.

- Media containing PRTS are passed over the immunoaffinity column, and the column is washed
15 under conditions that allow the preferential absorbance of PRTS (e.g., high ionic strength buffers in the presence of detergent). The column is eluted under conditions that disrupt antibody/PRTS binding (e.g., a buffer of pH 2 to pH 3, or a high concentration of a chaotrope, such as urea or thiocyanate ion), and PRTS is collected.

XVI. Identification of Molecules Which Interact with PRTS

- 20 PRTS, or biologically active fragments thereof, are labeled with ¹²⁵I Bolton-Hunter reagent. (See, e.g., Bolton A.E. and W.M. Hunter (1973) *Biochem. J.* 133:529-539.) Candidate molecules previously arrayed in the wells of a multi-well plate are incubated with the labeled PRTS, washed, and any wells with labeled PRTS complex are assayed. Data obtained using different concentrations of PRTS are used to calculate values for the number, affinity, and association of PRTS with the candidate
25 molecules.

Alternatively, molecules interacting with PRTS are analyzed using the yeast two-hybrid system as described in Fields, S. and O. Song (1989) *Nature* 340:245-246, or using commercially available kits based on the two-hybrid system, such as the MATCHMAKER system (Clontech).

- PRTS may also be used in the PATHCALLING process (CuraGen Corp., New Haven CT)
30 which employs the yeast two-hybrid system in a high-throughput manner to determine all interactions between the proteins encoded by two large libraries of genes (Nandabalan, K. et al. (2000) U.S. Patent No. 6,057,101).

XVII. Demonstration of PRTS Activity

Protease activity is measured by the hydrolysis of appropriate synthetic peptide substrates conjugated with various chromogenic molecules in which the degree of hydrolysis is quantified by spectrophotometric (or fluorometric) absorption of the released chromophore (Beynon, R.J. and J.S. Bond (1994) Proteolytic Enzymes: A Practical Approach, Oxford University Press, New York NY, pp.25-55). Peptide substrates are designed according to the category of protease activity as endopeptidase (serine, cysteine, aspartic proteases, or metalloproteases), aminopeptidase (leucine aminopeptidase), or carboxypeptidase (carboxypeptidases A and B, procollagen C-proteinase). Commonly used chromogens are 2-naphthylamine, 4-nitroaniline, and furylacrylic acid. For example, arginine- β -naphthylamide can be used as a substrate for SEQ ID NO:3 (Fukasawa, K.M. et al. (1996) J. Biol. Chem. 271:30731-30735) and 4-phenylazobenzoyloxycarbonyl-Pro-Leu-Gly-Pro-D-Arg can be used as a substrate for SEQ ID NO:4. In an alternative example, a substrate for SEQ ID NO:9 would be 7-amino-4-trifluoromethyl coumarin-Phe-Pro-AFC. Assays are performed at ambient temperature and contain an aliquot of the enzyme and the appropriate substrate in a suitable buffer. Reactions are carried out in an optical cuvette, and the increase/decrease in absorbance of the chromogen released during hydrolysis of the peptide substrate is measured. The change in absorbance is proportional to the enzyme activity in the assay.

An alternate assay for ubiquitin hydrolase activity measures the hydrolysis of a ubiquitin precursor. The assay is performed at ambient temperature and contains an aliquot of PRTS and the appropriate substrate in a suitable buffer. For SEQ ID NO:1, chemically synthesized human ubiquitin-valine may be used as substrate. Cleavage of the C-terminal valine residue from the substrate is monitored by capillary electrophoresis (Franklin, K. et al. (1997) Anal. Biochem. 247:305-309).

Alternatively, the ubiquitin protease activity of SEQ ID NO:5 can be measured using the method of Sloper-Mould et al. ((1999) J. Biol. Chem. 274:26878-26884). Aliquots of PRTS are incubated with 5 μ l [35 S]-labeled ubiquitin-GST fusion substrate for 1 hour at 37 °C in an appropriate buffer. Samples are resolved by electrophoresis on a 12% SDS-PAGE gel. Ubiquitin cleavage is monitored by fluorography of the gel.

Alternatively, the activity of SEQ ID NO:10, for example, can be measured by the method of Colige et al. (1999, J. Biol. Chem. 270:16724-16730). An aliquot of PRTS is incubated with amino procollagen type I substrate radioactively labeled only in the propeptide in a 250 μ l reaction containing 50 mM sodium cacodylate, pH 7.5, 200 mM KCl, 2 mM CaCl₂, 2.5 mM NEM, 0.5 mM PMSF, and 0.02% Brij (standard assay buffer). After 16 h at 26 °C, the reaction is stopped by adding 50 μ l of EDTA solution (0.2 M EDTA, pH 8, 0.5% SDS, 0.5 M DTT) and 300 μ l of 99% ethanol. The samples are kept for 30 min at 4 °C and centrifuged for 30 min at 9500 g. Collagen and uncleaved

radioactive pN-collagen substrate are pelleted, whereas the freed amino propeptides remained in solution. An aliquot of the supernatant is assayed by liquid scintillation spectrometry.

In the alternative, an assay for protease activity takes advantage of fluorescence resonance energy transfer (FRET) that occurs when one donor and one acceptor fluorophore with an appropriate spectral overlap are in close proximity. A flexible peptide linker containing a cleavage site specific for PRTS is fused between a red-shifted variant (RSGFP4) and a blue variant (BFP5) of Green Fluorescent Protein. This fusion protein has spectral properties that suggest energy transfer is occurring from BFP5 to RSGFP4. When the fusion protein is incubated with PRTS, the substrate is cleaved, and the two fluorescent proteins dissociate. This is accompanied by a marked decrease in energy transfer which is quantified by comparing the emission spectra before and after the addition of PRTS (Mitra, R.D. et al. (1996) *Gene* 173:13-17). This assay can also be performed in living cells. In this case the fluorescent substrate protein is expressed constitutively in cells and PRTS is introduced on an inducible vector so that FRET can be monitored in the presence and absence of PRTS (Sagot, I. et al. (1999) *FEBS Lett.* 447:53-57).

In yet another alternative, an assay for PRTS dipeptidase activity measures the hydrolysis activity of PRTS on a variety of dipeptides such as leukotriene D4 (Kozak, E. and S. Tate (1982) *J. Biol. Chem.* 257:6322-6327), or hydrolysis of the beta-lactam ring of antibiotics such as penum and carbapenem (Campbell et al., (1984) *J. Biol. Chem.* 259:14586-14590).

XVIII. Identification of PRTS Substrates

Phage display libraries can be used to identify optimal substrate sequences for PRTS. A random hexamer followed by a linker and a known antibody epitope is cloned as an N-terminal extension of gene III in a filamentous phage library. Gene III codes for a coat protein, and the epitope will be displayed on the surface of each phage particle. The library is incubated with PRTS under proteolytic conditions so that the epitope will be removed if the hexamer codes for a PRTS cleavage site. An antibody that recognizes the epitope is added along with immobilized protein A. Uncleaved phage, which still bear the epitope, are removed by centrifugation. Phage in the supernatant are then amplified and undergo several more rounds of screening. Individual phage clones are then isolated and sequenced. Reaction kinetics for these peptide substrates can be studied using an assay in Example XVII, and an optimal cleavage sequence can be derived (Ke, S.H. et al. (1997) *J. Biol. Chem.* 272:16603-16609).

To screen for *in vivo* PRTS substrates, this method can be expanded to screen a cDNA expression library displayed on the surface of phage particles (T7SELECT 10-3 Phage display vector, Novagen, Madison WI) or yeast cells (pYD1 yeast display vector kit, Invitrogen, Carlsbad CA). In this case, entire cDNAs are fused between Gene III and the appropriate epitope.

XIX. Identification of PRTS Inhibitors

Compounds to be tested are arrayed in the wells of a multi-well plate in varying concentrations along with an appropriate buffer and substrate, as described in the assays in Example XVII. PRTS activity is measured for each well and the ability of each compound to inhibit PRTS activity can be
5 determined, as well as the dose-response kinetics. This assay could also be used to identify molecules which enhance PRTS activity.

In the alternative, phage display libraries can be used to screen for peptide PRTS-inhibitors. Candidates are found among peptides which bind tightly to a protease. In this case, multi-well plate wells are coated with PRTS and incubated with a random peptide phage display library or a cyclic
10 peptide library (Koivunen, E. et al. (1999) Nat. Biotechnol. 17:768-774). Unbound phage are washed away and selected phage amplified and rescreened for several more rounds. Candidates are tested for PRTS inhibitory activity using an assay described in Example XVII.

Various modifications and variations of the described methods and systems of the invention will
15 be apparent to those skilled in the art without departing from the scope and spirit of the invention. Although the invention has been described in connection with certain embodiments, it should be understood that the invention as claimed should not be unduly limited to such specific embodiments. Indeed, various modifications of the described modes for carrying out the invention which are obvious to those skilled in molecular biology or related fields are intended to be within the scope of the following
20 claims.

Table 1

Incyte Project ID	Polypeptide SEQ ID NO:	Incyte Polypeptide ID	Polynucleotide SEQ ID NO:	Incyte Polynucleotide ID
275791	1	275791CD1	22	275791CB1
1389845	2	1389845CD1	23	1389845CB1
1726609	3	1726609CD1	24	1726609CB1
4503848	4	4503848CD1	25	4503848CB1
5544089	5	5544089CD1	26	5544089CB1
7474081	6	7474081CD1	27	7474081CB1
5281209	7	5281209CD1	28	5281209CB1
2256251	8	2256251CD1	29	2256251CB1
7160544	9	7160544CD1	30	7160544CB1
7477386	10	7477386CD1	31	7477386CB1
7473089	11	7473089CD1	32	7473089CB1
7604035	12	7604035CD1	33	7604035CB1
3473847	13	3473847CD1	34	3473847CB1
3750004	14	3750004CD1	35	3750004CB1
4904126	15	4904126CD1	36	4904126CB1
71268415	16	71268415CD1	37	71268415CB1
7473301	17	7473301CD1	38	7473301CB1
7473308	18	7473308CD1	39	7473308CB1
7478021	19	7478021CD1	40	7478021CB1
4333459	20	4333459CD1	41	4333459CB1
6817347	21	6817347CD1	42	6817347CB1

Table 2

Polypeptide SEQ ID NO:	Incyte Polypeptide ID	GenBank ID NO:	Probability score	GenBank Homolog
1	275791CD1	g9971757	1.00E-49	ubiquitin-specific processing protease [Homo sapiens]
2	1389845CD1	g1143194	1.30E-46	proctasin [Homo sapiens]
3	1726609CD1	g10719660	0	RNPEP-like protein [Homo sapiens] (Horikawa, Y. et al. (2000) Nat. Genet. 26:163-175.)
		g1754515	3.30E-96	aminopeptidase-B [Rattus norvegicus] (Prieto, I. et al. (1998) Horm. Metab. Res. 30:246-248.)
4	4503848CD1	g1783122	0	endopeptidase 24.16 type M1 [Sus scrofa]
5	5544089CD1	g5410230	5.20E-43	ubiquitin-specific protease 3 [Homo sapiens]
6	7474081CD1	g603903	2.90E-33	Trypsinogen [Gallus gallus]
7	5281209CD1	g11071729	0	putative dipeptidase [Homo sapiens]
		g219600	1.40E-88	dipeptidase precursor [Homo sapiens] (Sato, S. et al. (1993) Biochim. Biophys. Acta 1172:181-183.)
8	2256251CD1	g6103629	3.90E-166	transmembrane tryptase [Homo sapiens] (Wong, G.W. et al. (1999) J. Biol. Chem. 274:30784-30793.)
9	7160544CD1	g11095188	0	dipeptidyl peptidase 8 [Homo sapiens] (Abbott, C.A. et al. (2000) Eur. J. Biochem. 267:6140-6150.)
		g1753197	6.80E-64	dipeptidyl peptidase IV [Stenotrophomonas maltophilia] (Mentlein, R. (1999) Regul. Pept. 85:9-24; Kahne, T. Int. J. Mol. Med (1999) 4:3-15.)
10	7477386CD1	g1865716	0	procollagen I N-proteinase [Bos taurus] (Colige, A. et al. (1999) Am. J. Hum. Genet. 65:308-317.)
11	7473089CD1	g7768706	3.60E-255	metalloprotease with thrombospondin type 1 motifs [Homo sapiens] (Vazquez, F. et al. (1999) J. Biol. Chem. 274:23349-23357.)
12	7604035CD1	g6164595	4.70E-68	Lacunin [Manduca sexta]

Table 2 (cont.)

Polypeptide SEQ ID NO:	Incyte Polypeptide ID	GenBank ID NO:	Probability score	GenBank Homolog
13	3473847CD1	g217172	9.20E-50	aqualysin precursor (aa 1 to 513) [Thermus aquaticus]
14	3750004CD1	g5923786	4.30E-51	zinc metalloprotease ADAMTS6 [Homo sapiens]
15	4904126CD1	g186286	3.90E-40	interleukin 1-beta convertase [Homo sapiens] (Cerretti, D.P. et al. (1992) Science 256:97-100.)
16	71268415CD1	g6651071	0	disintegrin and metalloprotease domain 19 [Homo sapiens] (Inoue, D. et al. (1998) J. Biol. Chem. 273:4180-4187.)
17	7473301CD1	g6002714	5.10E-94	membrane-type serine protease 1 [Homo sapiens] (Takeuchi, T. et al. (1999) Proc. Natl. Acad. Sci. USA 96:11054-11061.)
18	7473308CD1	g1552517	6.60E-77	trypsinogen E [Homo sapiens]
19	7478021CD1	g3211705	5.60E-189	matrix metalloprotease [Xenopus laevis] (Yang, M. (1997) J. Biol. Chem. 272:13527-13533.)
20	4333459CD1	g1754714	2.30E-67	Oviductin [Xenopus laevis] (Lindsay, L.L. et al. (1999) Biol. Reprod. 60:989-995.)
21	6817347CD1	g7673618	5.10E-283	ubiquitin specific protease [Mus musculus]

Table 3

SEQ ID NO:	Incyte Polypeptide ID	Amino Acid Residues	Potential Phosphorylation Sites	Potential Glycosylation Sites	Signature Sequences, Domains and Motifs	Analytical Methods and Databases
1	275791CD1	232	T15 T17 S23 S43 S71 T90 S93 S100 S107 S111 T122 T174 S190 T10 S141 S190 T213 T227	N98 N99	Ubiquitin carboxyl-terminal hydrolase family 2 signature 2 Uch_2: Y142-Y160 Ubiquitin carboxyl-terminal hydrolase family 2 signature 2 UCH-2: L138-H203 Ubiquitin carboxyl-terminal hydrolase family 2 signature 2 BL00972: Y142-D166, K169-S190	MOTIFS HMMER-PFAM BLIMPS-BLOCKS
2	1389845CD1	365	S120 S187 S225 S253 S82 T31 T37 T42 Y283		TRYPSIN DM00018 A57014 45-284: I123-Q314 PROTEASE SERINE PRECURSOR SIGNAL HYDROLASE ZYMOGEN GLYCOPROTEIN FAMILY MULTIGENE FACTOR PD000046: I123-Q314 Serine proteases, trypsin family BL00134A: C148-C164 Kringle domain proteins BL00021: C148-F165, V231-G252 CHYMOTRYPSIN SERINE PROTEASE PR00722: G149-C164, G208-V222 V8 Serine proteases PR00839B: C148-F165 Serine proteases, trypsin family, active sites trypsin.his.prf: I140-P188 Trypsin: I123-Q314 Trypsin_His: L159-C164	BLAST_DOMO BLAST_PRODOR BLIMPS_BLOCKS BLIMPS_BLOCKS BLIMPS_PRINTS BLIMPS_PRINTS PROFILES SCAN HMMER-PFAM MOTIFS

Table 3 (cont.)

SEQ ID NO:	Incyte Polypeptide ID	Amino Acid Residues	Potential Phosphorylation Sites	Potential Glycosylation Sites	Signature Sequences, Domains and Motifs	Analytical Methods and Databases
3	1726609CD1	416	S244 S283 S30 S370 S408 T389 T59 T78 T87	N203 N413 N57	do HYDROLASE; LEUKOTRIENE; A-4; ZINC; DM08707 P19602 7-609: M1-P354 AMINOPEPTIDASE HYDROLASE METALLOPROTEASE ZINC N GLYCOPROTEIN PROTEIN TRANSMEMBRANE SIGNAL ANCHOR MEMBRANE PD001134: R4-S177 Neutral zinc metallopeptidase family BL00142: D41-F51 MEMBRANE ALANYL DIPEPTIDASE PR00756: F14-L24, D41-T56, W60-Y72 signal_cleavage: M1-S34	BLAST_DOMO BLAST_PRODUM BLIMPS_BLOCKS BLIMPS_PRINTS SPScan
4	4503848CD1	714	S124 S140 S147 S179 S200 S206 S226 S333 S551 S556 S592 T114 T133 T244 T252 T270 T308 T318 T322 T376 T406 T432 T528 T585 T602 T69 Y175 Y249 Y505	N425 N485 N601	do ZINC; METALLOPEPTIDASE; NEUTRAL; OLIGO-PEPTIDASE DM01184 Q02038 36-702: A46-A713 HYDROLASE METALLOPROTEASE ZINC OLIGOPEPTIDASE PRECURSOR MITOCHONDRIAL ENDOPEPTIDASE MITOCHONDRION TRANSIT PEPTIDE PD002945: W60-N529 transmembrane domain: L14-M34 Peptidase family M3 Peptidase_M3: C98- L711 Neutral zinc metallopeptidase family BL00142: T504-H514 Zinc_Protease: T504-M513 signal_cleavage: M1-G27	BLAST_PRODUM HMMER HMMER_PFAM BLIMPS_BLOCKS MOTIFS SPScan

Table 3 (cont.)

SEQ ID NO:	Incyte Polypeptide ID	Amino Acid Residues	Potential Phosphorylation Sites	Potential Glycosylation Sites	Signature Sequences, Domains and Motifs	Analytical Methods and Databases
5	5544089CD1	367	S108 S161 S197 S203 S235 S266 S361 S49 T180 T263 T316 T331	N139 N142 N308	UBIQUITIN CARBOXYL-TERMINAL HYDROLASES FAMILY 2 DM00659 P40818 782-1103: L36-L328 Ubiquitin carboxyl-terminal hydrolase family BL00972: Y74-L83, V120-C134, Y274-N298, G301-K322 Ubiquitin carboxyl-terminal hydrolase family UCH-2: E270-Q332 Ubiquitin carboxyl-terminal hydrolase family Uch_2_2: Y274-Y292 signal_cleavage: M1-S16	BLAST_DOMO BLIMPS_BLOCKS HMMER_PFAM MOTIFS SPScan
6	7474081CD1	235	S134 S143 S159 S171 S226 T231	N90	TRYPSIN DM00018 S55065 26-244: A27-T231 Serine proteases, trypsin family signature BL00134: C40-C56, V215-I228 Type I fibronectin domain BL01253: C40-P53, I197-T231 Kringle domain proteins BL00021: S159-Q164, C40-Y57 CHYMOTRYPSIN SERINE PROTEASE PR00722A: V41-C56, A95-A109 Serine proteases, trypsin family, active site trypsin_his.prf: S35-T76 Trypsin: G42-V178, G216-I228 Leucine_Zipper: L44-L65 signal_cleavage: M1-S19	BLAST_DOMO BLIMPS_BLOCKS BLIMPS_BLOCKS BLIMPS_BLOCKS BLIMPS_BLOCKS BLIMPS_PRINTS PROFILES SCAN HMMER_PFAM MOTIFS SPScan

Table 3 (cont.)

SEQ ID NO:	Incyte Polypeptide ID	Amino Acid Residues	Potential Phosphorylation Sites	Potential Glycosylation Sites	Signature Sequences, Domains and Motifs	Analytical Methods and Databases
7	5281209CD1	488	S13 T74 S186 S233 T363 T456 T4 T34 T125 S170 S172 T178 S249 T337 S387 S389 S419 T447	N119 N184 N243 N334	Renal dipeptidase proteins BL00869: P92-L247, E280-R412, S415-N457 DIPEPTIDASE MICROSOMAL PRECURSOR MDP HYDROLASE MICROSOME SIGNAL GPI-ANCHOR GLYCOPROTEIN ZINC PD005626: S143-E450 RENAL DIPEPTIDASE DM02775: T77-K410 Renal dipeptidase: V195-R217 Renal dipeptidase: A63-V475 Signal peptide: M1-A36 Signal cleavage: M1-A36	BLIMPS-BLOCKS BLAST-PRODROM BLAST-DOMO MOTIFS HMMER-PFAM HMMER SPSCAN
8	2256251CD1	346	S203 S210 S266 S45 S79 T131 T147 T216	N110	TRYPSIN DM00018 P15944 31-270: I63-I294 PROTEASE SERINE PRECURSOR SIGNAL HYDROLASE ZYMOGEN GLYCOPROTEIN FAMILY MULTIGENE FACTOR PD000046: L156-I290, I63-F178, N288-F314, P274-P305 Serine proteases, trypsin family BL00134: C88-C104, D241-I264, P277-I290 Type I fibronectin domain BL01253: C88-A101, V158-E194, G240-C253, W259-H293 Kringle domain proteins BL00021: C88-F105, V169-G190, G249-I290 CHYMOTRYPSIN SERINE PROTEASE PR00722: G89-C104, G146-V160, G240-V252 transmembrane domain: P308-L328 trypsin: I63-I290 Trypsin_His L99-C104 Trypsin family serine protease active sites trypsin_his.prf: L80-H128 trypsin_ser.prf: L229-R273 signal_cleavage: M1-S45	BLIMPS_BLOCKS BLIMPS_BLOCKS BLIMPS_BLOCKS BLIMPS_PRINTS HMMER HMMER_PFAM MOTIFS PROFILES SCAN SPSCAN

Table 3 (cont.)

SEQ ID NO:	Incyte Polypeptide ID	Amino Acid Residues	Potential Phosphorylation Sites	Potential Glycosylation Sites	Signature Sequences, Domains and Motifs	Analytical Methods and Databases
9	7160544CD1	882	S115 S133 S293 S312 S412 S443 S479 S530 S587 S588 S723 S80 S850 T227 T234 T307 T326 T499 T52 T551 T594 T603 T615 T776 Y315 Y36 Y55 Y555 Y844		PROLYL ENDOPEPTIDASE FAMILY SERINE DM02461 P27487 192-765: F488-E870, G251-E370 DIPEPTIDYL IV HYDROLASE PROTEASE SERINE PEPTIDASE DIPEPTIDASE TRANSMEMBRANE GLYCOPROTEIN PROTEIN PD003048: L744-E870 DIPEPTIDYL IV HYDROLASE PROTEASE SERINE PEPTIDASE DIPEPTIDASE TRANSMEMBRANE GLYCOPROTEIN PROTEIN PD003086: Y423- V661, I212-T326 Prolyl endopeptidase family BL00708: G501-I513, Q714-L744 Dipeptidyl peptidase IV PF00930: I498- R508, F756-P783, R808-L828 PROLYL OLIGOPEPTIDASE SERINE PROTEASE PR00862: P647-P665, G737-R757 Prolyl oligopeptidase family Peptidase_S9: R672-L744 Dipeptidyl peptidase IV DPPIV_N_term: M88-N663	BLAST_DOMO BLAST_PRODUM BLAST_PRODUM BLIMPS_BLOCKS BLIMPS_PFAM BLIMPS_PRINTS HMMER_PFAM HMMER_PFAM

Table 3 (cont.)

SEQ ID NO:	Incyte Polypeptide ID	Amino Acid Residues	Potential Phosphorylation Sites	Potential Glycosylation Sites	Signature Sequences, Domains and Motifs	Analytical Methods and Databases
10	7477386CD1	1189	S132 S169 S200 S32 S323 S350 S445 S480 S511 S626 S675 S699 S798 S1064 T247 T362 T521 T612 T718 T777 T946 T986 T1104 Y552	N109 N478 N944	do ZINC; METALLOPEPTIDASE; NEUTRAL; ATROLYSIN DM00368 Q05910 189-395:I261-P463 THROMBOSPONDIN TYPE 1 REPEAT DM00275 P07996 477-540: D555-C604 PROTEIN PROCOLLAGEN THROMBOSPONDIN MOTIFS NPROTEINASE C02B4.1 A DISINTEGRIN METALLOPROTEASE PD013511: L474-E549 PROTEIN PROCOLLAGEN THROMBOSPONDIN: MOTIFS NPROTEINASE A DISINTEGRIN METALLOPROTEASE WITH ADAMTS1 PD011654: Q647-C716 PROCOLLAGEN C37C3.6 SERINE PROTEASE INHIBITOR PD007018: W854-Q974, W914-C1029, W558-K623 METALLOPROTEASE PRECURSOR HYDROLASE SIGNAL ZINC VENOM CELL PROTEIN TRANSMEMBRANE ADHESION PD000791:P256-P463 Neutral zinc metallopeptidase BL00142: T398-N408 signal_peptide: M1-A22 Reprolysin family propeptide Pep_M12B_propep: R120-V240 Reprolysin (M12B) family zinc metalloprotease Reprolysin: I261-P463 Thrombospondin type 1 domain tsp_1: A973-C1024, S559-C609, Y852-C909, W914-C971 signal_cleavage: M1-G23	BLAST_DOMO BLAST_DOMO BLAST_PRODOM BLAST_PRODOM BLAST_PRODOM BLAST_PRODOM BLIMPS_BLOCKS HMMER HMMER_PFAM HMMER_PFAM HMMER_PFAM

Table 3 (cont.)

SEQ ID NO:	Incyte Polypeptide ID	Amino Acid Residues	Potential Phosphorylation Sites	Potential Glycosylation Sites	Signature Sequences, Domains and Motifs	Analytical Methods and Databases
11	7473089CD1	952	S19 S203 S207 S303 S346 S432 S492 S575 S578 S611 S666 S682 S708 S745 S919 T171 T288 T317 T325 T337 T359 T471 T594 T687 T765	N141 N591 N623 N680	do ZINC; METALLOPEPTIDASE; NEUTRAL; ATROLYSIN DMO0368 JC2550 1-201:R218-P427 PROTEIN PROCOLLAGEN THROMBOSPONDIN MOTIFS NPROTEINASE A DISINTEGRIN METALLOPROTEASE WITH ADAMTS1 PD014161: K684-E804 PROTEIN PROCOLLAGEN THROMBOSPONDIN MOTIFS NPROTEINASE A DISINTEGRIN METALLOPROTEASE WITH ADAMTS1 PD011654: V610-C683 METALLOPROTEASE PRECURSOR HYDROLASE SIGNAL ZINC VENOM CELL PROTEIN TRANSMEMBRANE ADHESION PD000791: V214-P427 PROTEIN PROCOLLAGEN THROMBOSPONDIN MOTIFS NPROTEINASE C02B4.1 A DISINTEGRIN METALLOPROTEASE PD013511: L437-V505 Neutral zinc metalloproteinase BL00142: T358-N368 signal_peptide: M1-G18 Reprolysin family propeptide Pep_M12B_propep: H67-N181 Reprolysin (M12B) family zinc metalloproteinase Reprolysin: R218-P427 Thrombospondin type 1 domain tsp_1: A520-C570, W845-C896, W899-C952 Zinc_Protease: T358-F367 SpScan signal_cleavage: M1-G17	BLAST_DOMO BLAST_PRODROM BLAST_PRODROM BLAST_PRODROM BLAST_PRODROM BLAST_PRODROM BLIMPS_BLOCKS HMMER HMMER_FFAM HMMER_FFAM HMMER_FFAM MOTIFS SPSCAN

Table 3 (cont.)

SEQ ID NO:	Incyte Polypeptide ID	Amino Acid Residues	Potential Phosphorylation Sites	Potential Glycosylation Sites	Signature Sequences, Domains and Motifs	Analytical Methods and Databases
12	7604035CD1	898	S187 S188 S258 S268 S285 S415 S467 S547 S696 S796 S819 S851 S892 T106 T198 T35 T434 T483 T492 T5	N3 N490 N773	PROCOLLAGEN C37C3.6 SERINE PROTEASE INHIBITOR ALTERNATIVE PD007018: Y726-C841, W786-A874, Y667-C778, W50-Q72, S368-Q383 PROTEIN PROCOLLAGEN THROMBOSPONDIN MOTIFS NPROTEINASE A DISINTEGRIN METALLOPROTEASE WITH ADAMTS1 PD011654: Q416-C484 PROTEIN PROCOLLAGEN THROMBOSPONDIN MOTIFS NPROTEINASE A DISINTEGRIN METALLOPROTEASE WITH ADAMTS1 PD014161: R485-I599 signal_peptide: M1-D24 Thrombospondin type 1 domain tsp_1: G48-R87, W727-C783, E787-C841 signal_cleavage: M1-D24	BLAST_PRODUM BLAST_PRODUM
13	3473847CD1	631	S117 S160 S174 S185 S188 S268 S28 S30 S358 S431 S503 S605 T142 T33 T346 T512 T606	N472	Subtilase family Peptidase_S8: S86-N364 SERINE PROTEASES, SUBTILASE FAMILY, HISTIDINE DM00108 P80146 150-377: G116-T346 Serine proteases, subtilase family BL00136: L123-I135, D163-G175, G323-G333 SUBTILISIN SERINE PROTEASE FAMILY PR00723: G116-I135, K161-S174, S322-M338 Serine proteases, subtilase family, active site subtilase_ser.prf: A302-Q352	SPSCAN HMMER_PFAM BLAST_DOMO BLIMPS_BLOCKS BLIMPS_PRINTS PROFILESCAN

Table 3 (cont.)

SEQ ID NO:	Incyte Polypeptide ID	Amino Acid Residues	Potential Phosphorylation Sites	Potential Glycosylation Sites	Signature Sequences, Domains and Motifs	Analytical Methods and Databases
14	3750004CD1	470	S454 S51 S54 T104 T276 T386 T464	N182 N203	Thrombospondin type 1 domain tsp_1: T34-C81 PROTEIN PROCOLLAGEN THROMBOSPONDIN MOTIFS NPROTEINASE A DISINTEGRIN METALLOPROTEASE WITH ADAMTS1 PD011654: Q119-C185 signal_peptide: M1-G29 signal_cleavage: M1-G24	HMMER_PFAM BLAST_PRODOR
15	4904126CD1	110	S16 S36 T100 T49 N47		Caspase recruitment domain CARD: A2-A91 INTERLEUKIN-1 BETA CONVERTING ENZYME FAMILY HISTIDINE DM07463 P29466 1-122: M1-S110	HMMER SPScan HMMER_PFAM BLAST_DOMO

Table 3 (cont.)

SEQ ID NO:	Incyte Polypeptide ID	Amino Acid Residues	Potential Phosphorylation Sites	Potential Glycosylation Sites	Signature Sequences, Domains and Motifs	Analytical Methods and Databases
16	71268415CD1	879	S132 S14 S208 S288 S571 S711 S747 S754 S755 S827 T106 T118 T29 T30 T373 T412 T42 T428 T444 T55 T688 Y167 Y39	N368 N371 N569 N68	Reprolysin family propeptide Pep_M12B_propep: R8-K119 Reprolysin (M12B) family zinc metalloprotease Reprolysin: K134-P332 Disintegrin: E349-Q424 do-ZINC; METALLOPEPTIDASE; NEUTRAL; ATROLYSIN; DM00368 S60257 204-414: K126-D333 do ZINC; REGULATED; EPIDIDYMAL; NEUTRAL; DM00591 S60257 492-628: F410-L549 MELTRIN, BETA METALLOPROTEASE DISINTEGRIN BETA INTEGRIN PROTEASE METALLOPROTEASE PD105322: P620-G812 METALLOPROTEASE PRECURSOR HYDROLASE SIGNAL ZINC VENOM CELL PROTEIN TRANSMEMBRANE ADHESION PD000791: K134-P332 MELTRIN, BETA METALLOPROTEASE DISINTEGRIN MELTRIN BETA INTEGRIN PROTEASE METALLOPROTEASE PD171676: K495-C567 CELL ADHESION PLATELET BLOOD COAGULATION VENOM DISINTEGRIN METALLOPROTEASE PRECURSOR SIGNAL PD000664: E349-Y423 Neutral zinc metalloprotease BL00142: T266-G276 DISINTEGRIN SIGNATURE PR00289: C380-R399, E409-N421 NEPRILYSIN METALLOPROTEASE PR00786C: N259-F275	HMIMER_PPFAM HMIMER_PPFAM HMIMER_PPFAM BLAST_DOMO BLAST_PRODOM BLAST_PRODOM BLAST_PRODOM BLIMPS_BLOCKS BLIMPS_PRINTS BLIMPS_PRINTS

Table 3 (cont.)

SEQ ID NO:	Incyte Polypeptide ID	Amino Acid Residues	Potential Phosphorylation Sites	Potential Glycosylation Sites	Signature Sequences, Domains and Motifs	Analytical Methods and Databases
					Disintegrins signature disintegrins.prf: E360-P419	PROFILESCAN
					Neutral zinc metalloproteinases, zinc-binding region signature	PROFILESCAN
					zinc_protease.prf: S249-G301	
					transmembrane domain: V624-Y645	HMME
					Zn binding region Zinc_Protease: T266-F275	MOTIFS
17	7473301CD1	850	S100 S275 S295 S358 S429 S448 S470 S474 S495 S536 S596 S64 S787 S802 S807 T117 T250 T312 T348 T382 T404 T426 T570 T714 T777	N19 N210 N422 N486 N533 N559 N568	TRYPSIN FAMILY SERINE PROTEASE trypsin: I613-I842 Low-density lipoprotein receptor domain ldl_recept_a: Q489-S527, P530-Q562, I564-C603 TRYPSIN DM00018 P98072 800-1033: R612-V846 PROTEASE SERINE PRECURSOR SIGNAL HYDROLASE ZMOGEN GLYCOPROTEIN FAMILY MULTIGENE FACTOR PD000046: Q675-I842, I613-G809 Serine proteases, trypsin family BL00134: C638-C654, D791-T814, P829-I842 Type I fibronectin domain BL01253: C638-A651, R790-C803, W811-Y845 Kringle domain proteins BL00021: I722-G743, L801-I842, C638-F655 LOW DENSITY LIPOPROTEIN RECEPTOR DOMAIN PR00261: G501-E522 CHYMOTRYPSIN SERINE PROTEASE PR00722: G639-C654, T697-W711, R790-S802	HMME_PPFAM HMME_PPFAM BLAST_DOMO BLAST_PRODOM BLIMPS_BLOCKS BLIMPS_BLOCKS BLIMPS_BLOCKS BLIMPS_PRINTS BLIMPS_PRINTS

Table 3 (cont.)

SEQ ID NO:	Incyte Polypeptide ID	Amino Acid Residues	Potential Phosphorylation Sites	Potential Glycosylation Sites	Signature Sequences, Domains and Motifs	Analytical Methods and Databases
					Trypsin family serine protease active sites trypsin_his.prf: L630-K679 trypsin_ser.prf: I776-R825 transmembrane domain: I77-L95	PROFILES CAN
					Trypsin family serine protease active sites Trypsin_His L649-C654 Trypsin_Ser D791-S802	HMER MOTIFS
18	7473308CD1	254	S136 S14 S153 S195 S227 T230 T249		TRYPSIN FAMILY SERINE PROTEASE trypsin: I21-Q183 CHYMOTRYPSIN SERINE PROTEASE FAMILY PRO0722B: T89-A103 TRYPSIN DM00018 P07478 24-242: I21-Q183 PROTEASE SERINE PRECURSOR SIGNAL HYDROLASE ZMOGEN GLYCOPROTEIN FAMILY MULTIGENE FACTOR PD000046: G23-Q183	HMER_PFAM BLIMPS_PRINTS BLAST_DOMO BLAST_PRODUM

Table 3 (cont.)

SEQ ID NO:	Incyte Polypeptide ID	Amino Acid Residues	Potential Phosphorylation Sites	Potential Glycosylation Sites	Signature Sequences, Domains and Motifs	Analytical Methods and Databases
19	7478021CD1	568	S142 S145 S153 S172 S177 S190 S244 S316 S34 S420 S448 S552 T209 T22 T293 T334 T401 T427 T489 T79 Y509	N371	<p>Matrixin Peptidase M10: F56-T266</p> <p>Hemopexin domain: F332-T390, I393-S448, L450-Q498, I505-K548</p> <p>MATRIXINS CYSTEINE SWITCH</p> <p>DM00558 P22757 100-337: A184-T334, P76-P124</p> <p>MATRIXINS CYSTEINE SWITCH</p> <p>DM00558 P08254 29-274: Q158-T334, L85-M122</p> <p>MATRIX METALLOPROTEINASE PD168921: S327-N392</p> <p>MATRIX METALLOPROTEINASE PD169970: A494-M568</p> <p>MATRIX PRECURSOR METALLOPROTEASE</p> <p>HYDROLASE ZINC ZYMOGEN CALCIUM COLLAGEN DEGRADATION SIGNAL PD000673: F171-T266, P73-M122</p> <p>Matrixins cysteine switch BL00546: F92-D121, V224-P267, G273-Y304, L313-G326, F443-Y455, F409-E428</p> <p>Hemopexin domain protein BL00024: M112-M122, G273-Y304, L313-G326, F443-Y455, Y408-D419</p> <p>MATRIXIN SIGNATURE PR00138: M112-P125, E198-F213, V224-W252, V279-Y304, L313-G326</p> <p>Matrixins cysteine switch</p> <p>cysteine_switch.prf: A95-M204</p> <p>Neutral zinc metalloproteinases, zinc-binding region signature</p> <p>zinc_protease.prf: D256-E312</p>	<p>HMIMER_PFAM</p> <p>HMIMER_PFAM</p> <p>BLAST_DOMO</p> <p>BLAST_DOMO</p> <p>BLAST_PRODOM</p> <p>BLAST_PRODOM</p> <p>BLAST_PRODOM</p> <p>BLIMPS_BLOCKS</p> <p>BLIMPS_BLOCKS</p> <p>BLIMPS_PRINTS</p> <p>PROFILES SCAN</p> <p>PROFILES SCAN</p>

Table 3 (cont.)

SEQ ID NO:	Incyte Polypeptide ID	Amino Acid Residues	Potential Phosphorylation Sites	Potential Glycosylation Sites	Signature Sequences, Domains and Motifs	Analytical Methods and Databases
					Hemopexin domain signature hemopexin.prf: F409-R477 signal_peptide: M1-P21 Zn binding region Zinc_Protease: V279-L288 signal_cleavage: M1-P24	PROFILES SCAN HMMER MOTIFS
20	4333459CD1	306	S117 S138 S2 S223 S60 S72 T110 T139 T207 T217	N108	TRYPSIN FAMILY SERINE PROTEASE trypsin: I56-I298 TRYPSIN DM00018 Q05319 543-784: I56-I302 PROTEASE SERINE PRECURSOR SIGNAL HYDROLASE ZYMOGEN GLYCOPROTEIN FAMILY MULTIGENE FACTOR PD000046: S117-I298, I56-G192 Serine proteases, trypsin family BL00134: C81-C97, D238-G261, P285-I298 Type I fibronectin domain BL01253: C81-A94, G154-E190, R237-C250 Kring domain proteins BL00021: C81-I98, I165-G186, S247-E288 CHYMOTRYPSIN SERINE PROTEASE PR00722: G82-C97, P142-F156, R237-M249 Trypsin family serine protease active sites trypsin_his.prf: L73-G122 trypsin_ser.prf: K225-R271 Trypsin family serine protease active sites Trypsin_His: I92-C97 Trypsin_Ser: D238-M249 signal_cleavage: M1-S26	SPSCAN HMMER_PFAAM BLAST_DOMO BLAST_PRODOR BLIMPS_BLOCKS BLIMPS_BLOCKS BLIMPS_BLOCKS BLIMPS_PRINTS PROFILES SCAN MOTIFS SPSCAN

Table 3 (cont.)

SEQ ID NO:	Incyte Polypeptide ID	Amino Acid Residues	Potential Phosphorylation Sites	Potential Glycosylation Sites	Signature Sequences, Domains and Motifs	Analytical Methods and Databases
21	6817347CD1	953	S102 S114 S150 S172 S369 S429 S47 S623 S794 S804 S808 S831 S856 S919 S942 T289 T42 T455 T488 T544 T567 T568 T585 T59 T736 T777 T786 T839 Y929	N95	Ubiquitin carboxyl-terminal hydrolase family 1 UCH-1: R593-D624 Ubiquitin carboxyl-terminal hydrolase family 2 UCH-2: N875-K935 UBIQUITIN CARBOXYL-TERMINAL HYDROLASES FAMILY 2 DM00659 P40818 782-1103: T777-L931, L598-H709, I713-T753, V101-L128 PROTEASE UBIQUITIN HYDROLASE UBIQUITIN SPECIFIC ENZYME DEUBIQUITINATING C-TERMINAL THIOLESTERASE PROCESSING CONJUGATION PD017412: T777-E859 Ubiquitin carboxyl-terminal hydrolase family 2 BL00972: G594-L611, Y675-L684, I714-C728, K878-H902, K904-D925 Ubiquitin carboxyl-terminal hydrolase family 2 signature 1 Uch_2_1: G594-Q609 Ubiquitin carboxyl-terminal hydrolase family 2 signature 2 Uch_2_2: Y879-Y896	HMMER_PFAM HMMER_PFAM BLAST_DOMO BLAST_PRODOM BLIMPS_BLOCKS MOTIFS MOTIFS

Table 4

Polynucleotide SEQ ID NO:	Incyte Polynucleotide ID	Sequence Length	Selected Fragment(s)	Sequence Fragments	5' Position	3' Position
22	275791CB1	2204	1168-1197, 1503-1522, 1-281, 1716-1738	6455514H1 (COLNDIC01)	890	1510
				7246467T8 (PROSTMY01)	692	1390
				4943009F8 (BRAIFEN05)	1319	1877
				55047202J1	1	811
				6053385H1 (BRABDIR03)	1636	2204
23	1389845CB1	2036	1-392, 1468-1491, 1334-1400, 1974-2036	FL1389845_00001	6	2036
				1389845H1 (EOSINOT01)	1	244
24	1726609CB1	2185	1-44, 1804- 2185	71762189V1	1	662
				5426388F9 (PROSTMT07)	1370	1992
				71053940V1	744	1373
				71041539V1	1954	2185
				5968441H1 (BRAZNOT01)	1326	1824
				6758865J1 (SINTER02)	642	1316
				2053131H1 (BEPINOT01)	2885	3136
25	4503848CB1	3486	1-1330	6440674H1 (BRAENOT02)	2576	3095
				95745066	1831	2254
				7191212H2 (BRATDIC01)	2295	2873
				5960039H1 (BRATNOT05)	1229	1797
				GBI.g7710158_edit	1	2015
				60200050D1	825	1146
				5969176H1 (BRAZNOT01)	1677	2104
				5649471H1 (BRAITUT23)	3157	3486
				2232143F6 (PROSNOT16)	2235	2670
				3022114H1 (PROSDIN01)	3116	3402
				60220456D1	1085	1462

Table 4 (cont.)

Polynucleotide SEQ ID NO:	Incyte Polynucleotide ID	Sequence Length	Selected Fragment(s)	Sequence Fragments	5' Position	3' Position
26	5544089CB1	2847	1260-1631, 2532-2847, 408-914	55051688J1	810	1767
				2344450F6 (TESTTUT02)	2195	2847
				7658834H1 (OVARNOE02)	2130	2642
				71763578V1	1562	2266
				6576463H1 (COLHTUS02)	1079	1838
27	7474081CB1	890	1-21	55051680J1.comp	1	958
				92103202	1	493
28	5281209CB1	1577	1-629	92142177	397	890
				FL5281209_g7712102_000	1	1467
				004_g436191		
29	2256251CB1	1958	1-399, 896-935	g3644494	1155	1577
				3142983R6 (HNT2AZS07)	1045	1576
				3220504T6 (COLNNON03)	1611	1958
				g4264312	400	848
				FL2256251_g7708357_000	910	1830
30	7160544CB1	3106	1-540, 1166-1428	002_g6103629		
				2256251R6 (OVARTUT01)	408	1003
				6471337H1 (PLACFEB01)	1654	2316
				6854305H1 (BRAIFEN08)	895	1561
				4443368H1 (SINDNOT01)	1332	1585
				6894004J1 (BRAITDR03)	470	1077
				7655990H1 (UTREDME06)	324	822
				7745974H1 (ADRETUE04)	2378	3052
				7160544H1 (HNT2TXC01)	1	427
				70490289V1	2636	3106
				70748463V1	2269	2891
				7745974J1 (ADRETUE04)	1566	2153

Table 4 (cont.)

Polynucleotide SEQ ID NO:	Incyte Polynucleotide ID	Sequence Length	Selected Fragment(s)	Sequence Fragments	5' Position	3' Position
31	7477386CB1	3567	1-971, 1953-2846, 3243-3567	GBI:g6682143_000029_ed it.20231-20345 GBI:g6682143_000023.co mp_edit.11365-11445 GBI:g6682143_000027_ed it.14110-14265 GBI:g6682143_000029_ed it.35032-35202 GBI:g6682143_000029_ed it.13651-13803 GBI:g6682143_000019_ed it.3461-3655 GBI:g6682143_000029_ed it.41513-41644 GBI:g6682143_000029_ed it.43912-44404 GBI:g6682143_000029_ed it.12846-12930 GBI:g6682143_000029_ed it.15804-15912 GBI:g6682143_000023.co mp_edit.9253-9693 GBI:g6682143_000029_ed it.27621-27746 GBI:g6682143_000029_ed it.18722-18853 GBI:g6682143_000029_ed it.17438-17581 GBI:g6682143_000029_ed it.35651-35783	1495 1 523 2272 958 679 2944 3076 874 1111 82 2068 1363 1219 2608	1608 81 678 2436 1110 873 3075 3567 957 1218 522 2193 1494 1362 2739

Table 4

Polynucleotide SEQ ID NO:	Incyte Polynucleotide ID	Sequence Length	Selected Fragment(s)	Sequence Fragments	5' Position	3' Position
22	275791CB1	2204	1168-1197, 1503-1522, 1-281, 1716-1738	6456514H1 (COLNDIC01)	890	1510
				7246467T8 (PROSTMY01)	692	1390
				4943009F8 (BRAIFEN05)	1319	1877
				55047202J1	1	811
				6053385H1 (BRABDIR03)	1636	2204
23	1389845CB1	2036	1-392, 1468-1491, 1334-1400, 1974-2036	FL1389845_00001	6	2036
				1389845H1 (EOSINOT01)	1	244
24	1726609CB1	2185	1-44, 1804- 2185	71762189V1	1	662
				5426388F9 (PROSTMT07)	1370	1992
				71053940V1	744	1373
				71041539V1	1954	2185
				5968441H1 (BRAZNOT01)	1326	1824
				6756865J1 (SINTFER02)	642	1316
				2053131H1 (BEPINOT01)	2885	3136
				6440674H1 (BRAENOT02)	2576	3095
25	4503848CB1	3486	1-1330	95745066	1831	2254
				7191212H2 (BRATDIC01)	2295	2873
				5960039H1 (BRATNOT05)	1229	1797
				GBI.g7710158_edit	1	2015
				60200050D1	825	1146
				5969176H1 (BRAZNOT01)	1677	2104
				5649471H1 (BRAITUT23)	3157	3486
				2232143F6 (PROSNOT16)	2235	2670
				3022114H1 (PROSDIN01)	3116	3402
				60220456D1	1085	1462

Table 4 (cont.)

Polynucleotide SEQ ID NO:	Incyte Polynucleotide ID	Sequence Length	Selected Fragment(s)	Sequence Fragments	5' Position	3' Position
26	5544089CB1	2847	1260-1631, 2532-2847, 408-914	55051688J1	810	1767
				2344450F6 (TESTTUT02)	2195	2847
				7658834H1 (OVARNOE02)	2130	2642
				71763578V1	1562	2266
				6576463H1 (COLHTUS02)	1079	1838
27	7474081CB1	890	1-21	55051680J1.comp	1	958
				g2103202	1	493
				g2142177	397	890
28	5281209CB1	1577	1-629	FL5281209_g7712102_000	1	1467
				004_g436191		
				g3644494	1155	1577
				3142983R6 (HNT2AZS07)	1045	1576
29	2256251CB1	1958	1-399, 896-935	3220504T6 (COLNNON03)	1611	1958
				g4264312	400	848
				FL2256251_g7708357_000	910	1830
				002_g6103629		
				2256251R6 (OVRTUT01)	408	1003
30	7160544CB1	3106	1-540, 1166-1428	6471337H1 (PLACFEB01)	1654	2316
				6854305H1 (BRAIFEN08)	895	1561
				4443368H1 (SINDNOT01)	1332	1585
				6894004J1 (BRAITDR03)	470	1077
				7655990H1 (UTREDME06)	324	822
				7745974H1 (ADRETUE04)	2378	3052
				7160544H1 (HNT2TXC01)	1	427
				70490289V1	2636	3106
				70748463V1	2269	2891
				7745974J1 (ADRETUE04)	1566	2153

Table 4 (cont.)

Polynucleotide SEQ ID NO:	Incyte Polynucleotide ID	Sequence Length	Selected Fragment(s)	Sequence Fragments	5' Position	3' Position
31	7477386CB1	3567	1-971, 1953-2846, 3243-3567	GBI:g6682143_000029_ed it.20231-20345	1495	1608
				GBI:g6682143_000023.co mp_edit.11365-11445	1	81
				GBI:g6682143_000027_ed it.14110-14265	523	678
				GBI:g6682143_000029_ed it.35032-35202	2272	2436
				GBI:g6682143_000029_ed it.13651-13803	958	1110
				GBI:g6682143_000019_ed it.3461-3655	679	873
				GBI:g6682143_000029_ed it.41513-41644	2944	3075
				GBI:g6682143_000029_ed it.43912-44404	3076	3567
				GBI:g6682143_000029_ed it.12846-12930	874	957
				GBI:g6682143_000029_ed it.15804-15912	1111	1218
				GBI:g6682143_000023.co mp_edit.9253-9693	82	522
				GBI:g6682143_000029_ed it.27621-27746	2068	2193
				GBI:g6682143_000029_ed it.18722-18853	1363	1494
				GBI:g6682143_000029_ed it.17438-17581	1219	1362
				GBI:g6682143_000029_ed it.35651-35783	2608	2739

Table 4 (cont.)

Polynucleotide SEQ ID NO:	Incyte Polynucleotide ID	Sequence Length	Selected Fragment(s)	Sequence Fragments	5' Position	3' Position
32	7473089CB1	2930	1-632, 1082-1138, 2453-2555, 1373-1615, 1716-1740	GBI:g6682143_000029_ed it.27099-27237	1759	1932
				GBI:g6682143_000029_ed it.24540-24713	1609	1758
				GBI:g6682143_000029_ed it.37355-37558	2740	2943
				GBI:g7387384_000011.co mp_edit.13491-13892	2529	2930
				GBI:g7387384_000010_ed it.2924-3211	1335	1619
				GBI:g7387384_000010_ed it.13479-13850	2157	2528
				GBI:g7387384_000010_ed it.11350-11535	1794	1979
				798641H1 (UTRSTUC01)	1064	1587
				GBI:g7387384_000010_ed it.9694-9867	1620	1793
				GBI:g7387384_000012.co mp_edit.9639-10595	75	1031
				GBI:g7387384_000010_ed it.1917-2074	1032	1163
				GBI:g7387384_000010_ed it.2514-2684	1164	1334
				7631548J1 (BRAFTUE03)	21	619
				GBI:g7387384_000010_ed it.11639-11815	1980	2156
				GBI:g7387384_edit	1	2930

Table 4 (cont.)

Polynucleotide SEQ ID NO:	Incyte Polynucleotide ID	Sequence Length	Selected Fragment(s)	Sequence Fragments	5' Position	3' Position
33	7604035CB1	4230	4185-4230, 894-2774	6254235H1 (LUNPTUT02)	3308	3897
				6213818H1 (MUSCDIT06)	3900	4230
				6314348H1 (NERDTDN03)	3426	3994
				7195502H1 (LUNGFER04)	2758	3376
				6800634J1 (COLENOT03)	2678	3323
				8113675H1 (OSTEUNC01)	1661	2049
				6804059H1 (COLENOT03)	478	1050
				7750274H1 (HEAONOE01)	1995	2503
				3843227F6 (DENDNOT01)	2094	2707
				7632961H1 (BLADTUE01)	1418	2024
				5509797J1	688	1535
				7716357J1 (SINTFEE02)	1	678
				71906145V1	1340	2118
				7101935F8 (BRAWTDRO2)	166	760
				70857826V1	2757	3441
				70855756V1	2650	3239
				820867R1 (KERANOT02)	2166	2732
34	3473847CB1	3699	1-2631	8055446J1 (ESOGTUE01)	579	1013
				70857738V1	3156	3699
				70858612V1	1998	2671
				GNN.g7208751_000002_00	555	1850
				2.edit		
				7101935R8 (BRAWTDRO2)	1	473

Table 4 (cont.)

Polynucleotide SEQ ID NO:	Incyte Polynucleotide ID	Sequence Length	Selected Fragment(s)	Sequence Fragments	5' Position	3' Position
35	3750004CB1	2410	1-264, 2116-2410, 1057-1167, 1590-1649	g7712021_edit	1	246
				7680089J1 (BRAFTUE01)	1327	1911
				6804411H1 (COLENO03)	1088	1618
				71909368V1	536	968
				g1187194	1655	2127
				g2241985	706	1144
				6314962H1 (NERDTN03)	973	1135
				6823371J1 (SINTNOR01)	65	855
				7655009J1 (UTREDME06)	1407	1990
				g1272147	1754	2410
36	4904126CB1	549		71620969V1	1	549
37	71268415CB1	2755	1-1097, 2326-2755	771527J1 (SINTFEE02)	590	1340
				7372052H2 (BRAIFEE04)	2044	2514
				g6651070_CD	102	2755
				7723192J2 (THYRDIE01)	905	1500
				GBI:g7709257_000011.ed it	1	139
				8037549H1 (SMCRUNE01)	206	819
				7720289J1 (THYRDIE01)	1596	2263
				8037549J1 (SMCRUNE01)	1456	2120

Table 4 (cont.)

Polynucleotide SEQ ID NO:	Incyte Polynucleotide ID	Sequence Length	Selected Fragment(s)	Sequence Fragments	5' Position	3' Position
38	7473301CB1	2553	1-2394	GBI.g7272157_000017.ed it 71704195V1 5544473H1 (TESTNOC01) GNN.g7272157_000017_00 2.edit 5544473T8 (TESTNOC01) 71703469V1 GNN.g8571511_000004_00 2.edit GNN.g6624046_000008_00 4	2001	2553
39	7473308CB1	1041	826-1041, 1-299	GNN.g1552511_035	1	1041
40	7478021CB1	1707	1-1188	g8176728_edit g7684439_edit g7684439_edit_2 71571956V1 5634861R8 (PLACFER01) 71571988V1 71573159V1	979 1 1084 704 1 256 247	1083 978 1707 1262 266 937 928
41	4333459CB1	1262	1-1262	55022864H1 55022792H2 55022814H1 55022795J2	2314 1392 2080 2044	3067 2091 2886 2726
42	6817347CB1	3067	1-2270	GNN.g7417337_004.edit 1	1	3067

Table 5

Polynucleotide SEQ ID NO:	Incyte Project ID	Representative Library
22	275791CB1	TESTNOT03
23	1389845CB1	EOSITXT01
24	1726609CB1	BRAITUT02
25	4503848CB1	PROSNOT16
26	5544089CB1	BRAIFEC01
28	5281209CB1	HNT2AZS07
29	2256251CB1	OVARTUT01
30	7160544CB1	BRAFNOT02
32	7473089CB1	UTRSTUC01
33	7604035CB1	PLACNOR01
34	3473847CB1	KERANOT02
35	3750004CB1	BRAFTUE01
36	4904126CB1	TYMNOT08
37	71268415CB1	THYRDIE01
38	7473301CB1	TESTNOC01
41	4333459CB1	KIDCTMT01
42	6817347CB1	ADRETUR01

Table 6

Library	Vector	Library Description
ADRETUR01	PCDNA2.1	This random primed library was constructed using RNA isolated from left upper pole, adrenal gland tumor tissue removed from a 52-year-old Caucasian male during nephroureterectomy and local destruction of renal lesion. Pathology indicated grade 3 adrenal cortical carcinoma forming a mass that infiltrated almost the whole adrenal parenchyma and extended to adjacent adipose tissue. A metastatic tumor nodule was identified in the hilar region. The renal vein was infiltrated by tumor and the neoplastic process was present at the resection margin of the renal vein. Fragments of adrenal cortical carcinoma and thrombus were found in the inferior vena cava. Patient history included abnormal weight loss. Family history included skin cancer, type I diabetes, and neurotic depression.
BRAFNOT02	pINCY	Library was constructed using RNA isolated from superior frontal cortex tissue removed from a 35-year-old Caucasian male who died from cardiac failure. Pathology indicated moderate leptomeningeal fibrosis and multiple microinfarctions of the cerebral neocortex. Microscopically, the cerebral hemisphere revealed moderate fibrosis of the leptomeninges with focal calcifications. There was evidence of shrunken and slightly eosinophilic pyramidal neurons throughout the cerebral hemispheres. In addition, scattered throughout the cerebral cortex, there were multiple small microscopic areas of cavitation with surrounding gliosis. Patient history included dilated cardiomyopathy, congestive heart failure, cardiomegaly, and an enlarged spleen and liver.
BRAFTUE01	PCDNA2.1	This 5' biased random primed library was constructed using RNA isolated from brain tumor tissue removed from the frontal lobe of a 58-year-old Caucasian male during excision of a cerebral meningeal lesion. Pathology indicated a grade 2 metastatic hypernephroma. The patient presented with migraine headache. The patient developed a cerebral hemorrhage and pulmonary edema, and died during this hospitalization. Patient history included a grade 2 renal cell carcinoma, insomnia, and chronic airway obstruction. Previous surgeries included a nephroureterectomy. Patient medications included Decadron and Dilantin. Family history included a malignant neoplasm of the kidney in the father.
BRAIFEC01	pINCY	This large size-fractionated library was constructed using RNA isolated from brain tissue removed from a Caucasian male fetus who was stillborn with a hypoplastic left heart at 23 weeks' gestation.

Table 6 (cont.)

Library	Vector	Library Description
BRAITUT02	PSPORT1	Library was constructed using RNA isolated from brain tumor tissue removed from the frontal lobe of a 58-year-old Caucasian male during excision of a cerebral meningeal lesion. Pathology indicated a grade 2 metastatic hypernephroma. Patient history included a grade 2 renal cell carcinoma, insomnia, and chronic airway obstruction. Family history included a malignant neoplasm of the kidney.
EOSITXT01	pINCY	Library was constructed using RNA isolated from eosinophils stimulated with IL-5.
HNT2AZS07	PSPORT1	This subtracted library was constructed from RNA isolated from an hNT2 cell line (derived from a human teratocarcinoma that exhibited properties characteristic of a committed neuronal precursor) treated for three days with 0.35 micromolar A2. The hybridization probe for subtraction was derived from a similarly constructed library from untreated hNT2 cells. 3.08M clones from the A2-treated library were subjected to three rounds of subtractive hybridization with 3.04M clones from the untreated library. Subtractive hybridization conditions were based on the methodologies of Swaroop et al. (NAR (1991) 19:1954) and Bonaldo et al. (Genome Research (1996) 6:791).
KERANOT02	PSPORT1	Library was constructed using RNA isolated from epidermal breast keratinocytes (NHEK). NHEK (Clontech #CC-2501) is human breast keratinocyte cell line derived from a 30-year-old black female during breast-reduction surgery.
KIDCTMT01	pINCY	Library was constructed using RNA isolated from kidney cortex tissue removed from a 65-year-old male during nephroureterectomy. Pathology for the associated tumor tissue indicated grade 3 renal cell carcinoma within the mid-portion of the kidney and the renal capsule.
OVARTUT01	PSPORT1	Library was constructed using RNA isolated from ovarian tumor tissue removed from a 43-year-old Caucasian female during removal of the fallopian tubes and ovaries. Pathology indicated grade 2 mucinous cystadenocarcinoma involving the entire left ovary. Patient history included mitral valve disorder, pneumonia, and viral hepatitis. Family history included atherosclerotic coronary artery disease, pancreatic cancer, stress reaction, cerebrovascular disease, breast cancer, and uterine cancer.

Table 6 (cont.)

Library	Vector	Library Description
PLACNOR01	PCDNA2.1	This random primed library was constructed using pooled cDNA from two different donors. cDNA was generated using mRNA isolated from placental tissue removed from a Caucasian fetus (donor A), who died after 16 weeks' gestation from fetal demise and hydrocephalus and from placental tissue removed from a Caucasian male fetus (donor B), who died after 18 weeks' gestation from fetal demise. Patient history for donor A included umbilical cord wrapped around the head (3 times) and the shoulders (1 time). Serology was positive for anti-CMV and remaining serologies were negative. Family history included multiple pregnancies and live births, and an abortion in the mother. Serology was negative for donor B.
PROSNOT16	pINCY	Library was constructed using RNA isolated from diseased prostate tissue removed from a 68-year-old Caucasian male during a radical prostatectomy. Pathology indicated adenofibromatous hyperplasia. Pathology for the associated tumor tissue indicated an adenocarcinoma (Gleason grade 3+4). The patient presented with elevated prostate specific antigen (PSA). During this hospitalization, the patient was diagnosed with myasthenia gravis. Patient history included osteoarthritis, and type II diabetes. Family history included benign hypertension, acute myocardial infarction, hyperlipidemia, and arteriosclerotic coronary artery disease.
TESTNOC01	PBLUESCRIPT	This large size fractionated library was constructed using RNA isolated from testicular tissue removed from a pool of eleven, 10 to 61-year-old Caucasian males.
TESTNOT03	PBLUESCRIPT	Library was constructed using RNA isolated from testicular tissue removed from a 37-year-old Caucasian male, who died from liver disease. Patient history included cirrhosis, jaundice, and liver failure.
THYRDIE01	PCDNA2.1	This 5' biased random primed library was constructed using RNA isolated from diseased thyroid tissue removed from a 22-year-old Caucasian female during closed thyroid biopsy, partial thyroidectomy, and regional lymph node excision. Pathology indicated adenomatous hyperplasia. The patient presented with malignant neoplasm of the thyroid. Patient history included normal delivery, alcohol abuse, and tobacco abuse. Previous surgeries included myringotomy. Patient medications included an unspecified type of birth control pills. Family history included hyperlipidemia and depressive disorder in the mother; and benign hypertension, congestive heart failure, and chronic leukemia in the grandparent(s).

Table 6 (cont.)

Library	Vector	Library Description
TLYMNOT08	pINCY	The library was constructed using RNA isolated from anergic/allogenic T-lymphocyte tissue removed from an adult (40-50-year-old) Caucasian male. The cells were incubated for 3 days in the presence of 1 microgram/ml OKT3 mAb and 5% human serum.
UTRSTUC01	PSPORT1	This large size fractionated library was constructed using pooled cDNA from two donors. cDNA was generated using mRNA isolated from uterus tumor tissue removed from a 37-year-old Black female (donor A) during myomectomy, dilation and curettage, right fimbrial region biopsy, and incidental appendectomy; and from endometrial tumor tissue removed from a 49-year-old Caucasian female (donor B) during vaginal hysterectomy and bilateral salpingo-oophorectomy. For donor A, pathology indicated multiple uterine leiomyomata. A fimbrial cyst was identified. The endometrium was in secretory phase with hormonal effect. The patient presented with deficiency anemia, an umbilical hernia, and premenopausal menorrhagia. Patient history included premenopausal menorrhagia and sarcooidosis of the lung. Previous surgeries included hysterectomy, dilation and curettage, and endoscopic lung biopsy. Patient medications included Chromagen and Claritin. For donor B, pathology indicated grade 3 adenocarcinoma forming a mass within the uterine fundus and involving the anterior uterine wall, as well as focally involving an adjacent endometrial polyp. The tumor invaded to a maximum depth of 7mm (uninvolved wall thickness, 2.2cm). The adjacent endometrium was inactive. Paraffin section immunostains for estrogen receptors and progesterone receptors were positive. Patient history included malignant breast neoplasm. Previous surgeries included unilateral extended simple mastectomy and bilateral tubal destruction. Patient medications included Megace and CAF (Cyclophosphamide, Adriamycin, Fluorocil).

Table 7

Program	Description	Reference	Parameter Threshold
ABI FACTURA	A program that removes vector sequences and masks ambiguous bases in nucleic acid sequences.	Applied Biosystems, Foster City, CA.	
ABI/PARACEL FDF	A Fast Data Finder useful in comparing and annotating amino acid or nucleic acid sequences.	Applied Biosystems, Foster City, CA; Paracel Inc., Pasadena, CA.	Mismatch <50%
ABI AutoAssembler	A program that assembles nucleic acid sequences.	Applied Biosystems, Foster City, CA.	
BLAST	A Basic Local Alignment Search Tool useful in sequence similarity search for amino acid and nucleic acid sequences. BLAST includes five functions: blastp, blastn, blastx, tblastn, and tblastx.	Altschul, S.F. et al. (1990) <i>J. Mol. Biol.</i> 215:403-410; Altschul, S.F. et al. (1997) <i>Nucleic Acids Res.</i> 25:3389-3402.	ESTs: Probability value= 1.0E-8 or less <i>Full Length sequences</i> : Probability value= 1.0E-10 or less
FASTA	A Pearson and Lipman algorithm that searches for similarity between a query sequence and a group of sequences of the same type. FASTA comprises at least five functions: fasta, tfasta, fastx, tfastx, and ssearch.	Pearson, W.R. and D.J. Lipman (1988) <i>Proc. Natl. Acad. Sci. USA</i> 85:2444-2448; Pearson, W.R. (1990) <i>Methods Enzymol.</i> 183:63-98; and Smith, T.F. and M.S. Waterman (1981) <i>Adv. Appl. Math.</i> 2:482-489.	ESTs: fasta E value=1.0E-6 <i>Assembled ESTs</i> : fasta Identity= 95% or greater and Match length=200 bases or greater; fastx E value=1.0E-8 or less <i>Full Length sequences</i> : fastx score=100 or greater
BLIMPS	A BLocks IMProved Searcher that matches a sequence against those in BLOCKS, PRINTS, DOMO, PRODOM, and PFAM databases to search for gene families, sequence homology, and structural fingerprint regions.	Henikoff, S. and J.G. Henikoff (1991) <i>Nucleic Acids Res.</i> 19:6565-6572; Henikoff, J.G. and S. Henikoff (1996) <i>Methods Enzymol.</i> 266:88-105; and Attwood, T.K. et al. (1997) <i>J. Chem. Inf. Comput. Sci.</i> 37:417-424.	Probability value= 1.0E-3 or less
HMMER	An algorithm for searching a query sequence against hidden Markov model (HMM)-based databases of protein family consensus sequences, such as PFAM.	Krogh, A. et al. (1994) <i>J. Mol. Biol.</i> 235:1501-1531; Sonnhammer, B.L.L. et al. (1998) <i>Nucleic Acids Res.</i> 26:320-322; Durbin, R. et al. (1998) <i>Our World View</i> , in a Nutshell, Cambridge Univ. Press, pp. 1-350.	PFAM hits: Probability value= 1.0E-3 or less <i>Signal peptide hits</i> : Score= 0 or greater

Table 7 (cont.)

Program	Description	Reference	Parameter Threshold
ProfileScan	An algorithm that searches for structural and sequence motifs in protein sequences that match sequence patterns defined in Prosite.	Gribskov, M. et al. (1988) CABIOS 4:61-66; Gribskov, M. et al. (1989) Methods Enzymol. 183:146-159; Bairoch, A. et al. (1997) Nucleic Acids Res. 25:217-221.	Normalized quality score2 GCG-specified "HIGH" value for that particular Prosite motif. Generally, score=1.4-2.1.
Phred	A base-calling algorithm that examines automated sequencer traces with high sensitivity and probability.	Ewing, B. et al. (1998) Genome Res. 8:175-185; Ewing, B. and P. Green (1998) Genome Res. 8:186-194.	
Phrap	A Phils Revised Assembly Program including SWAT and CrossMatch, programs based on efficient implementation of the Smith-Waterman algorithm, useful in searching sequence homology and assembling DNA sequences.	Smith, T.F. and M.S. Waterman (1981) Adv. Appl. Math. 2:482-489; Smith, T.F. and M.S. Waterman (1981) J. Mol. Biol. 147:195-197; and Green, P., University of Washington, Seattle, WA.	Score= 120 or greater; Match length= 56 or greater
Consed	A graphical tool for viewing and editing Phrap assemblies.	Gordon, D. et al. (1998) Genome Res. 8:195-202.	
SPScan	A weight matrix analysis program that scans protein sequences for the presence of secretory signal peptides.	Nielson, H. et al. (1997) Protein Engineering 10:1-6; Claverie, J.M. and S. Audic (1997) CABIOS 12:431-439.	Score=3.5 or greater
TMAP	A program that uses weight matrices to delineate transmembrane segments on protein sequences and determine orientation.	Persson, B. and P. Argos (1994) J. Mol. Biol. 237:182-192; Persson, B. and P. Argos (1996) Protein Sci. 5:363-371.	
TMHMMER	A program that uses a hidden Markov model (HMM) to delineate transmembrane segments on protein sequences and determine orientation.	Sonnhammer, E.L. et al. (1998) Proc. Sixth Intl. Conf. on Intelligent Systems for Mol. Biol., Glasgow et al., eds., The Am. Assoc. for Artificial Intelligence Press, Menlo Park, CA, pp. 175-182.	
Motifs	A program that searches amino acid sequences for patterns that matched those defined in Prosite.	Bairoch, A. et al. (1997) Nucleic Acids Res. 25:217-221; Wisconsin Package Program Manual, version 9, page M51-59, Genetics Computer Group, Madison, WI.	

What is claimed is:

1. An isolated polypeptide selected from the group consisting of:
 - a) a polypeptide comprising an amino acid sequence selected from the group consisting of
5 SEQ ID NO:1-21,
 - b) a polypeptide comprising a naturally occurring amino acid sequence at least 90% identical to an amino acid sequence selected from the group consisting of SEQ ID NO:1-21, ..
 - c) a biologically active fragment of a polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO:1-21, and
 - 10 d) an immunogenic fragment of a polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO:1-21.
2. An isolated polypeptide of claim 1 selected from the group consisting of SEQ ID NO:1-
21.
15
3. An isolated polynucleotide encoding a polypeptide of claim 1.
4. An isolated polynucleotide encoding a polypeptide of claim 2.
- 20 5. An isolated polynucleotide of claim 4 selected from the group consisting of SEQ ID NO:22-42.
6. A recombinant polynucleotide comprising a promoter sequence operably linked to a polynucleotide of claim 3.
25
7. A cell transformed with a recombinant polynucleotide of claim 6.
8. A transgenic organism comprising a recombinant polynucleotide of claim 6.
- 30 9. A method for producing a polypeptide of claim 1, the method comprising:
 - a) culturing a cell under conditions suitable for expression of the polypeptide, wherein said cell is transformed with a recombinant polynucleotide, and said recombinant polynucleotide comprises a promoter sequence operably linked to a polynucleotide encoding the polypeptide of claim 1, and
 - 35 b) recovering the polypeptide so expressed.

10. An isolated antibody which specifically binds to a polypeptide of claim 1.
11. An isolated polynucleotide selected from the group consisting of:
- a) a polynucleotide comprising a polynucleotide sequence selected from the group consisting
 - 5 of SEQ ID NO:22-42,
 - b) a polynucleotide comprising a naturally occurring polynucleotide sequence at least 90% identical to a polynucleotide sequence selected from the group consisting of SEQ ID NO:22-42,
 - c) a polynucleotide complementary to a polynucleotide of a),
 - d) a polynucleotide complementary to a polynucleotide of b), and
 - 10 e) an RNA equivalent of a)-d).
12. An isolated polynucleotide comprising at least 60 contiguous nucleotides of a polynucleotide of claim 11.
13. A method for detecting a target polynucleotide in a sample, said target polynucleotide
- 15 having a sequence of a polynucleotide of claim 11, the method comprising:
- a) hybridizing the sample with a probe comprising at least 20 contiguous nucleotides comprising a sequence complementary to said target polynucleotide in the sample, and which probe
 - specifically hybridizes to said target polynucleotide, under conditions whereby a hybridization
 - 20 complex is formed between said probe and said target polynucleotide or fragments thereof, and
 - b) detecting the presence or absence of said hybridization complex, and, optionally, if present, the amount thereof.
14. A method of claim 13, wherein the probe comprises at least 60 contiguous nucleotides.
- 25 15. A method for detecting a target polynucleotide in a sample, said target polynucleotide having a sequence of a polynucleotide of claim 11, the method comprising:
- a) amplifying said target polynucleotide or fragment thereof using polymerase chain reaction amplification, and
 - 30 b) detecting the presence or absence of said amplified target polynucleotide or fragment thereof, and, optionally, if present, the amount thereof.
16. A composition comprising a polypeptide of claim 1 and a pharmaceutically acceptable excipient.

35

17. A composition of claim 16, wherein the polypeptide has an amino acid sequence selected from the group consisting of SEQ ID NO:1-21.

18. A method for treating a disease or condition associated with decreased expression of functional PRTS, comprising administering to a patient in need of such treatment the composition of claim 16.

19. A method for screening a compound for effectiveness as an agonist of a polypeptide of claim 1, the method comprising:

- a) exposing a sample comprising a polypeptide of claim 1 to a compound, and
- b) detecting agonist activity in the sample.

20. A composition comprising an agonist compound identified by a method of claim 19 and a pharmaceutically acceptable excipient.

21. A method for treating a disease or condition associated with decreased expression of functional PRTS, comprising administering to a patient in need of such treatment a composition of claim 20.

22. A method for screening a compound for effectiveness as an antagonist of a polypeptide of claim 1, the method comprising:

- a) exposing a sample comprising a polypeptide of claim 1 to a compound, and
- b) detecting antagonist activity in the sample.

23. A composition comprising an antagonist compound identified by a method of claim 22 and a pharmaceutically acceptable excipient.

24. A method for treating a disease or condition associated with overexpression of functional PRTS, comprising administering to a patient in need of such treatment a composition of claim 23.

25. A method of screening for a compound that specifically binds to the polypeptide of claim 1, said method comprising the steps of:

- a) combining the polypeptide of claim 1 with at least one test compound under suitable conditions, and

b) detecting binding of the polypeptide of claim 1 to the test compound, thereby identifying a compound that specifically binds to the polypeptide of claim 1.

26. A method of screening for a compound that modulates the activity of the polypeptide of claim 1, said method comprising:

- a) combining the polypeptide of claim 1 with at least one test compound under conditions permissive for the activity of the polypeptide of claim 1,
- b) assessing the activity of the polypeptide of claim 1 in the presence of the test compound, and
- c) comparing the activity of the polypeptide of claim 1 in the presence of the test compound with the activity of the polypeptide of claim 1 in the absence of the test compound, wherein a change in the activity of the polypeptide of claim 1 in the presence of the test compound is indicative of a compound that modulates the activity of the polypeptide of claim 1.

27. A method for screening a compound for effectiveness in altering expression of a target polynucleotide, wherein said target polynucleotide comprises a sequence of claim 5, the method comprising:

- a) exposing a sample comprising the target polynucleotide to a compound, under conditions suitable for the expression of the target polynucleotide,
- b) detecting altered expression of the target polynucleotide, and
- c) comparing the expression of the target polynucleotide in the presence of varying amounts of the compound and in the absence of the compound.

28. A method for assessing toxicity of a test compound, said method comprising:

- a) treating a biological sample containing nucleic acids with the test compound;
- b) hybridizing the nucleic acids of the treated biological sample with a probe comprising at least 20 contiguous nucleotides of a polynucleotide of claim 11 under conditions whereby a specific hybridization complex is formed between said probe and a target polynucleotide in the biological sample, said target polynucleotide comprising a polynucleotide sequence of a polynucleotide of claim 11 or fragment thereof;
- c) quantifying the amount of hybridization complex; and
- d) comparing the amount of hybridization complex in the treated biological sample with the amount of hybridization complex in an untreated biological sample, wherein a difference in the amount of hybridization complex in the treated biological sample is indicative of toxicity of the test compound.

29. A diagnostic test for a condition or disease associated with the expression of PRTS in a biological sample comprising the steps of:

- a) combining the biological sample with an antibody of claim 10, under conditions suitable for the antibody to bind the polypeptide and form an antibody:polypeptide complex; and
- b) detecting the complex, wherein the presence of the complex correlates with the presence of the polypeptide in the biological sample.

30. The antibody of claim 10, wherein the antibody is:

- a) a chimeric antibody,
- b) a single chain antibody,
- c) a Fab fragment,
- d) a F(ab')₂ fragment, or
- e) a humanized antibody.

31. A composition comprising an antibody of claim 10 and an acceptable excipient.

32. A method of diagnosing a condition or disease associated with the expression of PRTS in a subject, comprising administering to said subject an effective amount of the composition of claim

31.

33. A composition of claim 31, wherein the antibody is labeled.

34. A method of diagnosing a condition or disease associated with the expression of PRTS in a subject, comprising administering to said subject an effective amount of the composition of claim

33.

35. A method of preparing a polyclonal antibody with the specificity of the antibody of claim 10 comprising:

- a) immunizing an animal with a polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO:1-21, or an immunogenic fragment thereof, under conditions to elicit an antibody response;
- b) isolating antibodies from said animal; and

c) screening the isolated antibodies with the polypeptide, thereby identifying a polyclonal antibody which binds specifically to a polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO:1-21.

5 36. An antibody produced by a method of claim 35.

37. A composition comprising the antibody of claim 36 and a suitable carrier.

10 38. A method of making a monoclonal antibody with the specificity of the antibody of claim 10 comprising:

a) immunizing an animal with a polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO:1-21, or an immunogenic fragment thereof, under conditions to elicit an antibody response;

b) isolating antibody producing cells from the animal;

15 c) fusing the antibody producing cells with immortalized cells to form monoclonal antibody-producing hybridoma cells;

d) culturing the hybridoma cells; and

e) isolating from the culture monoclonal antibody which binds specifically to a polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO:1-21.

20

39. A monoclonal antibody produced by a method of claim 38.

40. A composition comprising the antibody of claim 39 and a suitable carrier.

25 41. The antibody of claim 10, wherein the antibody is produced by screening a Fab expression library.

42. The antibody of claim 10, wherein the antibody is produced by screening a recombinant immunoglobulin library.

30

43. A method for detecting a polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO:1-21 in a sample, comprising the steps of:

a) incubating the antibody of claim 10 with a sample under conditions to allow specific binding of the antibody and the polypeptide; and

b) detecting specific binding, wherein specific binding indicates the presence of a polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO:1-21 in the sample.

- 5 44. A method of purifying a polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO:1-21 from a sample, the method comprising:
- a) incubating the antibody of claim 10 with a sample under conditions to allow specific binding of the antibody and the polypeptide; and
- b) separating the antibody from the sample and obtaining the purified polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO:1-21.
- 10
45. A polypeptide of claim 1, comprising the amino acid sequence of SEQ ID NO:1.
- 15 46. A polypeptide of claim 1, comprising the amino acid sequence of SEQ ID NO:2.
47. A polypeptide of claim 1, comprising the amino acid sequence of SEQ ID NO:3.
48. A polypeptide of claim 1, comprising the amino acid sequence of SEQ ID NO:4.
- 20 49. A polypeptide of claim 1, comprising the amino acid sequence of SEQ ID NO:5.
50. A polypeptide of claim 1, comprising the amino acid sequence of SEQ ID NO:6.
- 25 51. A polypeptide of claim 1, comprising the amino acid sequence of SEQ ID NO:7.
52. A polypeptide of claim 1, comprising the amino acid sequence of SEQ ID NO:8.
53. A polypeptide of claim 1, comprising the amino acid sequence of SEQ ID NO:9.
- 30 54. A polypeptide of claim 1, comprising the amino acid sequence of SEQ ID NO:10.
55. A polypeptide of claim 1, comprising the amino acid sequence of SEQ ID NO:11.

56. A polypeptide of claim 1, comprising the amino acid sequence of SEQ ID NO:12.
57. A polypeptide of claim 1, comprising the amino acid sequence of SEQ ID NO:13.
- 5 58. A polypeptide of claim 1, comprising the amino acid sequence of SEQ ID NO:14.
59. A polypeptide of claim 1, comprising the amino acid sequence of SEQ ID NO:15.
60. A polypeptide of claim 1, comprising the amino acid sequence of SEQ ID NO:16.
- 10 61. A polypeptide of claim 1, comprising the amino acid sequence of SEQ ID NO:17.
62. A polypeptide of claim 1, comprising the amino acid sequence of SEQ ID NO:18.
63. A polypeptide of claim 1, comprising the amino acid sequence of SEQ ID NO:19.
- 15 64. A polypeptide of claim 1, comprising the amino acid sequence of SEQ ID NO:20.
65. A polypeptide of claim 1, comprising the amino acid sequence of SEQ ID NO:21.
- 20 66. A polynucleotide of claim 11, comprising the polynucleotide sequence of SEQ ID NO:22.
67. A polynucleotide of claim 11, comprising the polynucleotide sequence of SEQ ID NO:23.
- 25 68. A polynucleotide of claim 11, comprising the polynucleotide sequence of SEQ ID NO:24.
69. A polynucleotide of claim 11, comprising the polynucleotide sequence of SEQ ID NO:25.
- 30 70. A polynucleotide of claim 11, comprising the polynucleotide sequence of SEQ ID NO:26.

71. A polynucleotide of claim 11, comprising the polynucleotide sequence of SEQ ID NO:27.

5 72. A polynucleotide of claim 11, comprising the polynucleotide sequence of SEQ ID NO:28.

73. A polynucleotide of claim 11, comprising the polynucleotide sequence of SEQ ID NO:29.

10 74. A polynucleotide of claim 11, comprising the polynucleotide sequence of SEQ ID NO:30.

15 75. A polynucleotide of claim 11, comprising the polynucleotide sequence of SEQ ID NO:31.

76. A polynucleotide of claim 11, comprising the polynucleotide sequence of SEQ ID NO:32.

20 77. A polynucleotide of claim 11, comprising the polynucleotide sequence of SEQ ID NO:33.

78. A polynucleotide of claim 11, comprising the polynucleotide sequence of SEQ ID NO:34.

25 79. A polynucleotide of claim 11, comprising the polynucleotide sequence of SEQ ID NO:35.

30 80. A polynucleotide of claim 11, comprising the polynucleotide sequence of SEQ ID NO:36.

81. A polynucleotide of claim 11, comprising the polynucleotide sequence of SEQ ID NO:37.

82. A polynucleotide of claim 11, comprising the polynucleotide sequence of SEQ ID
NO:38.

83. A polynucleotide of claim 11, comprising the polynucleotide sequence of SEQ ID
5 NO:39.

84. A polynucleotide of claim 11, comprising the polynucleotide sequence of SEQ ID
NO:40.

10 85. A polynucleotide of claim 11, comprising the polynucleotide sequence of SEQ ID
NO:41.

86. A polynucleotide of claim 11, comprising the polynucleotide sequence of SEQ ID
NO:42.

<110> INCYTE GENOMICS, INC.
 YUE, Henry
 ELLIOTT, Vicki
 GANDHI, Aameena R.
 LAL, Preeti
 AU-YOUNG, Janice
 TRIBOULEY, Catherine M.
 DELEGEANE, Angelo M.
 BAUGHN, Mariah R.
 NGUYEN, Danniel B.
 LEE, Ernestine A.
 HAFALIA, April
 KHAN, Farrah A.
 WALIA, Narinder K.
 YAO, Monique G.
 LU, Dyung Aina M.
 PATTERSON, Chandra
 TANG, Y. Tom
 WALSH, Roderick T.
 AZIMZAI, Yalda
 LU, Yan
 RAMKUMAR, Jayalaximi
 XU, Yuming
 REDDY, Roopa
 DAS, Depopriya
 KEARNEY, Liam
 KALLICK, Deborah A.

<120> Proteases

<130> PI-0123 PCT

<140> To Be Assigned

<141> Herewith

<150> 60/212,336; 60/213,955; 60/215,396; 60/216,821; 60/218,946

<151> 2000-06-16; 2000-06-22; 2000-06-29; 2000-07-07; 2000-07-14

<160> 42

<170> PERL Program

<210> 1

<211> 232

<212> PRT

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 275791CD1

<400> 1

Met	Pro	Glu	Asn	Pro	Asp	Thr	Met	Glu	Thr	Glu	Lys	Pro	Lys	Thr
1				5					10					15
Ile	Thr	Glu	Leu	Asp	Pro	Ala	Ser	Phe	Thr	Glu	Ile	Thr	Lys	Asp
				20					25					30
Cys	Asp	Glu	Asn	Lys	Glu	Asn	Lys	Thr	Pro	Glu	Gly	Ser	Gln	Gly
				35					40					45
Glu	Val	Asp	Trp	Leu	Gln	Gln	Tyr	Asp	Met	Glu	Arg	Glu	Arg	Glu
				50					55					60
Glu	Gln	Glu	Leu	Gln	Gln	Ala	Leu	Ala	Gln	Ser	Leu	Gln	Glu	Gln
				65					70					75
Glu	Ala	Trp	Glu	Gln	Lys	Glu	Asp	Asp	Asp	Leu	Lys	Arg	Ala	Thr
				80					85					90
Glu	Leu	Ser	Leu	Gln	Glu	Phe	Asn	Asn	Ser	Phe	Val	Asp	Ala	Leu
				95					100					105
Gly	Ser	Asp	Glu	Asp	Ser	Gly	Asn	Glu	Asp	Val	Phe	Asp	Met	Glu
				110					115					120

Tyr	Thr	Glu	Ala	Glu	Ala	Glu	Glu	Leu	Lys	Arg	Asn	Ala	Glu	Thr	
				125					130					135	
Gly	Asn	Leu	Pro	His	Ser	Tyr	Arg	Leu	Ile	Ser	Val	Val	Ser	His	
				140					145					150	
Ile	Gly	Ser	Thr	Ser	Ser	Ser	Gly	His	Tyr	Ile	Ser	Asp	Val	Tyr	
				155					160					165	
Asp	Ile	Lys	Lys	Gln	Ala	Trp	Phe	Thr	Tyr	Asn	Asp	Leu	Glu	Val	
				170					175					180	
Ser	Lys	Ile	Gln	Glu	Ala	Ala	Val	Gln	Ser	Asp	Arg	Asp	Arg	Ser	
				185					190					195	
Gly	Tyr	Ile	Phe	Phe	Tyr	Met	His	Lys	Glu	Ile	Phe	Asp	Glu	Leu	
				200					205					210	
Leu	Glu	Thr	Glu	Lys	Asn	Ser	Gln	Ser	Leu	Ser	Thr	Glu	Val	Gly	
				215					220					225	
Lys	Thr	Thr	Arg	Gln	Ala	Ser									
				230											

<210> 2

<211> 365

<212> PRT

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 1389845CD1

<400> 2

Met	Pro	Lys	Tyr	Leu	Gly	Gly	Gly	Cys	Cys	Ile	Pro	Gly	Pro	Trp	
1				5					10					15	
Ala	Glu	Arg	Arg	Val	Tyr	Ser	Leu	Gly	His	Gln	Asp	Lys	Ser	Arg	
				20					25					30	
Thr	His	Gln	Glu	Leu	Arg	Thr	Asp	Arg	Arg	Thr	Thr	Glu	Gly	Val	
				35					40					45	
Thr	Gly	Trp	Cys	Glu	Asp	Trp	Cys	Pro	Trp	Ala	Arg	Thr	Leu	Leu	
				50					55					60	
Ser	Ser	Pro	Cys	Trp	Leu	Gln	Thr	Arg	Val	Gln	Ala	Leu	Gly	Ser	
				65					70					75	
Ala	Thr	Leu	Thr	Gln	Pro	Ser	Leu	Glu	Asp	Arg	Met	Arg	Gly	Val	
				80					85					90	
Ser	Cys	Leu	Gln	Val	Leu	Leu	Leu	Leu	Val	Leu	Gly	Ala	Ala	Gly	
				95					100					105	
Thr	Gln	Gly	Arg	Lys	Ser	Ala	Ala	Cys	Gly	Gln	Pro	Arg	Met	Ser	
				110					115					120	
Ser	Arg	Ile	Val	Gly	Gly	Arg	Asp	Gly	Arg	Asp	Gly	Glu	Trp	Pro	
				125					130					135	
Trp	Gln	Ala	Ser	Ile	Gln	His	Arg	Gly	Ala	His	Val	Cys	Gly	Gly	
				140					145					150	
Ser	Leu	Ile	Ala	Pro	Gln	Trp	Val	Leu	Thr	Ala	Ala	His	Cys	Phe	
				155					160					165	
Pro	Arg	Arg	Ala	Leu	Pro	Ala	Glu	Tyr	Arg	Val	Arg	Leu	Gly	Ala	
				170					175					180	
Leu	Arg	Leu	Gly	Ser	Thr	Ser	Pro	Arg	Thr	Leu	Ser	Val	Pro	Val	
				185					190					195	
Arg	Arg	Val	Leu	Leu	Pro	Pro	Asp	Tyr	Ser	Glu	Asp	Gly	Ala	Arg	
				200					205					210	
Gly	Asp	Leu	Ala	Leu	Leu	Gln	Leu	Arg	Arg	Pro	Val	Pro	Leu	Ser	
				215					220					225	
Ala	Arg	Val	Gln	Pro	Val	Cys	Leu	Pro	Val	Pro	Gly	Ala	Arg	Pro	
				230					235					240	
Pro	Pro	Gly	Thr	Pro	Cys	Arg	Val	Thr	Gly	Trp	Gly	Ser	Leu	Arg	
				245					250					255	
Pro	Gly	Val	Pro	Leu	Pro	Glu	Trp	Arg	Pro	Leu	Gln	Gly	Val	Arg	
				260					265					270	
Val	Pro	Leu	Leu	Asp	Ser	Arg	Thr	Cys	Asp	Gly	Leu	Tyr	His	Val	
				275					280					285	
Gly	Ala	Asp	Val	Pro	Gln	Ala	Glu	Arg	Ile	Val	Leu	Pro	Gly	Ser	
				290					295					300	

Leu	Cys	Ala	Gly	Tyr	Pro	Gln	Gly	His	Lys	Asp	Ala	Cys	Gln	Val
				305					310					315
Cys	Thr	Gln	Pro	Pro	Gln	Pro	Pro	Glu	Ser	Pro	Pro	Cys	Ala	Gln
				320					325					330
His	Pro	Pro	Ser	Leu	Asn	Ser	Arg	Thr	Gln	Asp	Ile	Pro	Thr	Gln
				335					340					345
Ala	Gln	Asp	Pro	Gly	Leu	Gln	Pro	Arg	Gly	Thr	Thr	Pro	Gly	Val
				350					355					360
Trp	Asn	Pro	Glu	Asn										
				365										

<210> 3

<211> 416

<212> PRT

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 1726609CD1

<400> 3

Met	Trp	Gly	Arg	Tyr	Asp	Ile	Val	Phe	Leu	Pro	Pro	Ser	Phe	Pro
1				5					10					15
Ile	Val	Ala	Met	Glu	Asn	Pro	Cys	Leu	Thr	Phe	Ile	Ile	Ser	Ser
				20					25					30
Ile	Leu	Glu	Ser	Asp	Glu	Phe	Leu	Val	Ile	Asp	Val	Ile	His	Glu
				35					40					45
Val	Ala	His	Ser	Trp	Phe	Gly	Asn	Ala	Val	Thr	Asn	Ala	Thr	Trp
				50					55					60
Glu	Glu	Met	Trp	Leu	Ser	Glu	Gly	Leu	Ala	Thr	Tyr	Ala	Gln	Arg
				65					70					75
Arg	Ile	Thr	Thr	Glu	Thr	Tyr	Gly	Ala	Ala	Phe	Thr	Cys	Leu	Glu
				80					85					90
Thr	Ala	Phe	Arg	Leu	Asp	Ala	Leu	His	Arg	Gln	Met	Lys	Leu	Leu
				95					100					105
Gly	Glu	Asp	Ser	Pro	Val	Ser	Lys	Leu	Gln	Val	Lys	Leu	Glu	Pro
				110					115					120
Gly	Val	Asn	Pro	Ser	His	Leu	Met	Asn	Leu	Phe	Thr	Tyr	Glu	Lys
				125					130					135
Gly	Tyr	Cys	Phe	Val	Tyr	Tyr	Leu	Ser	Gln	Leu	Cys	Gly	Asp	Pro
				140					145					150
Gln	Arg	Phe	Asp	Asp	Phe	Leu	Arg	Ala	Tyr	Val	Glu	Lys	Tyr	Lys
				155					160					165
Phe	Thr	Ser	Val	Val	Ala	Gln	Asp	Leu	Leu	Asp	Ser	Phe	Leu	Ser
				170					175					180
Phe	Phe	Pro	Glu	Leu	Lys	Glu	Gln	Ser	Val	Asp	Cys	Arg	Ala	Gly
				185					190					195
Leu	Glu	Phe	Glu	Arg	Trp	Leu	Asn	Ala	Thr	Gly	Pro	Pro	Leu	Ala
				200					205					210
Glu	Pro	Asp	Leu	Ser	Gln	Gly	Ser	Ser	Leu	Thr	Arg	Pro	Val	Glu
				215					220					225
Ala	Leu	Phe	Gln	Leu	Trp	Thr	Ala	Glu	Pro	Leu	Asp	Gln	Ala	Ala
				230					235					240
Ala	Ser	Ala	Ser	Ala	Ile	Asp	Ile	Ser	Lys	Trp	Arg	Thr	Phe	Gln
				245					250					255
Thr	Ala	Leu	Phe	Leu	Asp	Arg	Leu	Leu	Asp	Gly	Ser	Pro	Leu	Pro
				260					265					270
Gln	Glu	Val	Val	Met	Ser	Leu	Ser	Lys	Cys	Tyr	Ser	Ser	Leu	Leu
				275					280					285
Asp	Ser	Met	Asn	Ala	Glu	Ile	Arg	Ile	Arg	Trp	Leu	Gln	Ile	Val
				290					295					300
Val	Arg	Asn	Asp	Tyr	Tyr	Pro	Asp	Leu	His	Arg	Val	Arg	Arg	Phe
				305					310					315
Leu	Glu	Ser	Gln	Met	Ser	Arg	Met	Tyr	Thr	Ile	Pro	Leu	Tyr	Glu
				320					325					330
Asp	Leu	Cys	Thr	Gly	Ala	Leu	Lys	Ser	Phe	Ala	Leu	Glu	Val	Phe
				335					340					345

Tyr	Gln	Thr	Gln	Gly	Arg	Leu	His	Pro	Asn	Leu	Arg	Arg	Ala	Ile	
				350					355					360	
Gln	Gln	Ile	Leu	Ser	Gln	Gly	Leu	Gly	Ser	Ser	Thr	Glu	Pro	Ala	
				365					370					375	
Ser	Glu	Pro	Ser	Thr	Glu	Leu	Gly	Lys	Ala	Glu	Ala	Asp	Thr	Asp	
				380					385					390	
Ser	Asp	Ala	Gln	Ala	Leu	Leu	Leu	Gly	Asp	Glu	Ala	Pro	Ser	Ser	
				395					400					405	
Ala	Ile	Ser	Leu	Arg	Asp	Val	Asn	Val	Ser	Ala					
				410					415						

<210> 4

<211> 714

<212> PRT

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 4503848CD1

<400> 4

Met	His	Ile	His	Met	Leu	Thr	Leu	Asp	Gln	Gln	Lys	Ser	Leu	Ile	
1				5					10					15	
Leu	Ile	Leu	Phe	Leu	Ile	Leu	Phe	Arg	Val	Gly	Gly	Ser	Arg	Ile	
				20					25					30	
Leu	Leu	Arg	Met	Thr	Leu	Gly	Arg	Glu	Val	Met	Ser	Pro	Leu	Gln	
				35					40					45	
Ala	Met	Ser	Ser	Tyr	Thr	Val	Ala	Gly	Arg	Asn	Val	Leu	Arg	Trp	
				50					55					60	
Asp	Leu	Ser	Pro	Glu	Gln	Ile	Lys	Thr	Arg	Thr	Glu	Glu	Leu	Ile	
				65					70					75	
Val	Gln	Thr	Lys	Gln	Val	Tyr	Asp	Ala	Val	Gly	Met	Leu	Gly	Ile	
				80					85					90	
Glu	Glu	Val	Thr	Tyr	Glu	Asn	Cys	Leu	Gln	Ala	Leu	Ala	Asp	Val	
				95					100					105	
Glu	Val	Lys	Tyr	Ile	Val	Glu	Arg	Thr	Met	Leu	Asp	Phe	Pro	Gln	
				110					115					120	
His	Val	Ser	Ser	Asp	Lys	Glu	Val	Arg	Ala	Ala	Ser	Thr	Glu	Ala	
				125					130					135	
Asp	Lys	Arg	Leu	Ser	Arg	Phe	Asp	Ile	Glu	Met	Ser	Met	Arg	Gly	
				140					145					150	
Asp	Ile	Phe	Glu	Arg	Ile	Val	His	Leu	Gln	Glu	Thr	Cys	Asp	Leu	
				155					160					165	
Gly	Lys	Ile	Lys	Pro	Glu	Ala	Arg	Arg	Tyr	Leu	Glu	Lys	Ser	Ile	
				170					175					180	
Lys	Met	Gly	Lys	Arg	Asn	Gly	Leu	His	Leu	Pro	Glu	Gln	Val	Gln	
				185					190					195	
Asn	Glu	Ile	Lys	Ser	Met	Lys	Lys	Arg	Met	Ser	Glu	Leu	Cys	Ile	
				200					205					210	
Asp	Phe	Asn	Lys	Asn	Leu	Asn	Glu	Asp	Asp	Thr	Phe	Leu	Val	Phe	
				215					220					225	
Ser	Lys	Ala	Glu	Leu	Gly	Ala	Leu	Pro	Asp	Asp	Phe	Ile	Asp	Ser	
				230					235					240	
Leu	Glu	Lys	Thr	Asp	Asp	Asp	Lys	Tyr	Lys	Ile	Thr	Leu	Lys	Tyr	
				245					250					255	
Pro	His	Tyr	Phe	Pro	Val	Met	Lys	Lys	Cys	Cys	Ile	Pro	Glu	Thr	
				260					265					270	
Arg	Arg	Arg	Met	Glu	Met	Ala	Phe	Asn	Thr	Arg	Cys	Lys	Glu	Glu	
				275					280					285	
Asn	Thr	Ile	Ile	Leu	Gln	Gln	Leu	Leu	Pro	Leu	Arg	Thr	Lys	Val	
				290					295					300	
Ala	Lys	Leu	Leu	Gly	Tyr	Ser	Thr	His	Ala	Asp	Phe	Val	Leu	Glu	
				305					310					315	
Met	Asn	Thr	Ala	Lys	Ser	Thr	Ser	Arg	Val	Thr	Ala	Phe	Leu	Asp	
				320					325					330	
Asp	Leu	Ser	Gln	Lys	Leu	Lys	Pro	Leu	Gly	Glu	Ala	Glu	Arg	Glu	
				335					340					345	

```

Phe Ile Leu Asn Leu Lys Lys Lys Glu Cys Lys Asp Arg Gly Phe
350 355 360
Glu Tyr Asp Gly Lys Ile Asn Ala Trp Asp Leu Tyr Tyr Tyr Met
365 370 375
Thr Gln Thr Glu Glu Leu Lys Tyr Ser Ile Asp Gln Glu Phe Leu
380 385 390
Lys Glu Tyr Phe Pro Ile Glu Val Val Thr Glu Gly Leu Leu Asn
395 400 405
Thr Tyr Gln Glu Leu Leu Gly Leu Ser Phe Glu Gln Met Thr Asp
410 415 420
Ala His Val Trp Asn Lys Ser Val Thr Leu Tyr Thr Val Lys Asp
425 430 435
Lys Ala Thr Gly Glu Val Leu Gly Gln Phe Tyr Leu Asp Leu Tyr
440 445 450
Pro Arg Glu Gly Lys Tyr Asn His Ala Ala Cys Phe Gly Leu Gln
455 460 465
Pro Gly Cys Leu Leu Pro Asp Gly Ser Arg Met Met Ala Val Ala
470 475 480
Ala Leu Val Val Asn Phe Ser Gln Pro Val Ala Gly Arg Pro Ser
485 490 495
Leu Leu Arg His Asp Glu Val Arg Thr Tyr Phe His Glu Phe Gly
500 505 510
His Val Met His Gln Ile Cys Ala Gln Thr Asp Phe Ala Arg Phe
515 520 525
Ser Gly Thr Asn Val Glu Thr Asp Phe Val Glu Val Pro Ser Gln
530 535 540
Met Leu Glu Asn Trp Val Trp Asp Val Asp Ser Leu Arg Arg Leu
545 550 555
Ser Lys His Tyr Lys Asp Gly Ser Pro Ile Ala Asp Asp Leu Leu
560 565 570
Glu Lys Leu Val Ala Ser Arg Leu Val Asn Thr Gly Leu Leu Thr
575 580 585
Leu Arg Gln Ile Val Leu Ser Lys Val Asp Gln Ser Leu His Thr
590 595 600
Asn Thr Ser Leu Asp Ala Ala Ser Glu Tyr Ala Lys Tyr Cys Ser
605 610 615
Glu Ile Leu Gly Val Ala Ala Thr Pro Gly Thr Asn Met Pro Ala
620 625 630
Thr Phe Gly His Leu Ala Gly Gly Tyr Asp Gly Gln Tyr Tyr Gly
635 640 645
Tyr Leu Trp Ser Glu Val Phe Ser Met Asp Met Phe Tyr Ser Cys
650 655 660
Phe Lys Lys Glu Gly Ile Met Asn Pro Glu Val Gly Met Lys Tyr
665 670 675
Arg Asn Leu Ile Leu Lys Pro Gly Gly Ser Leu Asp Gly Met Asp
680 685 690
Met Leu His Asn Phe Leu Lys Arg Glu Pro Asn Gln Lys Ala Phe
695 700 705
Leu Met Ser Arg Gly Leu His Ala Pro
710

```

<210> 5

<211> 367

<212> PRT

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 5544089CD1

<400> 5

```

Met Phe Ala Pro Ser Val Leu Ser Ser Gly Leu Ser Gly Gly Ala
1 5 10 15
Ser Lys Gly Arg Lys Met Glu Leu Ile Gln Pro Lys Glu Pro Thr
20 25 30
Ser Gln Tyr Ile Ser Leu Cys His Glu Leu His Thr Leu Phe Gln
35 40 45

```

Val Met Trp Ser Gly Lys Trp Ala Leu Val Ser Pro Phe Ala Met	
50	55
Leu His Ser Val Trp Arg Leu Ile Pro Ala Phe Arg Gly Tyr Ala	
65	70
Gln Gln Asp Ala Gln Glu Phe Leu Cys Glu Leu Leu Asp Lys Ile	
80	85
Gln Arg Glu Leu Glu Thr Thr Gly Thr Ser Leu Pro Ala Leu Ile	
95	100
Pro Thr Ser Gln Arg Lys Leu Ile Lys Gln Val Leu Asn Val Val	
110	115
Asn Asn Ile Phe His Gly Gln Leu Leu Ser Gln Val Thr Cys Leu	
125	130
Ala Cys Asp Asn Lys Ser Asn Thr Ile Glu Pro Phe Trp Asp Leu	
140	145
Ser Leu Glu Phe Pro Glu Arg Tyr Gln Cys Ser Gly Lys Asp Ile	
155	160
Ala Ser Gln Pro Cys Leu Val Thr Glu Met Leu Ala Lys Phe Thr	
170	175
Glu Thr Glu Ala Leu Glu Gly Lys Ile Tyr Val Cys Asp Gln Cys	
185	190
Asn Ser Lys Arg Arg Arg Phe Ser Ser Lys Pro Val Val Leu Thr	
200	205
Glu Ala Gln Lys Gln Leu Met Ile Cys His Leu Pro Gln Val Leu	
215	220
Arg Leu His Leu Lys Arg Phe Arg Trp Ser Gly Arg Asn Asn Arg	
230	235
Glu Lys Ile Gly Val His Val Gly Phe Glu Glu Ile Leu Asn Met	
245	250
Glu Pro Tyr Cys Cys Arg Glu Thr Leu Lys Ser Leu Arg Pro Glu	
260	265
Cys Phe Ile Tyr Asp Leu Ser Ala Val Val Met His His Gly Lys	
275	280
Gly Phe Gly Ser Gly His Tyr Thr Ala Tyr Cys Tyr Asn Ser Glu	
290	295
Gly Gly Phe Trp Val His Cys Asn Asp Ser Lys Leu Ser Met Cys	
305	310
Thr Met Asp Glu Val Cys Lys Ala Gln Ala Tyr Ile Leu Phe Tyr	
320	325
Thr Gln Arg Val Thr Glu Asn Gly His Ser Lys Leu Leu Pro Pro	
335	340
Glu Leu Leu Leu Gly Ser Gln His Pro Asn Glu Asp Ala Asp Thr	
350	355
Ser Ser Asn Glu Ile Leu Ser	
365	

<210> 6

<211> 235

<212> PRT

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 7474081CD1

<400> 6

Met Lys Tyr Val Phe Tyr Leu Gly Val Leu Ala Gly Thr Phe Phe	
1	10
Phe Ala Asp Ser Ser Val Gln Lys Glu Asp Pro Ala Pro Tyr Leu	
20	25
Val Tyr Leu Lys Ser His Phe Asn Pro Cys Val Gly Val Leu Ile	
35	40
Lys Pro Ser Trp Val Leu Ala Pro Ala His Cys Tyr Leu Pro Asn	
50	55
Leu Lys Val Met Leu Gly Asn Phe Lys Ser Arg Val Arg Asp Gly	
65	70
Thr Glu Gln Thr Ile Asn Pro Ile Gln Ile Val Arg Tyr Trp Asn	
80	85
	90

Tyr Ser His Ser Ala Pro Gln Asp Asp Leu Met Leu Ile Lys Leu
 95 100 105
 Ala Lys Pro Ala Met Leu Asn Pro Lys Val Gln Pro Leu Thr Leu
 110 115 120
 Ala Thr Thr Asn Val Arg Pro Gly Thr Val Cys Leu Leu Ser Gly
 125 130 135
 Leu Asp Trp Ser Gln Glu Asn Ser Gly Arg His Pro Asp Leu Arg
 140 145 150
 Gln Asn Leu Glu Ala Pro Val Met Ser Asp Arg Glu Cys Gln Lys
 155 160 165
 Thr Glu Gln Gly Lys Ser His Arg Asn Ser Leu Cys Val Lys Phe
 170 175 180
 Val Lys Val Phe Ser Arg Ile Phe Gly Glu Val Ala Val Ala Thr
 185 190 195
 Val Ile Cys Lys Asp Lys Leu Gln Gly Ile Glu Val Gly His Phe
 200 205 210
 Met Gly Gly Asp Val Gly Ile Tyr Thr Asn Val Tyr Lys Tyr Val
 215 220 225
 Ser Trp Ile Glu Asn Thr Ala Lys Asp Lys
 230 235

<210> 7

<211> 488

<212> PRT

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 5281209CD1

<400> 7

Met Gln Pro Thr Gly Arg Glu Gly Ser Arg Ala Leu Ser Arg Arg
 1 5 10 15
 Tyr Leu Arg Arg Leu Leu Leu Leu Leu Leu Leu Leu Arg
 20 25 30
 Gln Pro Val Thr Arg Ala Glu Thr Thr Pro Gly Ala Pro Arg Ala
 35 40 45
 Leu Ser Thr Leu Gly Ser Pro Ser Leu Phe Thr Thr Pro Gly Val
 50 55 60
 Pro Ser Ala Leu Thr Thr Pro Gly Leu Thr Thr Pro Gly Thr Pro
 65 70 75
 Lys Thr Leu Asp Leu Arg Gly Arg Ala Gln Ala Leu Met Arg Ser
 80 85 90
 Phe Pro Leu Val Asp Gly His Asn Asp Leu Pro Gln Val Leu Arg
 95 100 105
 Gln Arg Tyr Lys Asn Val Leu Gln Asp Val Asn Leu Arg Asn Phe
 110 115 120
 Ser His Gly Gln Thr Ser Leu Asp Arg Leu Arg Asp Gly Leu Val
 125 130 135
 Gly Ala Gln Phe Trp Ser Ala Ser Val Ser Cys Gln Ser Gln Asp
 140 145 150
 Gln Thr Ala Val Arg Leu Ala Leu Glu Gln Ile Asp Leu Ile His
 155 160 165
 Arg Met Cys Ala Ser Tyr Ser Glu Leu Glu Leu Val Thr Ser Ala
 170 175 180
 Glu Gly Leu Asn Ser Ser Gln Lys Leu Ala Cys Leu Ile Gly Val
 185 190 195
 Glu Gly Gly His Ser Leu Asp Ser Ser Leu Ser Val Leu Arg Ser
 200 205 210
 Phe Tyr Val Leu Gly Val Arg Tyr Leu Thr Leu Thr Phe Thr Cys
 215 220 225
 Ser Thr Pro Trp Ala Glu Ser Ser Thr Lys Phe Arg His His Met
 230 235 240
 Tyr Thr Asn Val Ser Gly Leu Thr Ser Phe Gly Glu Lys Val Val
 245 250 255
 Glu Glu Leu Asn Arg Leu Gly Met Met Ile Asp Leu Ser Tyr Ala
 260 265 270

```
<210> 8
<211> 346
<212> PRT
<213> Homo sapiens
```

<400> 8														
Met	Leu	Leu	Gly	Arg	Val	Trp	Gln	Thr	Arg	Glu	Leu	Lys	Ser	Lys
1				5					10					15
Val	Pro	Lys	Lys	Ala	Gly	Arg	Cys	Gly	Gln	Gly	Arg	Leu	His	Gly
				20					25					30
Gly	Ser	Ala	Val	Gly	Phe	Leu	Gly	Ser	Pro	Pro	Gly	Thr	Pro	Ser
				35					40					45
Ser	Phe	Asp	Leu	Gly	Cys	Gly	Arg	Pro	Gln	Val	Ser	Asp	Ala	Gly
				50					55					60
Gly	Arg	Ile	Val	Gly	Gly	His	Ala	Ala	Pro	Ala	Gly	Ala	Trp	Pro
				65					70					75
Trp	Gln	Ala	Ser	Leu	Arg	Leu	Arg	Arg	Val	His	Val	Cys	Gly	Gly
				80					85					90
Ser	Leu	Leu	Ser	Pro	Gln	Trp	Val	Leu	Thr	Ala	Ala	His	Cys	Phe
				95					100					105
Ser	Gly	Ser	Leu	Asn	Ser	Ser	Asp	Tyr	Gln	Val	His	Leu	Gly	Glu
				110					115					120
Leu	Glu	Ile	Thr	Leu	Ser	Pro	His	Phe	Ser	Thr	Val	Arg	Gln	Ile
				125					130					135
Ile	Leu	His	Ser	Ser	Pro	Ser	Gly	Gln	Pro	Gly	Thr	Ser	Gly	Asp
				140					145					150
Ile	Ala	Leu	Val	Glu	Leu	Ser	Val	Pro	Val	Thr	Leu	Phe	Ser	Arg
				155					160					165
Ile	Leu	Pro	Val	Cys	Leu	Pro	Glu	Ala	Ser	Asp	Asp	Phe	Cys	Pro
				170					175					180
Gly	Ile	Arg	Cys	Trp	Val	Thr	Gly	Trp	Gly	Tyr	Thr	Arg	Glu	Gly
				185					190					195

Glu	Pro	Leu	Pro	Pro	Pro	Tyr	Ser	Leu	Arg	Glu	Val	Lys	Val	Ser
				200					205					210
Val	Val	Asp	Thr	Glu	Thr	Cys	Arg	Arg	Asp	Tyr	Pro	Gly	Pro	Gly
				215					220					225
Gly	Ser	Ile	Leu	Gln	Pro	Asp	Met	Leu	Cys	Ala	Arg	Gly	Pro	Gly
				230					235					240
Asp	Ala	Cys	Gln	Asp	Asp	Ser	Gly	Gly	Pro	Leu	Val	Cys	Gln	Val
				245					250					255
Asn	Gly	Ala	Trp	Val	Gln	Ala	Gly	Ile	Val	Ser	Trp	Gly	Glu	Gly
				260					265					270
Cys	Gly	Arg	Pro	Asn	Arg	Pro	Gly	Val	Tyr	Thr	Arg	Val	Pro	Ala
				275					280					285
Tyr	Val	Asn	Trp	Ile	Arg	Arg	His	Ile	Thr	Ala	Ser	Gly	Gly	Ser
				290					295					300
Glu	Ser	Gly	Tyr	Pro	Arg	Leu	Pro	Leu	Leu	Ala	Gly	Leu	Phe	Leu
				305					310					315
Pro	Gly	Leu	Phe	Leu	Leu	Val	Ser	Cys	Val	Leu	Leu	Ala	Lys	
				320					325					330
Cys	Leu	Leu	His	Pro	Ser	Ala	Asp	Gly	Thr	Pro	Phe	Pro	Ala	Pro
				335					340					345

Asp

<210> 9

<211> 882

<212> PRT

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 7160544CD1

<400> 9

Met	Ala	Ala	Ala	Met	Glu	Thr	Glu	Gln	Leu	Gly	Val	Glu	Ile	Phe
1				5					10					15
Glu	Thr	Ala	Asp	Cys	Glu	Glu	Asn	Ile	Glu	Ser	Gln	Asp	Arg	Pro
				20					25					30
Lys	Leu	Glu	Pro	Phe	Tyr	Val	Glu	Arg	Tyr	Ser	Trp	Ser	Gln	Leu
				35					40					45
Lys	Lys	Leu	Leu	Ala	Asp	Thr	Arg	Lys	Tyr	His	Gly	Tyr	Met	Met
				50					55					60
Ala	Lys	Ala	Pro	His	Asp	Phe	Met	Phe	Val	Lys	Arg	Asn	Asp	Pro
				65					70					75
Asp	Gly	Pro	His	Ser	Asp	Arg	Ile	Tyr	Tyr	Leu	Ala	Met	Ser	Gly
				80					85					90
Glu	Asn	Arg	Glu	Asn	Thr	Leu	Phe	Tyr	Ser	Glu	Ile	Pro	Lys	Thr
				95					100					105
Ile	Asn	Arg	Ala	Ala	Val	Leu	Met	Leu	Ser	Trp	Lys	Pro	Leu	Leu
				110					115					120
Asp	Leu	Phe	Gln	Ala	Thr	Leu	Asp	Tyr	Gly	Met	Tyr	Ser	Arg	Glu
				125					130					135
Glu	Glu	Leu	Leu	Arg	Glu	Arg	Lys	Arg	Ile	Gly	Thr	Val	Gly	Ile
				140					145					150
Ala	Ser	Tyr	Asp	Tyr	His	Gln	Gly	Ser	Gly	Thr	Phe	Leu	Phe	Gln
				155					160					165
Ala	Gly	Ser	Gly	Ile	Tyr	His	Val	Lys	Asp	Gly	Gly	Pro	Gln	Gly
				170					175					180
Phe	Thr	Gln	Gln	Pro	Leu	Arg	Pro	Asn	Leu	Val	Glu	Thr	Ser	Cys
				185					190					195
Pro	Asn	Ile	Arg	Met	Asp	Pro	Lys	Leu	Cys	Pro	Ala	Asp	Pro	Asp
				200					205					210
Trp	Ile	Ala	Phe	Ile	His	Ser	Asn	Asp	Ile	Trp	Ile	Ser	Asn	Ile
				215					220					225
Val	Thr	Arg	Glu	Glu	Arg	Arg	Leu	Thr	Tyr	Val	His	Asn	Glu	Leu
				230					235					240
Ala	Asn	Met	Glu	Glu	Asp	Ala	Arg	Ser	Ala	Gly	Val	Ala	Thr	Phe
				245					250					255

Val	Leu	Gln	Glu	Glu	Phe	Asp	Arg	Tyr	Ser	Gly	Tyr	Trp	Trp	Cys
				260					265					270
Pro	Lys	Ala	Glu	Thr	Thr	Pro	Ser	Gly	Gly	Lys	Ile	Leu	Arg	Ile
				275					280					285
Leu	Tyr	Glu	Glu	Asn	Asp	Glu	Ser	Glu	Val	Glu	Ile	Ile	His	Val
				290					295					300
Thr	Ser	Pro	Met	Leu	Glu	Thr	Arg	Arg	Ala	Asp	Ser	Phe	Arg	Tyr
				305					310					315
Pro	Lys	Thr	Gly	Thr	Ala	Asn	Pro	Lys	Val	Thr	Phe	Lys	Met	Ser
				320					325					330
Glu	Ile	Met	Ile	Asp	Ala	Glu	Gly	Arg	Ile	Ile	Asp	Val	Ile	Asp
				335					340					345
Lys	Glu	Leu	Ile	Gln	Pro	Phe	Glu	Ile	Leu	Phe	Glu	Gly	Val	Glu
				350					355					360
Tyr	Ile	Ala	Arg	Ala	Gly	Trp	Thr	Pro	Glu	Gly	Lys	Tyr	Ala	Trp
				365					370					375
Ser	Ile	Leu	Leu	Asp	Arg	Ser	Gln	Thr	Arg	Leu	Gln	Ile	Val	Leu
				380					385					390
Ile	Ser	Pro	Glu	Leu	Phe	Ile	Pro	Val	Glu	Asp	Asp	Val	Met	Glu
				395					400					405
Arg	Gln	Arg	Leu	Ile	Glu	Ser	Val	Pro	Asp	Ser	Val	Thr	Pro	Leu
				410					415					420
Ile	Ile	Tyr	Glu	Glu	Thr	Thr	Asp	Ile	Trp	Ile	Asn	Ile	His	Asp
				425					430					435
Ile	Phe	His	Val	Phe	Pro	Gln	Ser	His	Glu	Glu	Glu	Ile	Glu	Phe
				440					445					450
Ile	Phe	Ala	Ser	Glu	Cys	Lys	Thr	Gly	Phe	Arg	His	Leu	Tyr	Lys
				455					460					465
Ile	Thr	Ser	Ile	Leu	Lys	Glu	Ser	Lys	Tyr	Lys	Arg	Ser	Ser	Gly
				470					475					480
Gly	Leu	Pro	Ala	Pro	Ser	Asp	Phe	Lys	Cys	Pro	Ile	Lys	Glu	Glu
				485					490					495
Ile	Ala	Ile	Thr	Ser	Gly	Glu	Trp	Glu	Val	Leu	Gly	Arg	His	Gly
				500					505					510
Ser	Asn	Ile	Gln	Val	Asp	Glu	Val	Arg	Arg	Leu	Val	Tyr	Phe	Glu
				515					520					525
Gly	Thr	Lys	Asp	Ser	Pro	Leu	Glu	His	His	Leu	Tyr	Val	Val	Ser
				530					535					540
Tyr	Val	Asn	Pro	Gly	Glu	Val	Thr	Arg	Leu	Thr	Asp	Arg	Gly	Tyr
				545					550					555
Ser	His	Ser	Cys	Cys	Ile	Ser	Gln	His	Cys	Asp	Phe	Phe	Ile	Ser
				560					565					570
Lys	Tyr	Ser	Asn	Gln	Lys	Asn	Pro	His	Cys	Val	Ser	Leu	Tyr	Lys
				575					580					585
Leu	Ser	Ser	Pro	Glu	Asp	Asp	Pro	Thr	Cys	Lys	Thr	Lys	Glu	Phe
				590					595					600
Trp	Ala	Thr	Ile	Leu	Asp	Ser	Ala	Gly	Pro	Leu	Pro	Asp	Tyr	Thr
				605					610					615
Pro	Pro	Glu	Ile	Phe	Ser	Phe	Glu	Ser	Thr	Thr	Gly	Phe	Thr	Leu
				620					625					630
Tyr	Gly	Met	Leu	Tyr	Lys	Pro	His	Asp	Leu	Gln	Pro	Gly	Lys	Lys
				635					640					645
Tyr	Pro	Thr	Val	Leu	Phe	Ile	Tyr	Gly	Gly	Pro	Gln	Val	Gln	Leu
				650					655					660
Val	Asn	Asn	Arg	Phe	Lys	Gly	Val	Lys	Tyr	Phe	Arg	Leu	Asn	Thr
				665					670					675
Leu	Ala	Ser	Leu	Gly	Tyr	Val	Val	Val	Val	Ile	Asp	Asn	Arg	Gly
				680					685					690
Ser	Cys	His	Arg	Gly	Leu	Lys	Phe	Glu	Gly	Ala	Phe	Lys	Tyr	Lys
				695					700					705
Met	Gly	Gln	Ile	Glu	Ile	Asp	Asp	Gln	Val	Glu	Gly	Leu	Gln	Tyr
				710					715					720
Leu	Ala	Ser	Arg	Tyr	Asp	Phe	Ile	Asp	Leu	Asp	Arg	Val	Gly	Ile
				725					730					735
His	Gly	Trp	Ser	Tyr	Gly	Gly	Tyr	Leu	Ser	Leu	Met	Ala	Leu	Met
				740					745					750
Gln	Arg	Ser	Asp	Ile	Phe	Arg	Val	Ala	Ile	Ala	Gly	Ala	Pro	Val

	755		760		765
Thr Leu Trp Ile	Phe Tyr Asp Thr Gly	Tyr Thr Glu Arg Tyr	Met		
	770	775			
Gly His Pro Asp	Gln Asn Glu Gln Gly	Tyr Tyr Leu Gly Ser	Val		
	785	790			
Ala Met Gln Ala	Glu Lys Phe Pro Ser	Glu Pro Asn Arg Leu	Leu		
	800	805			
Leu Leu His Gly	Phe Leu Asp Glu Asn	Val His Phe Ala His	Thr		
	815	820			
Ser Ile Leu Leu	Ser Phe Leu Val Arg	Ala Gly Lys Pro Tyr	Asp		
	830	835			
Leu Gln Ile Tyr	Pro Gln Glu Arg His	Ser Ile Arg Val Pro	Glu		
	845	850			
Ser Gly Glu His	Tyr Glu Leu His Leu	Leu His Tyr Leu Gln	Glu		
	860	865			
Asn Leu Gly Ser	Arg Ile Ala Ala Leu	Lys Val Ile			
	875	880			

<210> 10

<211> 1189

<212> PRT

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 7477386CD1

<400> 10

Met Ala Pro Leu Arg	Ala Leu Leu Ser Tyr	Leu Leu Pro Leu His
1	5	10
Cys Ala Leu Cys Ala	Ala Ala Gly Ser Arg	Thr Pro Glu Leu His
	20	25
Leu Ser Gly Lys Leu	Ser Asp Tyr Gly Val	Thr Val Pro Cys Ser
	35	40
Thr Asp Phe Arg Gly	Arg Phe Leu Ser His	Val Val Ser Gly Pro
	50	55
Ala Ala Ala Ser Ala	Gly Ser Met Val Val	Asp Thr Pro Pro Thr
	65	70
Leu Pro Arg His Ser	Ser His Leu Arg Val	Ala Arg Ser Pro Leu
	80	85
His Pro Gly Gly Thr	Leu Trp Pro Gly Arg	Val Gly Arg His Ser
	95	100
Leu Tyr Phe Asn Val	Thr Val Phe Gly Lys	Glu Leu His Leu Arg
	110	115
Leu Arg Pro Asn Arg	Arg Leu Val Val Pro	Gly Ser Ser Val Glu
	125	130
Trp Gln Glu Asp Phe	Arg Glu Leu Phe Arg	Gln Pro Leu Arg Gln
	140	145
Glu Cys Val Tyr Thr	Gly Gly Val Thr Gly	Met Pro Gly Ala Ala
	155	160
Val Ala Ile Ser Asn	Cys Asp Gly Leu Ala	Gly Leu Ile Arg Thr
	170	175
Asp Ser Thr Asp Phe	Phe Ile Glu Pro Leu	Glu Arg Gly Gln Gln
	185	190
Glu Lys Glu Ala Ser	Gly Arg Thr His Val	Val Tyr Arg Arg Glu
	200	205
Ala Val Gln Gln Glu	Trp Ala Glu Pro Asp	Gly Asp Leu His Asn
	215	220
Glu Ala Phe Gly Leu	Gly Asp Leu Pro Asn	Leu Leu Gly Leu Val
	230	235
Gly Asp Gln Leu Gly	Asp Thr Glu Arg Lys	Arg Arg His Ala Lys
	245	250
Pro Gly Ser Tyr Ser	Ile Glu Val Leu Leu	Val Val Asp Asp Ser
	260	265
Val Val Arg Phe His	Gly Lys Glu His Val	Gln Asn Tyr Val Leu
	275	280
Thr Leu Met Asn Ile	Val Val Asp Glu Ile	Tyr His Asp Glu Ser

	290		295		300
Leu Gly Val His	Ile Asn Ile Ala Leu	Val Arg Leu Ile Met	Val		
	305		310		315
Gly Tyr Arg Gln	Gln Ser Leu Ser Leu	Ile Glu Arg Gly Asn	Pro		
	320		325		330
Ser Arg Ser Leu	Glu Gln Val Cys Arg	Trp Ala His Ser Gln	Gln		
	335		340		345
Arg Gln Asp Pro	Ser His Ala Glu His	His Asp His Val Val	Phe		
	350		355		360
Leu Thr Arg Gln	Asp Phe Gly Pro Ser	Gly Gly Tyr Ala Pro	Val		
	365		370		375
Thr Gly Met Cys	His Pro Leu Arg Ser	Cys Ala Leu Asn His	Glu		
	380		385		390
Asp Gly Phe Ser	Ser Ala Phe Val Ile	Ala His Glu Thr Gly	His		
	395		400		405
Val Leu Gly Met	Glu His Asp Gly Gln	Gly Asn Gly Cys Ala	Asp		
	410		415		420
Glu Thr Ser Leu	Gly Ser Val Met Ala	Pro Leu Val Gln Ala	Ala		
	425		430		435
Phe His Arg Phe	His Trp Ser Arg Cys	Ser Lys Leu Glu Leu	Ser		
	440		445		450
Arg Tyr Leu Pro	Ser Tyr Asp Cys Leu	Leu Asp Asp Pro Phe	Asp		
	455		460		465
Pro Ala Trp Pro	Gln Pro Pro Glu Leu	Pro Gly Ile Asn Tyr	Ser		
	470		475		480
Met Asp Glu Gln	Cys Arg Phe Asp Phe	Gly Ser Gly Tyr Gln	Thr		
	485		490		495
Cys Leu Ala Phe	Arg Thr Phe Glu Pro	Cys Lys Gln Leu Trp	Cys		
	500		505		510
Ser His Pro Asp	Asn Pro Tyr Phe Cys	Lys Thr Lys Lys Gly	Pro		
	515		520		525
Pro Leu Asp Gly	Thr Glu Cys Ala Pro	Gly Lys Trp Cys Phe	Lys		
	530		535		540
Gly His Cys Ile	Trp Lys Ser Pro Glu	Gln Thr Tyr Gly Gln	Asp		
	545		550		555
Gly Gly Trp Ser	Ser Trp Thr Lys Phe	Gly Ser Cys Ser Arg	Ser		
	560		565		570
Cys Gly Gly Gly	Val Arg Ser Arg Ser	Arg Ser Cys Asn Asn	Pro		
	575		580		585
Ser Pro Ala Tyr	Gly Gly Arg Leu Cys	Leu Gly Pro Met Phe	Glu		
	590		595		600
Tyr Gln Val Cys	Asn Ser Glu Glu Cys	Pro Gly Thr Tyr Glu	Asp		
	605		610		615
Phe Arg Ala Gln	Gln Cys Ala Lys Arg	Asn Ser Tyr Tyr Val	His		
	620		625		630
Gln Asn Ala Lys	His Ser Trp Val Pro	Tyr Glu Pro Asp Asp	Asp		
	635		640		645
Ala Gln Lys Cys	Glu Leu Ile Cys Gln	Ser Ala Asp Thr Gly	Asp		
	650		655		660
Val Val Phe Met	Asn Gln Val Val His	Asp Gly Thr Arg Cys	Ser		
	665		670		675
Tyr Arg Asp Pro	Tyr Ser Val Cys Ala	Arg Gly Glu Cys Val	Pro		
	680		685		690
Val Gly Cys Asp	Lys Glu Val Gly Ser	Met Lys Ala Asp Asp	Lys		
	695		700		705
Cys Gly Val Cys	Gly Gly Asp Asn Ser	His Cys Arg Thr Val	Lys		
	710		715		720
Gly Thr Leu Gly	Lys Ala Ser Lys Gln	Ala Gly Ala Leu Lys	Leu		
	725		730		735
Val Gln Ile Pro	Ala Gly Ala Arg His	Ile Gln Ile Glu Ala	Leu		
	740		745		750
Glu Lys Ser Pro	His Arg Ile Val Val	Lys Asn Gln Val Thr	Gly		
	755		760		765
Ser Phe Ile Leu	Asn Pro Lys Gly Lys	Glu Ala Thr Ser Arg	Thr		
	770		775		780
Phe Thr Ala Met	Gly Leu Glu Trp Glu	Asp Ala Val Glu Asp	Ala		
	785		790		795

Lys	Glu	Ser	Leu	Lys	Thr	Ser	Gly	Pro	Leu	Pro	Glu	Ala	Ile	Ala
				800					805					810
Ile	Leu	Ala	Leu	Pro	Pro	Thr	Glu	Gly	Gly	Pro	Arg	Ser	Ser	Leu
				815					820					825
Ala	Tyr	Lys	Tyr	Val	Ile	His	Glu	Asp	Leu	Leu	Pro	Leu	Ile	Gly
				830					835					840
Ser	Asn	Asn	Val	Leu	Leu	Glu	Glu	Met	Asp	Thr	Tyr	Glu	Trp	Ala
				845					850					855
Leu	Lys	Ser	Trp	Ala	Pro	Cys	Ser	Lys	Ala	Cys	Gly	Gly	Gly	Ile
				860					865					870
Gln	Phe	Thr	Lys	Tyr	Gly	Cys	Arg	Arg	Arg	Arg	Asp	His	His	Met
				875					880					885
Val	Gln	Arg	His	Leu	Cys	Asp	His	Lys	Lys	Arg	Pro	Lys	Pro	Ile
				890					895					900
Arg	Arg	Arg	Cys	Asn	Gln	His	Pro	Cys	Ser	Gln	Pro	Val	Trp	Val
				905					910					915
Thr	Glu	Glu	Trp	Gly	Ala	Cys	Ser	Arg	Ser	Cys	Gly	Lys	Leu	Gly
				920					925					930
Val	Gln	Thr	Arg	Gly	Ile	Gln	Cys	Leu	Leu	Pro	Leu	Ser	Asn	Gly
				935					940					945
Thr	His	Lys	Val	Met	Pro	Ala	Lys	Ala	Cys	Ala	Gly	Asp	Arg	Pro
				950					955					960
Glu	Ala	Arg	Arg	Pro	Cys	Leu	Arg	Val	Pro	Cys	Pro	Ala	Gln	Trp
				965					970					975
Arg	Leu	Gly	Ala	Trp	Ser	Gln	Cys	Ser	Ala	Thr	Cys	Gly	Glu	Gly
				980					985					990
Ile	Gln	Gln	Arg	Gln	Val	Val	Cys	Arg	Thr	Asn	Ala	Asn	Ser	Leu
				995					1000					1005
Gly	His	Cys	Glu	Gly	Asp	Arg	Pro	Asp	Thr	Val	Gln	Val	Cys	Ser
				1010					1015					1020
Leu	Pro	Ala	Cys	Gly	Ala	Glu	Pro	Cys	Thr	Gly	Asp	Arg	Ser	Val
				1025					1030					1035
Phe	Cys	Gln	Met	Glu	Val	Leu	Asp	Arg	Tyr	Cys	Ser	Ile	Pro	Gly
				1040					1045					1050
Tyr	His	Arg	Leu	Cys	Cys	Val	Ser	Cys	Ile	Lys	Lys	Ala	Ser	Gly
				1055					1060					1065
Pro	Asn	Pro	Gly	Pro	Asp	Pro	Gly	Pro	Thr	Ser	Leu	Pro	Pro	Phe
				1070					1075					1080
Ser	Thr	Pro	Gly	Ser	Pro	Leu	Pro	Gly	Pro	Gln	Asp	Pro	Ala	Asp
				1085					1090					1095
Ala	Ala	Glu	Pro	Pro	Gly	Lys	Pro	Thr	Gly	Ser	Glu	Asp	His	Gln
				1100					1105					1110
His	Gly	Arg	Ala	Thr	Gln	Leu	Pro	Gly	Ala	Leu	Asp	Thr	Ser	Ser
				1115					1120					1125
Pro	Gly	Thr	Gln	His	Pro	Phe	Ala	Pro	Glu	Thr	Pro	Ile	Pro	Gly
				1130					1135					1140
Ala	Ser	Trp	Ser	Ile	Ser	Pro	Thr	Thr	Pro	Gly	Gly	Leu	Pro	Trp
				1145					1150					1155
Gly	Trp	Thr	Gln	Thr	Pro	Thr	Pro	Val	Pro	Glu	Asp	Lys	Gly	Gln
				1160					1165					1170
Pro	Gly	Glu	Asp	Leu	Arg	His	Pro	Gly	Thr	Ser	Leu	Pro	Ala	Ala
				1175					1180					1185
Ser	Pro	Val	Thr											

<210> 11

<211> 952

<212> PRT

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 7473089CD1

<400> 11

Met	Leu	Leu	Leu	Gly	Ile	Leu	Thr	Leu	Ala	Phe	Ala	Gly	Arg	Thr
1				5					10					15

Ala	Gly	Gly	Ser	Glu	Pro	Glu	Arg	Glu	Val	Val	Val	Pro	Ile	Arg	20	25	30
Leu	Asp	Pro	Asp	Ile	Asn	Gly	Arg	Arg	Tyr	Tyr	Trp	Arg	Gly	Pro	35	40	45
Glu	Asp	Ser	Gly	Asp	Gln	Gly	Leu	Ile	Phe	Gln	Ile	Thr	Ala	Phe	50	55	60
Gln	Glu	Asp	Phe	Tyr	Leu	His	Leu	Thr	Pro	Asp	Ala	Gln	Phe	Leu	65	70	75
Ala	Pro	Ala	Phe	Ser	Thr	Glu	His	Leu	Gly	Val	Pro	Leu	Gln	Gly	80	85	90
Leu	Thr	Gly	Gly	Ser	Ser	Asp	Leu	Arg	Arg	Cys	Phe	Tyr	Ser	Gly	95	100	105
Asp	Val	Asn	Ala	Glu	Pro	Asp	Ser	Phe	Ala	Ala	Val	Ser	Leu	Cys	110	115	120
Gly	Gly	Leu	Arg	Gly	Ala	Phe	Gly	Tyr	Arg	Gly	Ala	Glu	Tyr	Val	125	130	135
Ile	Ser	Pro	Leu	Pro	Asn	Ala	Ser	Ala	Pro	Ala	Ala	Gln	Arg	Asn	140	145	150
Ser	Gln	Gly	Ala	His	Leu	Leu	Gln	Arg	Arg	Gly	Val	Pro	Gly	Gly	155	160	165
Pro	Ser	Gly	Asp	Pro	Thr	Ser	Arg	Cys	Gly	Val	Ala	Ser	Gly	Trp	170	175	180
Asn	Pro	Ala	Ile	Leu	Arg	Ala	Leu	Asp	Pro	Tyr	Lys	Pro	Arg	Arg	185	190	195
Ala	Gly	Phe	Gly	Glu	Ser	Arg	Ser	Arg	Arg	Arg	Ser	Gly	Arg	Ala	200	205	210
Lys	Arg	Phe	Val	Ser	Ile	Pro	Arg	Tyr	Val	Glu	Thr	Leu	Val	Val	215	220	225
Ala	Asp	Glu	Ser	Met	Val	Lys	Phe	His	Gly	Ala	Asp	Leu	Glu	His	230	235	240
Tyr	Leu	Leu	Thr	Leu	Leu	Ala	Thr	Ala	Ala	Arg	Leu	Tyr	Arg	His	245	250	255
Pro	Ser	Ile	Leu	Asn	Pro	Ile	Asn	Ile	Val	Val	Val	Lys	Val	Leu	260	265	270
Leu	Leu	Arg	Asp	Arg	Asp	Ser	Gly	Pro	Lys	Val	Thr	Gly	Asn	Ala	275	280	285
Ala	Leu	Thr	Leu	Arg	Asn	Phe	Cys	Ala	Trp	Gln	Lys	Lys	Leu	Asn	290	295	300
Lys	Val	Ser	Asp	Lys	His	Pro	Glu	Tyr	Trp	Asp	Thr	Ala	Ile	Leu	305	310	315
Phe	Thr	Arg	Gln	Asp	Leu	Cys	Gly	Ala	Thr	Thr	Cys	Asp	Thr	Leu	320	325	330
Gly	Met	Ala	Asp	Val	Gly	Thr	Met	Cys	Asp	Pro	Lys	Arg	Ser	Cys	335	340	345
Ser	Val	Ile	Glu	Asp	Asp	Gly	Leu	Pro	Ser	Ala	Phe	Thr	Thr	Ala	350	355	360
His	Glu	Leu	Gly	His	Val	Phe	Asn	Met	Pro	His	Asp	Asn	Val	Lys	365	370	375
Val	Cys	Glu	Glu	Val	Phe	Gly	Lys	Leu	Arg	Ala	Asn	His	Met	Met	380	385	390
Ser	Pro	Thr	Leu	Ile	Gln	Ile	Asp	Arg	Ala	Asn	Pro	Trp	Ser	Ala	395	400	405
Cys	Ser	Ala	Ala	Ile	Ile	Thr	Asp	Phe	Leu	Asp	Ser	Gly	His	Gly	410	415	420
Asp	Cys	Leu	Leu	Asp	Gln	Pro	Ser	Lys	Pro	Ile	Ser	Leu	Pro	Glu	425	430	435
Asp	Leu	Pro	Gly	Ala	Ser	Tyr	Thr	Leu	Ser	Gln	Gln	Cys	Glu	Leu	440	445	450
Ala	Phe	Gly	Val	Gly	Ser	Lys	Pro	Cys	Pro	Tyr	Met	Gln	Tyr	Cys	455	460	465
Thr	Lys	Leu	Trp	Cys	Thr	Gly	Lys	Ala	Lys	Gly	Gln	Met	Val	Cys	470	475	480
Gln	Thr	Arg	His	Phe	Pro	Trp	Ala	Asp	Gly	Thr	Ser	Cys	Gly	Glu	485	490	495
Gly	Lys	Leu	Cys	Leu	Lys	Gly	Ala	Cys	Val	Glu	Arg	His	Asn	Leu	500	505	510
Asn	Lys	His	Arg	Val	Asp	Gly	Ser	Trp	Ala	Lys	Trp	Asp	Pro	Tyr			

Gly Pro Cys Ser	515	Arg Thr Cys Gly Gly	520	Gly Val Gln Leu Ala	525
Arg Gln Cys Thr	530	Asn Pro Thr Pro Ala	535	Asn Gly Gly Lys Tyr	540
Glu Gly Val Arg	545	Val Lys Tyr Arg Ser	550	Cys Asn Leu Glu Pro	555
Pro Ser Ser Ala	560	Ser Gly Lys Ser Phe	565	Arg Glu Glu Gln Cys	570
Ala Phe Asn Gly	575	Tyr Asn His Ser Thr	580	Asn Arg Leu Thr Leu	585
Val Ala Trp Val	590	Pro Lys Tyr Ser Gly	595	Val Ser Pro Arg Asp	600
Cys Lys Leu Ile	605	Cys Arg Ala Asn Gly	610	Thr Gly Tyr Phe Tyr	615
Leu Ala Pro Lys	620	Val Val Val Asp Gly	625	Thr Leu Cys Ser Pro	630
Ser Thr Ser Val	635	Cys Val Gln Gly Lys	640	Cys Ile Lys Ala Gly	645
Asp Gly Asn Leu	650	Gly Ser Lys Lys Arg	655	Phe Asp Lys Cys Gly	660
Cys Gly Gly Asp	665	Asn Lys Ser Cys Lys	670	Lys Val Thr Gly Leu	675
Thr Lys Pro Met	680	His Gly Tyr Asn Phe	685	Val Val Ala Ile Pro	690
Gly Ala Ser Ser	695	Ile Asp Ile Arg Gln	700	Arg Gly Tyr Lys Gly	705
Ile Gly Asp Asp	710	Asn Tyr Leu Ala Leu	715	Lys Asn Ser Gln Gly	720
Tyr Leu Leu Asn	725	Gly His Phe Val Val	730	Ser Ala Val Glu Arg	735
Leu Val Val Lys	740	Gly Ser Leu Leu Arg	745	Tyr Ser Gly Thr Gly	750
Ala Val Glu Ser	755	Leu Gln Ala Ser Arg	760	Pro Ile Leu Glu Pro	765
Thr Val Glu Val	770	Leu Ser Val Gly Lys	775	Met Thr Pro Pro Arg	780
Arg Tyr Ser Phe	785	Tyr Leu Pro Lys Glu	790	Pro Arg Glu Asp Lys	795
Ser His Pro Pro	800	His Pro Arg Gly Gly	805	Pro Ser Val Leu His	810
Ser Val Leu Ser	815	Leu Ser Asn Gln Val	820	Gln Pro Asp Asp	825
Pro Pro Ala Arg	830	Val Ala Gly Ser	835	Gly Pro Cys Ser	840
Ser Cys Gly Ser	845	Gly Leu Gln Lys Arg	850	Ala Val Asp Trp Arg	855
Ser Ala Gly Gln	860	Arg Thr Val Pro Ala	865	Cys Asp Ala Ala His	870
Pro Val Glu Thr	875	Gln Ala Cys Gly Glu	880	Pro Cys Pro Thr Trp	885
Leu Ser Ala Trp	890	Ser Pro Cys Ser Lys	895	Ser Cys Gly Arg Gly	900
Gln Arg Arg Ser	905	Leu Lys Cys Val Gly	910	His Gly Gly Arg Leu	915
Ala Arg Asp Gln	920	Cys Asn Leu His Arg	925	Lys Pro Gln Glu Leu	930
Phe Cys Val Leu	935	Arg Pro Cys	940		945
	950				

<210> 12
 <211> 898
 <212> PRT
 <213> Homo sapiens

<220>
 <221> misc_feature

<223> Incyte ID No: 7604035CD1

<400> 12

Met	Glu	Asn	Trp	Thr	Gly	Arg	Pro	Trp	Leu	Tyr	Leu	Leu	Leu	Leu	
1				5					10						15
Leu	Ser	Leu	Pro	Gln	Leu	Cys	Leu	Asp	Gln	Glu	Val	Leu	Ser	Gly	
				20					25						30
His	Ser	Leu	Gln	Thr	Pro	Thr	Glu	Glu	Gly	Gln	Gly	Pro	Glu	Gly	
				35					40						45
Val	Trp	Gly	Pro	Trp	Val	Gln	Trp	Ala	Ser	Cys	Ser	Gln	Pro	Cys	
				50					55						60
Gly	Val	Gly	Val	Gln	Arg	Arg	Ser	Arg	Thr	Cys	Gln	Leu	Pro	Thr	
				65					70						75
Val	Gln	Leu	His	Pro	Ser	Leu	Pro	Leu	Pro	Pro	Arg	Pro	Pro	Arg	
				80					85						90
His	Pro	Glu	Ala	Leu	Leu	Pro	Arg	Gly	Gln	Gly	Pro	Arg	Pro	Gln	
				95					100						105
Thr	Ser	Pro	Glu	Thr	Leu	Pro	Leu	Tyr	Arg	Thr	Gln	Ser	Arg	Gly	
				110					115						120
Arg	Gly	Gly	Pro	Leu	Arg	Gly	Pro	Ala	Ser	His	Leu	Gly	Arg	Glu	
				125					130						135
Glu	Thr	Gln	Glu	Ile	Arg	Ala	Ala	Arg	Arg	Ser	Arg	Leu	Arg	Asp	
				140					145						150
Pro	Ile	Lys	Pro	Gly	Met	Phe	Gly	Tyr	Gly	Arg	Val	Pro	Phe	Ala	
				155					160						165
Leu	Pro	Leu	His	Arg	Asn	Arg	Arg	His	Pro	Arg	Ser	Pro	Pro	Arg	
				170					175						180
Ser	Glu	Leu	Ser	Leu	Ile	Ser	Ser	Arg	Gly	Glu	Glu	Pro	Ile	Pro	
				185					190						195
Ser	Pro	Thr	Pro	Arg	Ala	Glu	Pro	Phe	Ser	Ala	Asn	Gly	Ser	Pro	
				200					205						210
Gln	Thr	Glu	Leu	Pro	Pro	Thr	Glu	Leu	Ser	Val	His	Thr	Pro	Ser	
				215					220						225
Pro	Gln	Ala	Glu	Pro	Leu	Ser	Pro	Glu	Thr	Ala	Gln	Thr	Glu	Val	
				230					235						240
Ala	Pro	Arg	Thr	Arg	Pro	Ala	Pro	Leu	Arg	His	His	Pro	Arg	Ala	
				245					250						255
Gln	Ala	Ser	Gly	Thr	Glu	Pro	Pro	Ser	Pro	Thr	His	Ser	Leu	Gly	
				260					265						270
Glu	Gly	Gly	Phe	Phe	Arg	Ala	Ser	Pro	Gln	Pro	Arg	Arg	Pro	Ser	
				275					280						285
Ser	Gln	Gly	Trp	Ala	Ser	Pro	Gln	Val	Ala	Gly	Arg	Arg	Pro	Asp	
				290					295						300
Pro	Phe	Pro	Ser	Val	Pro	Arg	Gly	Arg	Gly	Gln	Gln	Gly	Gln	Gly	
				305					310						315
Pro	Trp	Gly	Thr	Gly	Gly	Thr	Pro	His	Gly	Pro	Arg	Leu	Glu	Pro	
				320					325						330
Asp	Pro	Gln	His	Pro	Gly	Ala	Trp	Leu	Pro	Leu	Leu	Ser	Asn	Gly	
				335					340						345
Pro	His	Ala	Ser	Ser	Leu	Trp	Ser	Leu	Phe	Ala	Pro	Ser	Ser	Pro	
				350					355						360
Ile	Pro	Arg	Cys	Ser	Gly	Glu	Ser	Glu	Gln	Leu	Arg	Ala	Cys	Ser	
				365					370						375
Gln	Ala	Pro	Cys	Pro	Pro	Glu	Gln	Pro	Asp	Pro	Arg	Ala	Leu	Gln	
				380					385						390
Cys	Ala	Ala	Phe	Asn	Ser	Gln	Glu	Phe	Met	Gly	Gln	Leu	Tyr	Gln	
				395					400						405
Trp	Glu	Pro	Phe	Thr	Glu	Val	Gln	Gly	Ser	Gln	Arg	Cys	Glu	Leu	
				410					415						420
Asn	Cys	Arg	Pro	Arg	Gly	Phe	Arg	Phe	Tyr	Val	Arg	His	Thr	Glu	
				425					430						435
Lys	Val	Gln	Asp	Gly	Thr	Leu	Cys	Gln	Pro	Gly	Ala	Pro	Asp	Ile	
				440					445						450
Cys	Val	Ala	Gly	Arg	Cys	Leu	Ser	Pro	Gly	Cys	Asp	Gly	Ile	Leu	
				455					460						465
Gly	Ser	Gly	Arg	Arg	Pro	Asp	Gly	Cys	Gly	Val	Cys	Gly	Gly	Asp	
				470					475						480

Asp	Ser	Thr	Cys	Arg	Leu	Val	Ser	Gly	Asn	Leu	Thr	Asp	Arg	Gly
				485					490					495
Gly	Pro	Leu	Gly	Tyr	Gln	Lys	Ile	Leu	Trp	Ile	Pro	Ala	Gly	Ala
				500					505					510
Leu	Arg	Leu	Gln	Ile	Ala	Gln	Leu	Arg	Pro	Ser	Ser	Asn	Tyr	Leu
				515					520					525
Ala	Leu	Arg	Gly	Pro	Gly	Gly	Arg	Ser	Ile	Ile	Asn	Gly	Asn	Trp
				530					535					540
Ala	Val	Asp	Pro	Pro	Gly	Ser	Tyr	Arg	Ala	Gly	Gly	Thr	Val	Phe
				545					550					555
Arg	Tyr	Asn	Arg	Pro	Pro	Arg	Glu	Glu	Gly	Lys	Gly	Glu	Ser	Leu
				560					565					570
Ser	Ala	Glu	Gly	Pro	Thr	Thr	Gln	Pro	Val	Asp	Val	Tyr	Met	Ile
				575					580					585
Phe	Gln	Glu	Glu	Asn	Pro	Gly	Val	Phe	Tyr	Gln	Tyr	Val	Ile	Ser
				590					595					600
Ser	Pro	Pro	Pro	Ile	Leu	Glu	Asn	Pro	Thr	Pro	Glu	Pro	Pro	Val
				605					610					615
Pro	Gln	Leu	Gln	Pro	Glu	Ile	Leu	Arg	Val	Glu	Pro	Pro	Leu	Ala
				620					625					630
Pro	Ala	Pro	Arg	Pro	Ala	Arg	Thr	Pro	Gly	Thr	Leu	Gln	Arg	Gln
				635					640					645
Val	Arg	Ile	Pro	Gln	Met	Pro	Ala	Pro	Pro	His	Pro	Arg	Thr	Pro
				650					655					660
Leu	Gly	Ser	Pro	Ala	Ala	Tyr	Trp	Lys	Arg	Val	Gly	His	Ser	Ala
				665					670					675
Cys	Ser	Ala	Ser	Cys	Gly	Lys	Gly	Val	Trp	Arg	Pro	Ile	Phe	Leu
				680					685					690
Cys	Ile	Ser	Arg	Glu	Ser	Gly	Glu	Glu	Leu	Asp	Glu	Arg	Ser	Cys
				695					700					705
Ala	Ala	Gly	Ala	Arg	Pro	Pro	Ala	Ser	Pro	Glu	Pro	Cys	His	Gly
				710					715					720
Thr	Pro	Cys	Pro	Pro	Tyr	Trp	Glu	Ala	Gly	Glu	Trp	Thr	Ser	Cys
				725					730					735
Ser	Arg	Ser	Cys	Gly	Pro	Gly	Thr	Gln	His	Arg	Gln	Leu	Gln	Cys
				740					745					750
Arg	Gln	Glu	Phe	Gly	Gly	Gly	Gly	Ser	Ser	Val	Pro	Pro	Glu	Arg
				755					760					765
Cys	Gly	His	Leu	Pro	Arg	Pro	Asn	Ile	Thr	Gln	Ser	Cys	Gln	Leu
				770					775					780
Arg	Leu	Cys	Gly	His	Trp	Glu	Val	Gly	Ser	Pro	Trp	Ser	Gln	Cys
				785					790					795
Ser	Val	Arg	Cys	Gly	Arg	Gly	Gln	Arg	Ser	Arg	Gln	Val	Arg	Cys
				800					805					810
Val	Gly	Asn	Asn	Gly	Asp	Glu	Val	Ser	Glu	Gln	Glu	Cys	Ala	Ser
				815					820					825
Gly	Pro	Pro	Gln	Pro	Pro	Ser	Arg	Glu	Ala	Cys	Asp	Met	Gly	Pro
				830					835					840
Cys	Thr	Thr	Ala	Trp	Phe	His	Ser	Asp	Trp	Ser	Ser	Lys	Cys	Ser
				845					850					855
Ala	Glu	Cys	Gly	Thr	Gly	Ile	Gln	Arg	Arg	Ser	Val	Val	Cys	Leu
				860					865					870
Gly	Ser	Gly	Ala	Ala	Thr	Arg	Ala	Arg	Pro	Gly	Gly	Ser	Arg	Ser
				875					880					885
Arg	Asn	Trp	Ala	Glu	Leu	Ser	Asn	Arg	Lys	Pro	Ala	Pro		
				890					895					

<210> 13

<211> 631

<212> PRT

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 3473847CD1

<400> 13

Met	Phe	Leu	Leu	Ala	Trp	Gly	Gln	Asp	Pro	Trp	Arg	Leu	Pro	Gly
1				5					10					15
Thr	Tyr	Val	Val	Val	Leu	Lys	Glu	Glu	Thr	His	Leu	Ser	Gln	Ser
				20					25					30
Glu	Arg	Thr	Ala	Arg	Arg	Leu	Gln	Ala	Gln	Ala	Ala	Arg	Arg	Gly
				35					40					45
Tyr	Leu	Thr	Lys	Ile	Leu	His	Val	Phe	His	Gly	Leu	Leu	Pro	Gly
				50					55					60
Phe	Leu	Val	Lys	Met	Ser	Gly	Asp	Leu	Leu	Glu	Leu	Ala	Leu	Lys
				65					70					75
Leu	Pro	His	Val	Asp	Tyr	Ile	Glu	Glu	Asp	Ser	Ser	Val	Phe	Ala
				80					85					90
Gln	Ser	Ile	Pro	Trp	Asn	Leu	Glu	Arg	Ile	Thr	Pro	Pro	Arg	Tyr
				95					100					105
Arg	Ala	Asp	Glu	Tyr	Gln	Pro	Pro	Asp	Gly	Gly	Ser	Leu	Val	Glu
				110					115					120
Val	Tyr	Leu	Leu	Asp	Thr	Ser	Ile	Gln	Ser	Asp	His	Arg	Glu	Ile
				125					130					135
Glu	Gly	Arg	Val	Met	Val	Thr	Asp	Phe	Glu	Asn	Val	Pro	Glu	Glu
				140					145					150
Asp	Gly	Thr	Arg	Phe	His	Arg	Gln	Ala	Ser	Lys	Cys	Asp	Ser	His
				155					160					165
Gly	Thr	His	Leu	Ala	Gly	Val	Val	Ser	Gly	Arg	Asp	Ala	Gly	Val
				170					175					180
Ala	Lys	Gly	Ala	Ser	Met	Arg	Ser	Leu	Arg	Val	Leu	Asn	Cys	Gln
				185					190					195
Gly	Lys	Gly	Thr	Val	Ser	Gly	Thr	Leu	Ile	Gly	Leu	Glu	Phe	Ile
				200					205					210
Arg	Lys	Ser	Gln	Leu	Val	Gln	Pro	Val	Gly	Pro	Leu	Val	Val	Leu
				215					220					225
Leu	Pro	Leu	Ala	Gly	Gly	Tyr	Ser	Arg	Val	Leu	Asn	Ala	Ala	Cys
				230					235					240
Gln	Arg	Leu	Ala	Arg	Ala	Gly	Val	Val	Leu	Val	Thr	Ala	Ala	Gly
				245					250					255
Asn	Phe	Arg	Asp	Asp	Ala	Cys	Leu	Tyr	Ser	Pro	Ala	Ser	Ala	Pro
				260					265					270
Glu	Val	Ile	Thr	Val	Gly	Ala	Thr	Asn	Ala	Gln	Asp	Gln	Pro	Val
				275					280					285
Thr	Leu	Gly	Thr	Leu	Gly	Thr	Asn	Phe	Gly	Arg	Cys	Val	Asp	Leu
				290					295					300
Phe	Ala	Pro	Gly	Glu	Asp	Ile	Ile	Gly	Ala	Ser	Ser	Asp	Cys	Ser
				305					310					315
Thr	Cys	Phe	Val	Ser	Gln	Ser	Gly	Thr	Ser	Gln	Ala	Ala	Ala	His
				320					325					330
Val	Ala	Gly	Ile	Ala	Ala	Met	Met	Leu	Ser	Ala	Glu	Pro	Glu	Leu
				335					340					345
Thr	Leu	Ala	Glu	Leu	Arg	Gln	Arg	Leu	Ile	His	Phe	Ser	Ala	Lys
				350					355					360
Asp	Val	Ile	Asn	Glu	Ala	Trp	Phe	Pro	Glu	Asp	Gln	Arg	Val	Leu
				365					370					375
Thr	Pro	Asn	Leu	Val	Ala	Ala	Leu	Pro	Pro	Ser	Thr	His	Gly	Ala
				380					385					390
Gly	Trp	Gln	Leu	Phe	Cys	Arg	Thr	Val	Trp	Ser	Ala	His	Ser	Gly
				395					400					405
Pro	Thr	Arg	Met	Ala	Thr	Ala	Ile	Ala	Arg	Cys	Ala	Pro	Asp	Glu
				410					415					420
Glu	Leu	Leu	Ser	Cys	Ser	Ser	Phe	Ser	Arg	Ser	Gly	Lys	Arg	Arg
				425					430					435
Gly	Glu	Arg	Met	Glu	Ala	Gln	Gly	Gly	Lys	Leu	Val	Cys	Arg	Ala
				440					445					450
His	Asn	Ala	Phe	Gly	Gly	Glu	Gly	Val	Tyr	Ala	Ile	Ala	Arg	Cys
				455					460					465
Cys	Leu	Leu	Pro	Gln	Ala	Asn	Cys	Ser	Val	His	Thr	Ala	Pro	Pro
				470					475					480
Ala	Glu	Ala	Ser	Met	Gly	Thr	Arg	Val	His	Cys	His	Gln	Gln	Gly
				485					490					495
His	Val	Leu	Thr	Gly	Cys	Ser	Ser	His	Trp	Glu	Val	Glu	Asp	Leu

	500		505		510
Gly Thr His Lys	Pro Val Leu Arg	Pro Arg Gly Gln Pro	Asn		
	515		520		525
Gln Cys Val Gly	His Arg Glu Ala Ser	Ile His Ala Ser Cys	Cys		
	530		535		540
His Ala Pro Gly	Leu Glu Cys Lys Val	Lys Glu His Gly Ile	Pro		
	545		550		555
Ala Pro Gln Glu	Gln Val Thr Val Ala	Cys Glu Glu Gly Trp	Thr		
	560		565		570
Leu Thr Gly Cys	Ser Ala Leu Pro Gly	Thr Ser His Val Leu	Gly		
	575		580		585
Ala Tyr Ala Val	Asp Asn Thr Cys Val	Val Arg Ser Arg Asp	Val		
	590		595		600
Ser Thr Thr Gly	Ser Thr Ser Glu Glu	Ala Val Thr Ala Val	Ala		
	605		610		615
Ile Cys Cys Arg	Ser Arg His Leu Ala	Gln Ala Ser Gln Glu	Leu		
	620		625		630
Gln					

<210> 14

<211> 470

<212> PRT

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 3750004CD1

<400> 14

Met Arg His Arg Thr	Asp Leu Gly Gln Asn	Leu Leu Leu Phe	Leu
1	5	10	15
Trp Ala Leu Leu Asn	Cys Gly Leu Gly Val	Ser Ala Gln Gly	Pro
	20	25	30
Gly Glu Trp Thr Pro	Trp Val Ser Trp Thr	Arg Cys Ser Ser	Ser
	35	40	45
Cys Gly Arg Gly Val	Ser Val Arg Ser Arg	Arg Cys Leu Arg	Leu
	50	55	60
Pro Gly Glu Glu Pro	Cys Trp Gly Asp Ser	His Glu Tyr Arg	Leu
	65	70	75
Cys Gln Leu Pro Asp	Cys Pro Pro Gly Ala	Val Pro Phe Arg	Asp
	80	85	90
Leu Gln Cys Ala Leu	Tyr Asn Gly Arg Pro	Val Leu Gly Thr	Gln
	95	100	105
Lys Thr Tyr Gln Trp	Val Pro Phe His Gly	Ala Pro Asn Gln	Cys
	110	115	120
Asp Leu Asn Cys Leu	Ala Glu Gly His Ala	Phe Tyr His Ser	Phe
	125	130	135
Gly Arg Val Leu Asp	Gly Thr Ala Cys Ser	Pro Gly Ala Gln	Gly
	140	145	150
Val Cys Val Ala Gly	Arg Cys Leu Ser Ala	Gly Cys Asp Gly	Leu
	155	160	165
Leu Gly Ser Gly Ala	Leu Glu Asp Arg Cys	Gly Arg Cys Gly	Gly
	170	175	180
Ala Asn Asp Ser Cys	Leu Phe Val Gln Arg	Val Phe Arg Asp	Ala
	185	190	195
Gly Ala Phe Ala Gly	Tyr Trp Asn Val Thr	Leu Ile Pro Glu	Gly
	200	205	210
Ala Arg His Ile Arg	Val Glu His Arg Ser	Arg Asn His Leu	Gly
	215	220	225
Ile Leu Gly Ser Leu	Met Gly Gly Asp Gly	Arg Tyr Val Leu	Asn
	230	235	240
Gly His Trp Val Val	Ser Pro Pro Gly Thr	Tyr Glu Ala Ala	Gly
	245	250	255
Thr His Val Val Tyr	Thr Arg Asp Thr Gly	Pro Gln Glu Thr	Leu
	260	265	270
Gln Ala Ala Gly Pro	Thr Ser His Asp Leu	Leu Leu Gln Val	Leu

275	280	285
Leu Gln Glu Pro Asn Pro Gly Ile Glu Phe Glu Phe Trp Leu Pro		
290	295	300
Arg Glu Arg Tyr Ser Pro Phe Gln Ala Arg Val Gln Ala Leu Gly		
305	310	315
Trp Pro Leu Arg Gln Pro Gln Pro Arg Gly Val Glu Pro Gln Pro		
320	325	330
Pro Ala Ala Pro Ala Val Thr Pro Ala Gln Thr Pro Thr Leu Ala		
335	340	345
Pro Asp Pro Cys Pro Pro Cys Pro Asp Thr Arg Gly Arg Ala His		
350	355	360
Arg Leu Leu His Tyr Cys Gly Ser Asp Phe Val Phe Gln Ala Arg		
365	370	375
Val Leu Gly His His His Gln Ala Gln Glu Thr Arg Tyr Glu Val		
380	385	390
Arg Ile Gln Leu Val Tyr Lys Asn Arg Ser Pro Leu Arg Ala Arg		
395	400	405
Glu Tyr Val Trp Ala Pro Gly His Cys Pro Cys Pro Met Leu Ala		
410	415	420
Pro His Arg Asp Tyr Leu Met Ala Val Gln Arg Leu Val Ser Pro		
425	430	435
Asp Gly Thr Gln Asp Gln Leu Leu Leu Pro His Ala Gly Tyr Ala		
440	445	450
Arg Pro Trp Ser Pro Ala Glu Asp Ser Arg Ile Arg Leu Thr Ala		
455	460	465
Arg Arg Cys Pro Gly		
470		

<210> 15
 <211> 110
 <212> PRT
 <213> Homo sapiens

<220>
 <221> misc_feature
 <223> Incyte ID No: 4904126CD1

<400> 15	
Met Ala Asp Lys Val Leu Lys Glu Lys Arg Lys Gln Phe Ile Arg	
1 5 10 15	
Ser Val Gly Glu Gly Thr Ile Asn Gly Leu Leu Gly Glu Leu Leu	
20 25 30	
Glu Thr Arg Val Leu Ser Gln Glu Glu Ile Glu Ile Val Lys Cys	
35 40 45	
Glu Asn Ala Thr Val Met Asp Lys Ala Arg Ala Leu Leu Asp Ser	
50 55 60	
Val Ile Arg Lys Gly Ala Pro Ala Cys Gln Ile Cys Ile Thr Tyr	
65 70 75	
Ile Cys Glu Glu Asp Ser His Leu Ala Gly Thr Leu Gly Leu Ser	
80 85 90	
Ala Gly Pro Thr Ser Gly Asn His Leu Thr Thr Gln Asp Ser Gln	
95 100 105	
Ile Val Leu Pro Ser	
110	

<210> 16
 <211> 879
 <212> PRT
 <213> Homo sapiens

<220>
 <221> misc_feature
 <223> Incyte ID No: 71268415CD1

<400> 16	
Met Ser Leu Phe Ile Phe Cys Arg Gln Leu Phe Ala Pro Ser Tyr	
1 5 10 15	

Thr	Glu	Thr	His	Tyr	Thr	Ser	Ser	Gly	Asn	Pro	Gln	Thr	Thr	Thr
				20					25					30
Arg	Lys	Leu	Glu	Asp	His	Cys	Phe	Tyr	His	Gly	Thr	Val	Arg	Glu
				35					40					45
Thr	Glu	Leu	Ser	Ser	Val	Thr	Leu	Ser	Thr	Cys	Arg	Gly	Ile	Arg
				50					55					60
Gly	Leu	Ile	Thr	Val	Ser	Ser	Asn	Leu	Ser	Tyr	Val	Ile	Glu	Pro
				65					70					75
Leu	Pro	Asp	Ser	Lys	Gly	Gln	His	Leu	Ile	Tyr	Arg	Ser	Glu	His
				80					85					90
Leu	Lys	Pro	Pro	Pro	Gly	Asn	Cys	Gly	Phe	Glu	His	Ser	Lys	Pro
				95					100					105
Thr	Thr	Arg	Asp	Trp	Ala	Leu	Gln	Phe	Thr	Gln	Gln	Thr	Lys	Lys
				110					115					120
Arg	Pro	Arg	Arg	Met	Lys	Arg	Glu	Asp	Leu	Asn	Ser	Met	Lys	Tyr
				125					130					135
Val	Glu	Leu	Tyr	Leu	Val	Ala	Asp	Tyr	Leu	Glu	Phe	Gln	Lys	Asn
				140					145					150
Arg	Arg	Asp	Gln	Asp	Ala	Thr	Lys	His	Lys	Leu	Ile	Glu	Ile	Ala
				155					160					165
Asn	Tyr	Val	Asp	Lys	Phe	Tyr	Arg	Ser	Leu	Asn	Ile	Arg	Ile	Ala
				170					175					180
Leu	Val	Gly	Leu	Glu	Val	Trp	Thr	His	Gly	Asn	Met	Cys	Glu	Val
				185					190					195
Ser	Glu	Asn	Pro	Tyr	Ser	Thr	Leu	Trp	Ser	Phe	Leu	Ser	Trp	Arg
				200					205					210
Arg	Lys	Leu	Leu	Ala	Gln	Lys	Tyr	His	Asp	Asn	Ala	Gln	Leu	Ile
				215					220					225
Thr	Gly	Met	Ser	Phe	His	Gly	Thr	Thr	Ile	Gly	Leu	Ala	Pro	Leu
				230					235					240
Met	Ala	Met	Cys	Ser	Val	Tyr	Gln	Ser	Gly	Gly	Val	Asn	Met	Asp
				245					250					255
His	Ser	Glu	Asn	Ala	Ile	Gly	Val	Ala	Ala	Thr	Met	Ala	His	Glu
				260					265					270
Met	Gly	His	Asn	Phe	Gly	Met	Thr	His	Asp	Ser	Ala	Asp	Cys	Cys
				275					280					285
Ser	Ala	Ser	Ala	Ala	Asp	Gly	Gly	Cys	Ile	Met	Ala	Ala	Ala	Thr
				290					295					300
Gly	His	Pro	Phe	Pro	Lys	Val	Phe	Asn	Gly	Cys	Asn	Arg	Arg	Glu
				305					310					315
Leu	Asp	Arg	Tyr	Leu	Gln	Ser	Gly	Gly	Gly	Met	Cys	Leu	Ser	Asn
				320					325					330
Met	Pro	Asp	Thr	Arg	Met	Leu	Tyr	Gly	Gly	Arg	Arg	Cys	Gly	Asn
				335					340					345
Gly	Tyr	Leu	Glu	Asp	Gly	Glu	Glu	Cys	Asp	Cys	Gly	Glu	Glu	Glu
				350					355					360
Glu	Cys	Asn	Asn	Pro	Cys	Cys	Asn	Ala	Ser	Asn	Cys	Thr	Leu	Arg
				365					370					375
Pro	Gly	Ala	Glu	Cys	Ala	His	Gly	Ser	Cys	Cys	His	Gln	Cys	Lys
				380					385					390
Leu	Leu	Ala	Pro	Gly	Thr	Leu	Cys	Arg	Glu	Gln	Ala	Arg	Gln	Cys
				395					400					405
Asp	Leu	Pro	Glu	Phe	Cys	Thr	Gly	Lys	Ser	Pro	His	Cys	Pro	Thr
				410					415					420
Asn	Phe	Tyr	Gln	Met	Asp	Gly	Thr	Pro	Cys	Glu	Gly	Gly	Gln	Ala
				425					430					435
Tyr	Cys	Tyr	Asn	Gly	Met	Cys	Leu	Thr	Tyr	Gln	Glu	Gln	Cys	Gln
				440					445					450
Gln	Leu	Trp	Gly	Pro	Gly	Ala	Arg	Pro	Ala	Pro	Asp	Leu	Cys	Phe
				455					460					465
Glu	Lys	Val	Asn	Val	Ala	Gly	Asp	Thr	Phe	Gly	Asn	Cys	Gly	Lys
				470					475					480
Asp	Met	Asn	Gly	Glu	His	Arg	Lys	Cys	Asn	Met	Arg	Asp	Ala	Lys
				485					490					495
Cys	Gly	Lys	Ile	Gln	Cys	Gln	Ser	Ser	Glu	Ala	Arg	Pro	Leu	Glu
				500					505					510
Ser	Asn	Ala	Val	Pro	Ile	Asp	Thr	Thr	Ile	Ile	Met	Asn	Gly	Arg

Gln Ile Gln Cys	515	520	525
Arg Gly Thr His Val	530	Tyr Arg Gly Pro Glu	Glu
Glu Gly Asp Met	545	Val Met Thr Gly Thr	Lys
Cys Gly Tyr Asn	560	Gly Gln Cys Arg Asn	Thr
Ser Phe Phe Glu	575	Lys Lys Cys Asn Gly	His
Gly Val Cys Asn	590	His Cys Leu Pro Gly	Trp
Ala Pro Pro Phe	605	Gly Gly Ser Ile	Asp
Ser Gly Pro Met	620	Gly Pro Val Val Ala	Gly
Val Leu Val Ala	635	Val Leu Met Leu Met	Tyr
Tyr Cys Cys Arg	650	Gly Gln Leu Lys Pro	Ser
Ala Leu Pro Ser	665	Phe Ser Cys Pro Phe	Arg
Val Ser Gln Asn	680	Ala Asn Pro Thr Phe	Lys
Leu Gln Thr Pro	695	Val Ile Asn Thr Pro	Glu
Ile Leu Arg Lys	710	Pro Arg Pro Pro Pro	Asp
Tyr Leu Arg Gly	725	Pro Leu Pro Ala His	Leu
Ser Arg Ala Ala	740	Pro Gly Ser Gln Ile	Glu
Arg Thr Glu Ser	755	Pro Ser Arg Pro Ile	Pro
Pro Ala Pro Asn	770	Asp Phe Ser Arg Pro	Arg
Pro Pro Gln Lys	785	Pro Val Pro Gly Arg	Arg
Ser Leu Pro Arg	800	Pro Leu Arg Pro Pro	Gly
Ala Gly Pro Gln	815	Ala Ala Leu Ala Pro	Lys
Val Ser Pro Arg	830	Lys Ala Gly Thr Arg	Gly
Leu Gln Gly Gly	845	Lys Thr Lys Gln Phe	Met
Leu Leu Val Val	860	Glu Gln Lys Pro Arg	Ala
Lys His Ser Cys	875		

<210> 17

<211> 850

<212> PRT

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 7473301CD1

<400> 17

Met Asp Lys Glu Asn	Ser Asp Val Ser	Ala Ala Pro Ala Asp	Leu
1	5	10	15
Lys Ile Ser Asn Ile	Ser Val Gln Val	Ser Ala Gln Lys	Lys
	20	25	30
Leu Pro Val Arg Arg	Pro Pro Leu Pro	Gly Arg Arg Leu Pro	Leu
	35	40	45
Pro Gly Arg Arg Pro	Pro Gln Arg Pro	Ile Gly Lys Ala Lys	Pro

				50					55					60
Lys	Lys	Gln	Ser	Lys	Lys	Lys	Val	Pro	Phe	Trp	Asn	Val	Gln	Asn
				65					70					75
Lys	Ile	Ile	Leu	Phe	Thr	Val	Phe	Leu	Phe	Ile	Leu	Ala	Val	Ile
				80					85					90
Ala	Trp	Thr	Leu	Leu	Trp	Leu	Tyr	Ile	Ser	Lys	Thr	Glu	Ser	Lys
				95					100					105
Asp	Ala	Phe	Tyr	Phe	Ala	Gly	Met	Phe	Arg	Ile	Thr	Asn	Ile	Glu
				110					115					120
Phe	Leu	Pro	Glu	Tyr	Arg	Gln	Lys	Glu	Ser	Arg	Glu	Phe	Leu	Ser
				125					130					135
Val	Ser	Arg	Thr	Val	Gln	Gln	Val	Ile	Asn	Leu	Val	Tyr	Thr	Thr
				140					145					150
Ser	Ala	Phe	Ser	Lys	Phe	Tyr	Glu	Gln	Ser	Val	Val	Ala	Asp	Val
				155					160					165
Ser	Ser	Asn	Asn	Lys	Gly	Gly	Leu	Leu	Val	His	Phe	Trp	Ile	Val
				170					175					180
Phe	Val	Met	Pro	Arg	Ala	Lys	Gly	His	Ile	Phe	Cys	Glu	Asp	Cys
				185					190					195
Val	Ala	Ala	Ile	Leu	Lys	Asp	Ser	Ile	Gln	Thr	Ser	Ile	Ile	Asn
				200					205					210
Arg	Thr	Ser	Val	Gly	Ser	Leu	Gln	Gly	Leu	Ala	Val	Asp	Met	Asp
				215					220					225
Ser	Val	Val	Leu	Asn	Gly	Asp	Cys	Trp	Ser	Phe	Leu	Lys	Lys	Lys
				230					235					240
Lys	Arg	Lys	Glu	Asn	Gly	Ala	Val	Ser	Thr	Asp	Lys	Gly	Cys	Ser
				245					250					255
Gln	Tyr	Phe	Tyr	Ala	Glu	His	Leu	Ser	Leu	His	Tyr	Pro	Leu	Glu
				260					265					270
Ile	Ser	Ala	Ala	Ser	Gly	Arg	Leu	Met	Cys	His	Phe	Lys	Leu	Val
				275					280					285
Ala	Ile	Val	Gly	Tyr	Leu	Ile	Arg	Leu	Ser	Ile	Lys	Ser	Ile	Gln
				290					295					300
Ile	Glu	Ala	Asp	Asn	Cys	Val	Thr	Asp	Ser	Leu	Thr	Ile	Tyr	Asp
				305					310					315
Ser	Leu	Leu	Pro	Ile	Arg	Ser	Ser	Ile	Leu	Tyr	Arg	Ile	Cys	Glu
				320					325					330
Pro	Thr	Arg	Thr	Leu	Met	Ser	Phe	Val	Ser	Thr	Asn	Asn	Leu	Met
				335					340					345
Leu	Val	Thr	Phe	Lys	Ser	Pro	His	Ile	Arg	Arg	Leu	Ser	Gly	Ile
				350					355					360
Arg	Ala	Tyr	Phe	Glu	Val	Ile	Pro	Glu	Gln	Lys	Cys	Glu	Asn	Thr
				365					370					375
Val	Leu	Val	Lys	Asp	Ile	Thr	Gly	Phe	Glu	Gly	Lys	Ile	Ser	Ser
				380					385					390
Pro	Tyr	Tyr	Pro	Ser	Tyr	Tyr	Pro	Pro	Lys	Cys	Lys	Cys	Thr	Trp
				395					400					405
Lys	Phe	Gln	Thr	Ser	Leu	Ser	Thr	Leu	Gly	Ile	Ala	Leu	Lys	Phe
				410					415					420
Tyr	Asn	Tyr	Ser	Ile	Thr	Lys	Lys	Ser	Met	Lys	Gly	Cys	Glu	His
				425					430					435
Gly	Trp	Trp	Glu	Ile	Tyr	Glu	His	Met	Tyr	Cys	Gly	Ser	Tyr	Met
				440					445					450
Asp	His	Gln	Thr	Ile	Phe	Arg	Val	Pro	Ser	Pro	Leu	Val	His	Ile
				455					460					465
Gln	Leu	Gln	Cys	Ser	Ser	Arg	Leu	Ser	Gly	Lys	Pro	Leu	Leu	Ala
				470					475					480
Glu	Tyr	Gly	Ser	Tyr	Asn	Ile	Ser	Gln	Pro	Cys	Pro	Val	Gly	Ser
				485					490					495
Phe	Arg	Cys	Ser	Ser	Gly	Leu	Cys	Val	Pro	Gln	Ala	Gln	Arg	Gly
				500					505					510
Asp	Gly	Val	Asn	Asp	Cys	Phe	Asp	Glu	Ser	Asp	Glu	Leu	Phe	Cys
				515					520					525
Val	Ser	Pro	Gln	Pro	Ala	Cys	Asn	Thr	Ser	Ser	Phe	Arg	Gln	His
				530					535					540
Gly	Pro	Leu	Ile	Cys	Asp	Gly	Phe	Arg	Asp	Cys	Glu	Asn	Gly	Arg
				545					550					555

Asp	Glu	Gln	Asn	Cys	Thr	Gln	Ser	Ile	Pro	Cys	Asn	Asn	Arg	Thr
				560					565					570
Phe	Lys	Cys	Gly	Asn	Asp	Ile	Cys	Phe	Arg	Lys	Gln	Asn	Ala	Lys
				575					580					585
Cys	Asp	Gly	Thr	Val	Asp	Cys	Pro	Asp	Gly	Ser	Asp	Glu	Glu	Gly
				590					595					600
Cys	Thr	Cys	Ser	Arg	Ser	Ser	Ser	Ala	Leu	His	Arg	Ile	Ile	Gly
				605					610					615
Gly	Thr	Asp	Thr	Leu	Glu	Gly	Gly	Trp	Pro	Trp	Gln	Val	Ser	Leu
				620					625					630
His	Phe	Val	Gly	Ser	Ala	Tyr	Cys	Gly	Ala	Ser	Val	Ile	Ser	Arg
				635					640					645
Glu	Trp	Leu	Leu	Ser	Ala	Ala	His	Cys	Phe	His	Gly	Asn	Arg	Leu
				650					655					660
Ser	Asp	Pro	Thr	Pro	Trp	Thr	Ala	His	Leu	Gly	Met	Tyr	Val	Gln
				665					670					675
Gly	Asn	Ala	Lys	Phe	Val	Ser	Pro	Val	Arg	Arg	Ile	Val	Val	His
				680					685					690
Glu	Tyr	Tyr	Asn	Ser	Gln	Thr	Phe	Asp	Tyr	Asp	Ile	Ala	Leu	Leu
				695					700					705
Gln	Leu	Ser	Ile	Ala	Trp	Pro	Glu	Thr	Leu	Lys	Gln	Leu	Ile	Gln
				710					715					720
Pro	Ile	Cys	Ile	Pro	Pro	Thr	Gly	Gln	Arg	Val	Arg	Ser	Gly	Glu
				725					730					735
Lys	Cys	Trp	Val	Thr	Gly	Trp	Gly	Arg	Arg	His	Glu	Ala	Asp	Asn
				740					745					750
Lys	Gly	Ser	Leu	Val	Leu	Gln	Gln	Ala	Glu	Val	Glu	Leu	Ile	Asp
				755					760					765
Gln	Thr	Leu	Cys	Val	Ser	Thr	Tyr	Gly	Ile	Ile	Thr	Ser	Arg	Met
				770					775					780
Leu	Cys	Ala	Gly	Ile	Met	Ser	Gly	Lys	Arg	Asp	Ala	Cys	Lys	Gly
				785					790					795
Asp	Ser	Gly	Gly	Pro	Leu	Ser	Cys	Arg	Arg	Lys	Ser	Asp	Gly	Lys
				800					805					810
Trp	Ile	Leu	Thr	Gly	Ile	Val	Ser	Trp	Gly	His	Gly	Cys	Gly	Arg
				815					820					825
Pro	Asn	Phe	Pro	Gly	Val	Tyr	Thr	Arg	Val	Ser	Asn	Phe	Val	Pro
				830					835					840
Trp	Ile	His	Lys	Tyr	Val	Pro	Ser	Leu	Leu					
				845					850					

<210> 18

<211> 254

<212> PRT

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 7473308CD1

<400> 18

Met	Gln	Asp	His	Arg	Lys	Gly	Lys	Ala	Ala	Val	Gly	Val	Ser	Phe
1				5					10					15
Asp	Asp	Asp	Asp	Lys	Ile	Val	Gly	Gly	Tyr	Asn	Cys	Glu	Glu	Asn
				20					25					30
Ser	Val	Pro	Tyr	Gln	Val	Ser	Leu	Asn	Ser	Gly	Tyr	His	Phe	Cys
				35					40					45
Val	Gly	Ser	Leu	Asn	Arg	Glu	Tyr	Cys	Ile	Gln	Val	Arg	Leu	Gly
				50					55					60
Glu	His	Asn	Ile	Glu	Val	Leu	Glu	Gly	Asn	Glu	Gln	Phe	Ile	Tyr
				65					70					75
Ala	Val	Lys	Ile	Ile	Arg	His	Pro	Lys	Tyr	Asn	Ser	Trp	Thr	Leu
				80					85					90
Asp	Asn	Asp	Ile	Leu	Leu	Ile	Lys	Leu	Ser	Thr	Pro	Ala	Ile	Ile
				95					100					105
Asn	Ala	His	Val	Ser	Thr	Ile	Ser	Leu	Pro	Thr	Thr	Pro	Pro	Ala
				110					115					120

Ala	Gly	Thr	Glu	Cys	Leu	Ile	Ser	Gly	Trp	Gly	Asn	Thr	Leu	Ser
				125					130					135
Ser	Gly	Ala	Asp	Tyr	Pro	Asp	Glu	Leu	Gln	Cys	Leu	Asp	Ala	Pro
				140					145					150
Val	Leu	Ser	Gln	Ala	Glu	Tyr	Glu	Ala	Ser	Tyr	Pro	Gly	Lys	Ile
				155					160					165
Thr	Asn	Asn	Val	Phe	Cys	Val	Gly	Phe	Leu	Glu	Gly	Gly	Lys	Asp
				170					175					180
Ser	Cys	Gln	Ile	Ile	Pro	Ile	Lys	Val	Gln	Gln	Leu	Val	Thr	Ser
				185					190					195
Ser	Gln	Glu	Thr	Asp	Ile	Arg	Ile	Pro	Met	Ala	Leu	Gln	Thr	Ala
				200					205					210
Ala	Ser	Thr	Ser	Tyr	Leu	Gly	Pro	Leu	Asp	Ser	Leu	His	Arg	Lys
				215					220					225
Val	Ser	His	Pro	Thr	Glu	Lys	Arg	Cys	Gln	Gln	Lys	Gln	Gly	Met
				230					235					240
Lys	Ile	Thr	Asp	Asn	His	Gly	Ile	Thr	Ser	Lys	Trp	Ser	Val	
				245					250					

<210> 19

<211> 568

<212> PRT

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 7478021CD1

<400> 19

Met	Leu	Ala	Ala	Ser	Ile	Phe	Arg	Pro	Thr	Leu	Leu	Leu	Cys	Trp
1				5					10					15
Leu	Ala	Ala	Pro	Trp	Pro	Thr	Gln	Pro	Glu	Ser	Leu	Phe	His	Ser
				20					25					30
Arg	Asp	Arg	Ser	Asp	Leu	Glu	Pro	Ser	Pro	Leu	Arg	Gln	Ala	Lys
				35					40					45
Pro	Ile	Ala	Asp	Leu	His	Ala	Ala	Gln	Arg	Phe	Leu	Ser	Arg	Tyr
				50					55					60
Gly	Trp	Ser	Gly	Val	Trp	Ala	Ala	Trp	Gly	Pro	Ser	Pro	Glu	Gly
				65					70					75
Pro	Pro	Glu	Thr	Pro	Lys	Gly	Ala	Ala	Leu	Ala	Glu	Ala	Val	Arg
				80					85					90
Arg	Phe	Gln	Arg	Ala	Asn	Ala	Leu	Pro	Ala	Ser	Gly	Glu	Leu	Asp
				95					100					105
Ala	Ala	Thr	Leu	Ala	Ala	Met	Asn	Arg	Pro	Arg	Cys	Gly	Val	Pro
				110					115					120
Asp	Met	Arg	Pro	Pro	Pro	Pro	Ser	Ala	Pro	Pro	Ser	Pro	Pro	Gly
				125					130					135
Pro	Pro	Pro	Arg	Ala	Arg	Ser	Arg	Arg	Ser	Pro	Arg	Ala	Pro	Leu
				140					145					150
Ser	Leu	Ser	Arg	Arg	Gly	Trp	Gln	Pro	Arg	Gly	Tyr	Pro	Asp	Gly
				155					160					165
Gly	Ala	Ala	Gln	Ala	Phe	Ser	Lys	Arg	Thr	Leu	Ser	Trp	Arg	Leu
				170					175					180
Leu	Gly	Glu	Ala	Leu	Ser	Ser	Gln	Leu	Ser	Val	Ala	Asp	Gln	Arg
				185					190					195
Arg	Ile	Glu	Ala	Leu	Ala	Phe	Arg	Met	Trp	Ser	Glu	Val	Thr	Pro
				200					205					210
Leu	Asp	Phe	Arg	Glu	Asp	Leu	Ala	Ala	Pro	Gly	Ala	Ala	Val	Asp
				215					220					225
Ile	Lys	Leu	Gly	Phe	Gly	Arg	Arg	His	Leu	Gly	Cys	Pro	Arg	Ala
				230					235					240
Phe	Asp	Gly	Ser	Gly	Gln	Glu	Phe	Ala	His	Ala	Trp	Arg	Leu	Gly
				245					250					255
Asp	Ile	His	Phe	Asp	Asp	Asp	Glu	His	Phe	Thr	Pro	Pro	Thr	Ser
				260					265					270
Asp	Thr	Gly	Ile	Ser	Leu	Leu	Lys	Val	Ala	Val	His	Glu	Ile	Gly
				275					280					285

His Val Leu Gly	Leu Pro His Thr Tyr	Arg Thr Gly Ser Ile Met	
	290	295	300
Gln Pro Asn Tyr	Ile Pro Gln Glu Pro	Ala Phe Glu Leu Asp Trp	
	305	310	315
Ser Asp Arg Lys	Ala Ile Gln Lys Leu	Tyr Gly Ser Cys Glu Gly	
	320	325	330
Ser Phe Asp Thr	Ala Phe Asp Trp Ile	Arg Lys Glu Arg Asn Gln	
	335	340	345
Tyr Gly Glu Val	Met Val Arg Phe Ser	Thr Tyr Phe Phe Arg Asn	
	350	355	360
Ser Trp Tyr Trp	Leu Tyr Glu Asn Arg	Asn Asn Arg Thr Arg Tyr	
	365	370	375
Gly Asp Pro Ile	Gln Ile Leu Thr Gly	Trp Pro Gly Ile Pro Thr	
	380	385	390
His Asn Ile Asp	Ala Phe Val His Ile	Trp Thr Trp Lys Arg Asp	
	395	400	405
Glu Arg Tyr Phe	Phe Gln Gly Asn Gln	Tyr Trp Arg Tyr Asp Ser	
	410	415	420
Asp Lys Asp Gln	Ala Leu Thr Glu Asp	Glu Gln Gly Lys Ser Tyr	
	425	430	435
Pro Lys Leu Ile	Ser Glu Gly Phe Pro	Gly Ile Pro Ser Pro Leu	
	440	445	450
Asp Thr Ala Phe	Tyr Asp Arg Arg Gln	Lys Leu Ile Tyr Phe Phe	
	455	460	465
Lys Glu Ser Leu	Val Phe Ala Phe Asp	Val Asn Arg Asn Arg Val	
	470	475	480
Leu Asn Ser Tyr	Pro Lys Arg Ile Thr	Glu Val Phe Pro Ala Val	
	485	490	495
Ile Pro Gln Asn	His Pro Phe Arg Asn	Ile Asp Ser Ala Tyr Tyr	
	500	505	510
Ser Tyr Ala Tyr	Asn Ser Ile Phe Phe	Phe Lys Gly Asn Ala Tyr	
	515	520	525
Trp Lys Val Val	Asn Asp Lys Asp Lys	Gln Gln Asn Ser Trp Leu	
	530	535	540
Pro Ala Asn Gly	Leu Phe Pro Lys Lys	Phe Ile Ser Glu Lys Trp	
	545	550	555
Phe Asp Val Cys	Asp Val His Ile Ser	Thr Leu Asn Met	
	560	565	

<210> 20

<211> 306

<212> PRT

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 4333459CD1

<400> 20

Met Ser Leu Lys	Met Leu Ile Ser Arg Asn	Lys Leu Ile Leu Leu	
1	5	10	15
Leu Gly Ile Val	Phe Phe Glu Arg Gly Lys	Ser Ala Thr Leu Ser	
	20	25	30
Leu Pro Lys Ala	Pro Ser Cys Gly Gln Ser	Leu Val Lys Val Gln	
	35	40	45
Pro Trp Asn Tyr	Phe Asn Ile Phe Ser Arg	Ile Leu Gly Gly Ser	
	50	55	60
Gln Val Glu Lys	Gly Ser Tyr Pro Trp Gln	Val Ser Leu Lys Gln	
	65	70	75
Arg Gln Lys His	Ile Cys Gly Gly Ser Ile	Val Ser Pro Gln Trp	
	80	85	90
Val Ile Thr Ala	Ala His Cys Ile Ala Asn	Arg Asn Ile Val Ser	
	95	100	105
Thr Leu Asn Val	Thr Ala Gly Glu Tyr Asp	Leu Ser Gln Thr Asp	
	110	115	120
Pro Gly Glu Gln	Thr Leu Thr Ile Glu Thr	Val Ile Ile His Pro	
	125	130	135

```

His Phe Ser Thr Lys Lys Pro Met Asp Tyr Asp Ile Ala Leu Leu
140 145 150
Lys Met Ala Gly Ala Phe Gln Phe Gly His Phe Val Gly Pro Ile
155 160 165
Cys Leu Pro Glu Leu Arg Glu Gln Phe Glu Ala Gly Phe Ile Cys
170 175 180
Thr Thr Ala Gly Trp Gly Arg Leu Thr Glu Gly Gly Val Leu Ser
185 190 195
Gln Val Leu Gln Glu Val Asn Leu Pro Ile Leu Thr Trp Glu Glu
200 205 210
Cys Val Ala Ala Leu Leu Thr Leu Lys Arg Pro Ile Ser Gly Lys
215 220 225
Thr Phe Leu Cys Thr Gly Phe Pro Asp Gly Gly Arg Asp Ala Cys
230 235 240
Gln Gly Asp Ser Gly Gly Ser Leu Met Cys Arg Asn Lys Lys Gly
245 250 255
Ala Trp Thr Leu Ala Gly Val Thr Ser Trp Gly Leu Gly Cys Gly
260 265 270
Arg Gly Trp Arg Asn Asn Val Arg Lys Ser Asp Gln Gly Ser Pro
275 280 285
Gly Ile Phe Thr Asp Ile Ser Lys Val Leu Ser Trp Ile His Glu
290 295 300
His Ile Gln Thr Gly Asn
305

```

<210> 21
 <211> 953
 <212> PRT
 <213> Homo sapiens

<220>
 <221> misc_feature
 <223> Incyte ID No: 6817347CD1

```

<400> 21
Met Thr Leu Leu Ala Pro Trp Tyr Thr Gly Pro Met Ile Pro Met
1 5 10 15
Asp Val Asn Glu Pro Ser Ser Val Thr Thr Ala Pro Thr Leu Ser
20 25 30
Ser Ser Leu Gln His Ile Ser Ser Phe Leu Ala Thr Gly Lys Lys
35 40 45
Leu Ser Leu His Phe Gly His Pro Arg Glu Cys Glu Val Thr Arg
50 55 60
Ile Asp Asp Lys Asn Arg Arg Gly Leu Glu Asp Ser Glu Pro Gly
65 70 75
Ala Lys Leu Phe Asn Asn Asp Gly Val Cys Cys Cys Leu Gln Lys
80 85 90
Arg Gly Pro Val Asn Ile Thr Ser Val Cys Val Ser Pro Arg Thr
95 100 105
Leu Gln Ile Ser Val Phe Val Leu Ser Glu Lys Tyr Glu Gly Ile
110 115 120
Val Lys Phe Glu Ser Asp Glu Leu Pro Phe Gly Val Ile Gly Ser
125 130 135
Asn Ile Gly Asp Ala His Phe Gln Glu Phe Arg Ala Gly Ile Ser
140 145 150
Trp Lys Pro Val Val Asp Pro Asp Asp Pro Ile Pro Gln Phe Pro
155 160 165
Asp Cys Cys Ser Ser Ser Ser Ser Arg Ile Pro Ser Val Ser Val
170 175 180
Leu Val Ala Val Pro Leu Val Ala Gly His Lys Gly Gln Ala Phe
185 190 195
Ile Glu Arg Met Leu Gly Cys Phe Lys Glu Leu Lys Gln Glu Leu
200 205 210
Thr Gln Glu Gly Pro Gly Gly Gly His Pro Arg Ser Ala Trp Pro
215 220 225
Pro Arg Arg His Ala Gln Trp Pro Pro Glu Pro Cys Glu Gln Gly
230 235 240

```

Glu	Glu	Pro	Pro	Pro	Val	Glu	Ala	Glu	Glu	Val	Glu	Glu	Ala	Glu
				245					250					255
Thr	Ala	Glu	Lys	Ala	Glu	Arg	Lys	Val	Glu	Ala	Glu	Ala	Lys	Val
				260					265					270
Glu	Gly	Lys	Ala	Glu	Ala	Ala	Gly	Lys	Ala	Glu	Ala	Ala	Gly	Lys
				275					280					285
Val	Asp	Ala	Thr	Glu	Lys	Val	Glu	Thr	Ala	Gly	Lys	Val	Asp	Ala
				290					295					300
Ala	Gly	Lys	Val	Glu	Thr	Ala	Glu	Gly	Pro	Gly	Arg	Arg	Ala	Glu
				305					310					315
Leu	Lys	Leu	Glu	Pro	Glu	Pro	Glu	Pro	Val	Arg	Glu	Ala	Glu	Gln
				320					325					330
Glu	Pro	Lys	Gln	Glu	Leu	Glu	Asp	Glu	Asn	Pro	Ala	Arg	Ser	Gly
				335					340					345
Gly	Gly	Gly	Asn	Ser	Asp	Glu	Val	Pro	Pro	Pro	Thr	Leu	Pro	Ser
				350					355					360
Asp	Pro	Pro	Arg	Pro	Pro	Asp	Pro	Ser	Pro	Arg	Arg	Ser	Arg	Ala
				365					370					375
Pro	Arg	Arg	Arg	Pro	Arg	Pro	Arg	Pro	Gln	Thr	Arg	Leu	Arg	Thr
				380					385					390
Pro	Pro	Gln	Pro	Arg	Pro	Arg	Pro	Pro	Pro	Arg	Pro	Arg	Pro	Arg
				395					400					405
Arg	Gly	Pro	Gly	Gly	Gly	Cys	Leu	Asp	Val	Asp	Phe	Ala	Val	Gly
				410					415					420
Pro	Pro	Gly	Cys	Ser	His	Val	Asn	Ser	Phe	Lys	Val	Gly	Glu	Asn
				425					430					435
Trp	Arg	Gln	Glu	Leu	Arg	Val	Ile	Tyr	Gln	Cys	Phe	Val	Trp	Cys
				440					445					450
Gly	Thr	Pro	Glu	Thr	Arg	Lys	Ser	Lys	Ala	Lys	Ser	Cys	Ile	Cys
				455					460					465
His	Val	Cys	Gly	Thr	His	Leu	Asn	Arg	Leu	His	Ser	Cys	Leu	Ser
				470					475					480
Cys	Val	Phe	Phe	Gly	Cys	Phe	Thr	Glu	Lys	His	Ile	His	Glu	His
				485					490					495
Ala	Glu	Thr	Lys	Gln	His	Asn	Leu	Ala	Val	Asp	Leu	Tyr	Tyr	Gly
				500					505					510
Gly	Ile	Tyr	Cys	Phe	Met	Cys	Lys	Asp	Tyr	Val	Tyr	Asp	Lys	Asp
				515					520					525
Ile	Glu	Gln	Ile	Ala	Lys	Glu	Glu	Gln	Gly	Glu	Ala	Leu	Lys	Leu
				530					535					540
Gln	Ala	Ser	Thr	Ser	Thr	Glu	Val	Ser	His	Gln	Gln	Cys	Ser	Val
				545					550					555
Pro	Gly	Leu	Gly	Glu	Lys	Phe	Pro	Thr	Trp	Glu	Thr	Thr	Lys	Pro
				560					565					570
Glu	Leu	Glu	Leu	Leu	Gly	His	Asn	Pro	Arg	Arg	Arg	Arg	Ile	Thr
				575					580					585
Ser	Ser	Phe	Thr	Ile	Gly	Leu	Arg	Gly	Leu	Ile	Asn	Leu	Gly	Asn
				590					595					600
Thr	Cys	Phe	Met	Asn	Cys	Ile	Val	Gln	Ala	Leu	Thr	His	Thr	Pro
				605					610					615
Ile	Leu	Arg	Asp	Phe	Phe	Leu	Ser	Asp	Arg	His	Arg	Cys	Glu	Met
				620					625					630
Pro	Ser	Pro	Glu	Leu	Cys	Leu	Val	Cys	Glu	Met	Ser	Ser	Leu	Phe
				635					640					645
Arg	Glu	Leu	Tyr	Ser	Gly	Asn	Pro	Ser	Pro	His	Val	Pro	Tyr	Lys
				650					655					660
Leu	Leu	His	Leu	Val	Trp	Ile	His	Ala	Arg	His	Leu	Ala	Gly	Tyr
				665					670					675
Arg	Gln	Gln	Asp	Ala	His	Glu	Phe	Leu	Ile	Ala	Ala	Leu	Asp	Val
				680					685					690
Leu	His	Arg	His	Cys	Lys	Gly	Asp	Asp	Val	Gly	Lys	Ala	Ala	Asn
				695					700					705
Asn	Pro	Asn	His	Cys	Asn	Cys	Ile	Ile	Asp	Gln	Ile	Phe	Thr	Gly
				710					715					720
Gly	Leu	Gln	Ser	Asp	Val	Thr	Cys	Gln	Ala	Cys	His	Gly	Val	Ser
				725					730					735
Thr	Thr	Ile	Asp	Pro	Cys	Trp	Asp	Ile	Ser	Leu	Asp	Leu	Pro	Gly

	740		745		750
Ser Cys Thr Ser	Phe Trp Pro Met Ser	Pro Gly Arg Glu Ser	Ser		
	755		760		765
Val Asn Gly Glu	Ser His Ile Pro Gly	Ile Thr Thr Leu Thr	Asp		
	770		775		780
Cys Leu Arg Arg	Phe Thr Arg Pro Glu	His Leu Gly Ser Ser	Ala		
	785		790		795
Lys Ile Lys Cys	Gly Ser Cys Gln Ser	Tyr Gln Glu Ser Thr	Lys		
	800		805		810
Gln Leu Thr Met	Asn Lys Leu Pro Val	Val Ala Cys Phe His	Phe		
	815		820		825
Lys Arg Phe Glu	His Ser Ala Lys Gln	Arg Arg Lys Ile Thr	Thr		
	830		835		840
Tyr Ile Ser Phe	Pro Leu Glu Leu Asp	Met Thr Pro Phe Met	Ala		
	845		850		855
Ser Ser Lys Glu	Ser Arg Met Asn Gly	Gln Leu Gln Leu Pro	Thr		
	860		865		870
Asn Ser Gly Asn	Asn Glu Asn Lys Tyr	Ser Leu Phe Ala Val	Val		
	875		880		885
Asn His Gln Gly	Thr Leu Glu Ser Gly	His Tyr Thr Ser Phe	Ile		
	890		895		900
Arg His His Lys	Asp Gln Trp Phe Lys	Cys Asp Asp Ala Val	Ile		
	905		910		915
Thr Lys Ala Ser	Ile Lys Asp Val Leu	Asp Ser Glu Gly Tyr	Leu		
	920		925		930
Leu Phe Tyr His	Lys Gln Val Leu Glu	His Glu Ser Glu Lys	Val		
	935		940		945
Lys Glu Met Asn	Thr Gln Ala Tyr				
	950				

<210> 22
 <211> 2204
 <212> DNA
 <213> Homo sapiens

<220>
 <221> misc_feature
 <223> Incyte ID No: 275791CB1

<400> 22
 atatgccaat agacctgaca agtctgaatt ggaaactcag attgacagaa tgacgaagaa 60
 gagcttttagc agctgtcttg gagataagta agagagatgc ttcaccatct ctgagtcattg 120
 aagatgatga taagccaact agcagcccag ataccggatt tgcagaagat gatattcaag 180
 aaatgccgga aaatccagac actatggaaa ctgagaagcc caaaacaatc acagagctgg 240
 atcctgccag ttttactgag ataactaaag actgtgatga gaataaagaa aacaaaactc 300
 cagaaggatc tcaggggagaa gttgattggc tccagcagta tgatatggag cgtgaaaggg 360
 aagagcaaga gcttcagcag gcaactggctc agagccttca agagcaagag gcttgggaac 420
 agaaagaaga tgatgacctc aaaagagcta ccgagtttaag tcttcaagag ttttaacaact 480
 cctttgtgga tgcatgggt tctgatgagg actctggaaa tgaggatggt tttgatattgg 540
 agtacacaga agctgaagct gaggaactga aaagaaatgc tgagacagga aatctgcctc 600
 attcgtaccg gctcatcagt gttgtcagtc acattggtag cacttcttct tcaggtcatt 660
 acattagtga tgtatatgac attaagaagc aagcgtgggt tacttacaat gacctggagg 720
 tatcaaaaat ccaagaggct gccgtgcaga gtgatcgaga tcggagtggc tacatcttct 780
 tttatatgca caaggagatc tttgatgagc tgctggaaac agaaaagaac tctcagtcac 840
 ttagcacgga agtggggaag actaccgctc aggcctcgtg aggaacaaac tcttgggttg 900
 gcagcatgca ctgcatattt gttactgctg cccacctcac ctttctctctg ctgaaggaga 960
 atttggaaat ctacttgatg cgggagcaac aaacagctca gggccaaacc aaaagacaaa 1020
 aattggagta acgtagaatg ctccatgcta ttttatggaa actttggtct cacatccgta 1080
 gctgattatc ctctttttct cctatgagtg gcacttcttt tgtcttagga atacatgttg 1140
 taaatatata tctgtgtatg tgtgtataca cacacacaga cacacacgga cacaacggg 1200
 atgaatggag ccttaagag ttaggatgag ccaccagaat atgcctgctc aaaatataata 1260
 gcacagcagt ttggagaaga aatgaagggt tcaaagagtc cattcacctg agaaatgtgt 1320
 gaagacatac ttatcagttg gcttttagct tttatgttcc ttgagtgtt tcaactcaagt 1380
 ctgtaacctt ttgtgtttcc ttattagtaa aattcactgg aaagccagct cttcatgtta 1440
 cactaatgac agtttgttct ctttgcaaga gaggggcatt actgtcacct gacttgagga 1500
 gctgttttgt tgtgtgtgt gtctgcaaat ttcattgaatt tgtgatgtct ttgctgttta 1560
 catgcagtc caagaaatgg attgttggtg ctttgggaata tgttacagtc ccacatttga 1620

```

tatttcttat atacttttgtt ttctctaagg agattttcttc acacagtatg ttcacatcat 1680
atcatcatca ttattatgggt ggtaaagata gaattctttt tctttttttg tcattctggc 1740
catgggagcag cattacccta atggattgca accaaaactt taaacaagta gaaagataat 1800
atttctccaa ttgggactcc ccagcaggaa tacttaggga taaggaagaa tgctagcatc 1860
tctgtctctc aaacataggg aggataagaa gagtgttctt ctggtaaagc taaaattctg 1920
gaccactgaa gctaaaagcc ctattgcaag tatgaaatta agtacttgag ctataggaca 1980
aaccttgggc atttaacat ttactgtctg gctttgccct taaaataggg ttgcaattaa 2040
aatgtgattg gcttaggtaa tccccaaaac taacaaataa caaagggtgca taattttatt 2100
atctactttt taggtgctct gagttgagc aaagtagagc ggcaacatta agtgctatgc 2160
tagtcaacta gctgacgtaa ccagctttgg taagcagctt atga 2204

```

<210> 23
 <211> 2036
 <212> DNA
 <213> Homo sapiens

<220>
 <221> misc_feature
 <223> Incyte ID No: 1389845CB1

```

<400> 23
ccgatggggg ttaggtcca gggcttctgt cgagaccaag gatgccc aaa tatctgggtg 60
gtgggtgctg catacctggg ccctgggcag aacgaagggt atacagcctg ggccaccagg 120
ataagtccag aaccaccag gagctgagga cagacagaag gaccacggag ggggtgacgg 180
gctggtgtga ggattggtgc ccctgggcca ggactctcct ctcttctccc tgctggctcc 240
agaccagagt ccaagcccta ggcagtgcc aacctaccca gccagcctt gaagacagaa 300
tgagaggggt ttctgtctc caggctctgc tcttcttgt gctgggagct gctgggactc 360
agggaaaggaa gtctgcagcc tgcgggcagc ccgcagatgc cagtccggtc gttggggggc 420
gggatggccg ggacggagag tggccgtggc aggcgagcat ccagcatcgt ggggcacacg 480
tgttgggggg gtcgctcatc gccccccagt ggggtgctgac agcggcgcac tgcctcccca 540
ggagggcact gccagctgag taccgctgc gcctgggggc gctgcgtctg ggctccacct 600
cgccccgcac gctctcggtg ccctgtgcac ggggtgctgct gccccggac tactccgagg 660
acggggcccg cggcgacctg gcaactgctg agctgcgtcg cccggtgccc ctgagcgctc 720
gcgtccaacc cgtctgcctg ccctgtcccg gcgcccgcgc gcccgcgggc acaccatgcc 780
gggtcacccg ctggggcagc ctccgcccag gagtgccctt cccagagtgg cgaccgtac 840
aaggagtaag ggtgcccgtg ctggactcgc gcacctgcga cggcctctac cacgtggggc 900
cggacgtgcc ccaggctgag cgcattgtgc tgccctggag tctgtgtgcc ggtaccctcc 960
agggccacaa ggacgcctgc cagggtgtga ccagcctcc cagcctccg gagtccctc 1020
cctgtgcccc gcacctctcc tccctgaact ccaggacca ggacatccc actcaggctc 1080
aggatcctgg cctccaacct agaggcacca cgccaggggt ctggaacctt gagaactgaa 1140
gtcctgggag ggtgggact taggctcctc ttctctctgc aggggtgattc tgggggacct 1200
ctgacctgcc tgcagtctgg gagctgggtc ctgggtgggc tgggtgagctg gggcaagggt 1260
tgtgcctgc ccaaccgtcc aggggtctac accagtgtgg ccacatatag cccctggatt 1320
caggctcgcg tcagcttcta atgctagcgc gtgaggctga cctggagcca gctgctgggg 1380
tccctcagcc tccgtgttca tccaggcacc tgccataacc ccacatccct tctgctcga 1440
ggccaagatg cctaaaaaag ctaaaggcca ccccccccc caccaccac ctctgcctc 1500
ctctctctt tggggatcac cagctctgac tccaccaacc ctcatccagg aatctgccat 1560
gagtcaccag gagtcacact cccactccc ttcttggett gtatttactt ttcttgccc 1620
tggccagggc tgggcgcaag gcacgcagt atgggcaaac caattgctgc ccacttgccc 1680
tgtgtgcccc tctttttctg gagaaagtca gattcacagc atgacagaga tttgacacca 1740
gggagatcct ccatactgg ctttgaggac acggggacca cagccatgag cggcctctaa 1800
gagctgagag acagccggca gggaaatcga accctcagac ccacagccgc aaggcactgg 1860
attctggcag caccctgaag gagctgggaa gtaagttctt cccagcctc cagataagag 1920
ccccgcggc caatccctc atttcaacct aaagagaccc taagcagaga acctagctga 1980
gccactcctg acctacaaag ttgtgactta ataaatgtgt gctttaagct gctcca 2036

```

<210> 24
 <211> 2185
 <212> DNA
 <213> Homo sapiens

<220>
 <221> misc_feature
 <223> Incyte ID No: 1726609CB1

```

<400> 24
gccatgcctc ctgcccacgg ccaccagcaa gctgtcgggc gcagtggagc agtggctgag 60

```

tgcagctgag	cggctgtatg	ggccctacat	gtggggcagg	tacgacattg	tcttcctgcc	120
accttccttc	cccatcgtgg	ccatggagaa	ccctgcctc	accttcacat	tcttcctccat	180
cctggagagc	gatgagttcc	tggatcatga	tgtcatccac	gaggtggccc	acagttggtt	240
cggcaacgct	gtcaccaacg	ccacgtggga	agagatgtgg	ctgagcgagg	gcctggccac	300
ctatgcccag	cgccgtatca	ccaccgagac	ctacggtgct	gccttcacct	gcctggagac	360
tgccttcctc	ctggacgccc	tgcaccggca	gatgaagctt	ctgggagagg	acagcccggg	420
cagcaaactg	cagggtcaagc	tggagccagg	agtgaatccc	agccacctga	tgaacctgtt	480
cacctacgag	aagggctact	gcttcgtgta	ctacctgtcc	cagctctgcg	gagaccacac	540
gcgctttgat	gactttctcc	gagcctatgt	ggagaagtac	aagttcacca	gcgtggtggc	600
ccaggacctg	ctggactcct	tcttgagctt	cttcccggag	ctgaaggagc	agagcgtgga	660
ctgccgggca	gggctggaat	tcgagcgctg	gctcaatgcc	acaggcccgc	cgctggctga	720
gccggacctg	tctcagggat	ccagcctgac	ccggcccgtg	gaggcccttt	tccagctgtg	780
gaccgcagaa	cctctggacc	aggcagctgc	ctcgccagc	gccattgaca	tctccaagtg	840
gaggacacctc	cagacagcac	tcttctggga	cggctcctg	gatgggtccc	cgctgccgca	900
ggaggtggtg	atgagcctgt	ccaagtgtca	ctcctccctg	ctggactcga	tgaacgtgta	960
gatccgcac	cgctggctgc	agattgtggt	ccgcaacgac	tactatcctg	acctccacag	1020
ggtgcggcgc	ttcctggaga	gccagatgtc	acgcatgtac	accatcccgc	tgtacgagga	1080
cctctgcacc	ggtgccctca	agtcccttcg	gctggaggtc	ttctaccaga	cgaggggccg	1140
gctgcacccc	aacctgcgca	gagccatcca	gcagatcctg	tcccaggggc	tggtctccag	1200
cacagagccc	gcctcagagc	ccagcacgga	gctgggcaag	gctgaagcag	acacagactc	1260
ggaagcacag	gccctgctgc	ttggggacga	ggccccagc	agtgccatct	ctctcaggga	1320
cgtcaatgtg	tctgcctagc	cctgttggcg	ggctgacctc	cgacctccca	gacaccacaa	1380
ttgtgccttc	tgtggggccag	gcctgccatg	actgcgtctc	ggctctggcc	atgagctctg	1440
cccaggccca	caagcccctc	cctgggctc	tcccaggcag	ggagaatggg	gagagggacc	1500
tccttgtgtc	tggcagagac	ctgtggacct	ggcctcccca	ctcccagctc	tcttgccactg	1560
caggccctgg	ggccagcccg	cacacacccat	gcctcctgtc	tcaaacactga	cagctgtgtcc	1620
tagccccgga	tgccagcacc	tgccaggtgc	cgccccgggg	caaggggccc	agcagcccta	1680
tggtgaccgc	cacactgtgc	cttaatgtct	gccggggggc	caggctgtgc	tgtccctgca	1740
gcacgcctcc	ttgcagggat	ctgagccacc	ctccccgcac	agccctgcac	cccggccccta	1800
gggttggcag	cctcagttgg	cccctggcag	aggaacaagg	acacagacat	tccctcagtg	1860
tggggggcag	gggacacagg	gagaggatgg	ttgtccctgg	ggaggggcct	ctggcccacg	1920
gcaaccttag	cccctcagaa	cagggagtcc	caggaccacg	ggagagtgtg	gggacaggac	1980
agcctgtctc	ttgtagcttc	ctgggggtgg	aggcacaggg	gcaaagcaat	accccaggga	2040
aagtgggagg	tgggtgctgg	gctctctcca	ggccaccat	gctgggagag	gcgccagag	2100
cctggggcct	ccagcctggg	actgctgtga	tggggtatca	cggtgatggt	cccatataac	2160
ttccactctg	caaaaaaaaa	aaaaa				2185

<210> 25

<211> 3486

<212> DNA

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 4503848CB1

<400> 25

ctgtctttaa	aaaagaggga	gggaagatta	ctgatttaat	ttataaagga	gaattattat	60
agctccaaca	cctgacttta	tttatctata	tggtttaatt	acacaaacaa	ttcagtgttt	120
gaattataca	aatttcatta	aaactatgta	attatgcaag	aaaaatagga	aatacagggg	180
cacttagttt	tgtgcatatg	tgttcacctg	agagtatttg	cttgtttttt	taaaaagggt	240
ctttttaatt	taatatatta	ttttataatg	cacattcata	tgttgacttt	ggaccaacag	300
aaatccctaa	ttcttattct	ttttctgatt	cttttttagag	ttggtgggtc	caggatttta	360
ctcagaatga	cgttaggaag	agaagtgtatg	tctcctcttc	aggcaatgtc	ttcctatact	420
gtggctggca	gaaatgtttt	aagatgggat	ctttcaccag	agcaaatata	aacaagaact	480
gaggagctca	ttgtgcagac	caaacagggt	tacgatgctg	ttggaatgct	cggtattgag	540
gaagtaactt	acgagaactg	tctgcaggca	ctggcagatg	tagaagtaaa	gtatatagtg	600
gaaaggacca	tgctagactt	tccccagcat	gtatcctctg	acaaagaagt	acgagcagca	660
agtacagaag	cagacaaaag	actttctcgt	tttgataatg	agatgagcat	gagaggagat	720
atatttgaga	gaattgttca	tttacaggaa	acctgtgatc	tgggggaagt	aaaacctgag	780
gccagacgat	acttggaaaa	gtcaattaaa	atggggaaaa	gaaatgggct	ccatcttctc	840
gaacaagtac	agaatgaaat	caaatcaatg	aagaaaaagaa	tgagttagct	atgtattgat	900
tttaacaaaa	acctcaatga	ggatgatacc	ttccttgat	tttccaaggc	tgaacttggg	960
gctcttctcg	atgatttcat	tgacagttta	gaaaagacag	atgatgacaa	gtataaaatt	1020
accttaaaat	atccacacta	tttccctgtc	atgaagaaat	gttgatcccc	tgaaccagga	1080
agaaggatgg	aaatggcttt	taatacaagg	tgcaaaaggg	aaaacaccat	aattttgcag	1140
cagctactcc	cactgcgaac	caagggtggc	aaactactcg	gttatagcac	acatgctgac	1200

tctgctcttg	aatgaacac	tgcaagagc	acaagccgcg	taacagcctt	tctagatgat	1260
ttaagccaga	agttaaaacc	cttgggtgaa	gcagaacgag	agttttat	gaatttgaag	1320
aaaaagggaat	gcaaagacag	gggttttgaa	tatgatggga	aatcaatgc	ctgggatcta	1380
tattactaca	tgactcagac	agaggaactc	aagtattcca	tagaccaaga	gttcctcaag	1440
gaatacttcc	caattgaggt	ggtcactgaa	ggcttgctga	acacctacca	ggagttgttg	1500
ggactttcat	ttgaacaaat	gacagatgct	catgtttgga	acaagagtgt	tacactttat	1560
actgtgaagg	ataaagctac	aggagaagta	ttgggacagt	tctatttga	cctctatcca	1620
agggaaggaa	aatacaatca	tgcgccctgc	ttcggctctc	agcctggctg	ccttctgcct	1680
gatggaagcc	ggatgatggc	agtggctgcc	ctcgtggtga	acttctcaca	gccagtgcca	1740
ggctgctccct	ctctcctgag	acacgacgag	gtgaggactt	actttcatga	gtttggtcac	1800
gtgatgcac	agatttgtgc	acagactgat	tttgcacgat	ttagcggaac	aaatgtggaa	1860
actgactttg	tagaggtgcc	atcgcaaatg	cttgaaaatt	gggtgtggga	cgctcgattcc	1920
ctccgaagat	tgtcaaaaca	ttataaagat	ggaagcccta	ttgcagacga	tctgcttgaa	1980
aaactgtgtg	cttctaggt	ggtcaacaca	ggctctctga	ccctgcgcca	gattgttttg	2040
agcaaagttg	atcagtctct	tcataccaac	acatcgtctg	atgctgcaag	tgaatatgcc	2100
aaatactgct	cagaaatatt	aggagttgca	gctactccag	gcacaaatat	gccagctacc	2160
tttggacatt	ttggcagggg	atacgatggc	caatattatg	gatattcttg	gagtgaagta	2220
ttttccatgg	atatgtttta	cagctgtttt	aaaaaagaag	ggataatgaa	tccggaggtt	2280
ggaatgaaat	acagaaacct	aatcctgaaa	cctgggggat	ctctggacgg	catggacatg	2340
ctccacaatt	tccttgaacg	tgagccaaac	caaaaagcgt	tcctaatgag	tagaggcctg	2400
catgctccgt	gaactgggga	tctttggtag	ccgtccatgt	ctggaggaca	agtcgacatc	2460
accatgtgtt	actggccttg	aaactgaagg	gagttttgca	agtgaataat	tagattttcta	2520
ttgacatcct	tttgttttct	aattttaaaa	attataaaga	tgtaaatgga	attataaata	2580
ctgtgacctg	agaaaagacc	cactagaaag	taattgtact	ataaaaattc	ataaaaactgg	2640
atattgatttc	tttttatgaa	agtttcatat	gaatgttaact	tgatttttta	ctattataat	2700
ctagataata	tgatataaga	gggctaagaa	tttttaaat	gaatcatata	tatgatataa	2760
tttgatccct	cttgatctct	gaagttttgt	acttgggatt	tctggactga	taaatgaatc	2820
atcacattct	tctgttaaat	attttcttgg	agctctgtgt	caactttgat	cctttgtctc	2880
ccaggaaggt	gtgacctctc	ctttgacctgc	atacctcaag	gccaggggaa	tatgcctcag	2940
tgatgcattt	atctttgtat	atcaggccgc	atgattccca	actttctgcc	acacttaaat	3000
tacgttctct	catttcaagt	ttgtcttttc	tgctaaagt	tcagtcaaag	agtatcaaaa	3060
aattatgttt	cagctagact	gggtgaatgt	ataagttttt	gtatcttgta	ttagaggatt	3120
tcgtagcttt	tattagaggc	tcatttccac	ctcagcatac	aagatcgtaa	gtcttttggc	3180
atgtgtgcca	attagaatac	taaagcaagt	ccaagcacat	ttttctcttc	tcacgtttct	3240
aataagtggt	agggactttg	cctcttttac	ttaccacgtc	cccaaaagtg	tcaggtagac	3300
atgtcacaaa	tggctctgta	gagagccatg	ggaagagaga	ggaggtggat	gtggaacata	3360
aagggttcag	aaactccaga	agaggagtgg	gttttgata	gaagcatttg	aggacagctg	3420
ctccaaagcc	ttatgtgtat	gatgaaactt	aaccacgggg	aagagactct	tcagtgcct	3480
gttctg						3486

<210> 26

<211> 2847

<212> DNA

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 5544089CB1

<400> 26

caatgacgct	tggacgagga	tttattttcta	caagctaatt	gaatccagga	gcagctttta	60
ttattaacac	taacggaaga	gaaaaggagt	atttccaagg	gctcaaatgg	aagctgtact	120
cagtcgggtg	gaggcagggg	gaggtaaagt	ttctcacact	caagtcgtct	tcatagttaa	180
ctgtcccttt	ccaaacaaaa	gctaataacg	ccatacgcat	ccacacactc	cctcctggat	240
gaacctaaat	ctcgtcccca	ctgtcaccct	aaggccagtt	atcaaaaact	gttccttctc	300
tgcctcaaaa	gactgaagcc	gcaggccctg	ttctgctctt	gctcaggaat	ctgattgtct	360
ttaaagtgtc	cttacaagat	tccgtcgatg	tttgcctcct	ctgtcttgta	atcaggacta	420
agtgggtggg	catcaaaagg	tagaaagatg	gaacttatcc	agccaaagga	gccaaactta	480
cagtacattt	ctctttgtca	tgaattgcat	actttgttcc	aagtcagtgt	gtctggaaag	540
tgggcgttgg	tctcaccatt	tgctatgcta	cactcagttg	ggagactcat	tcctgccttt	600
cgtgggttacg	cccaacaaga	cgctcaggaa	tttctttgtg	aacttttaga	taaaatacaa	660
cgtgaattag	agacaactgg	taccagttta	ccagctctta	tccccacttc	tcaaaggaaa	720
ctcatcaaac	aagttctgaa	tgttgttaaat	aacatttttc	atggacaact	tcttagtcag	780
gttacatgtc	ttgcatgtga	caacaaatca	aataccatag	aacctttctg	ggacttgtca	840
ttggagtttc	cagaaaggta	tcaatgcagt	ggaaaagata	ttgcttccca	gccatgtctg	900
gttactgaaa	tgttggccaa	atttacagaa	actgaagctt	tagaaggaaa	aatctacgta	960
tgtgaccagt	gtaactcaaa	gcgtagaagg	ttttcctcca	aaccagttgt	actcacagaa	1020


```

gccagaaac aacttatgat atgccaccta cctcagggttc tcagactgca cctcaaacga 1080
ttcagggtggt caggacgtaa taaccgagag aagattgggtg ttcatgttgg ctttgaggaa 1140
atcttaaaaca tggagcccta ttgctgcagg gagaccctga aatccctcag accagaatgc 1200
tttatctatg acttgctccgc ggtggtgatg caccatggga aaggatttgg ctcagggcac 1260
tacactgcct actgctataa ttctgaagga gggttctggg tacactgcaa tgattccaaa 1320
ctaagcatgt gcactatgga tgaagtatgc aaggctcaag cttatatctt gttttatacc 1380
caacgagtta ctgagaatgg acattctaaa cttttgectc cagagctcct gttggggagc 1440
caacatccca atgaagacgc tgatacctcg tctaataaaa tccttagctg atccaaagac 1500
aatgggggttt tcttcctgtg atttatatat atacttttta aaagactgat gtaccatttt 1560
aaacttcatt ttttcttggt aatcagtgta tactacattt atacatttta tatctaacia 1620
tttttttttt tacaaagtat aaatgtatat atcaactgaa ggtaactact tttttcata 1680
ttggagtttt aaacttttgg tgtttacctc agactgatgt tacctctttt atatttttat 1740
gtcttaattg gctcggatga tgaacttggt caatcttcta ccaacaaagt tcaagtggca 1800
tcattttata tacatgtatc tttttcaggt attttctata caaattctta atagatggaa 1860
aattagactc tactttgggtc actaatagtc ttctatttgt atattgaagt taccttgccc 1920
cttggagtta ttgaagtgc atgtcaaggt atcacctaaa tattcttcag tcacactcac 1980
tggtatttct gaggcttggg gtgttaacag gccttgtaat tgacattatt ttggttaatt 2040
taaccccaaa atgtctttag taattgctct ttggcatagt caaactataa atgaaaatgg 2100
cagctttaca aatagtatat ttaagtgaac tctggaaacta tggacatgaa aaaaatgatg 2160
gctgggattt atgatttttg tctggcagca aacagggttg tccagaagtc taataattaa 2220
gcatttcata aaagtctgaa tttagttaaac cagtgtatga tgttattcaa atagtttacc 2280
ttgggtatga gttcatttta taatgtctga tgacattaga tctcttaaaa ctttatgtat 2340
tttttttagt tcaaaggaat agagtcttga agagaaaaaa ttatagggca gaaaagataa 2400
gtgttcaaaa ttggcaactg gactattatt atgtctagca tctcattcta aataactaaa 2460
gcttgattta ctcttgctag gattatgtga ctactaggta ggagcctctt aaaacactgg 2520
ccctgagcat taaaaaaaaa aaaaaaaaaa aaaaagctatc tatctaaact tgcaaaaaaa 2580
aaattccggg gggggtcacc cttttccttc ttctgaaaat ctcacggggg ttctttaaa 2640
ccctgttgct ccaaacctta tcttttttgg ggggggtaga atcaccta atctgttagac 2700
cagctatgtt tctaagctct gtttaaccac gggagatctg gtacccctt tttaaaagg 2760
ggtttatatt cgggttgaa gcttagtgaa aagtagtccc ctggagaatg cgggccaccc 2820
ctgggggcca tctgttaggt aaaactt 2847

```

<210> 27
 <211> 890
 <212> DNA
 <213> Homo sapiens

<220>
 <221> misc_feature
 <223> Incyte ID No: 7474081CB1

```

<400> 27
gaggccaaga attcggcacg aggcacttac tccctgagct aagggggaag agctggatca 60
ccatgaaata tgtcttctat ttgggtgtcc tcgctgggac atttttctt gctgactcat 120
ctgttcagaa agaagaccct gctccctatt tgggtgtacct caagtctcac ttcaaccctt 180
gtgtgggctg cctcatcaaa ccagctggg tgctggcccc agctcactgc tatttaccac 240
atctgaaagt gatgctggga aatttcaaga gcagagtcag agacggtact gaacagacaa 300
ttaaccccat tcagatcgtc cgctactgga actactagca tagcgcccca caggatgacc 360
tcattgctcat caagctggct aagcctgcca tgcctcaatc caaagtccag ccccttacc 420
tcgccaccac caatgtcagg ccaggcactg tctgtctact ctcagggttg gactggagcc 480
aagaaaacag tggccgacac cctgacttgc ggcagaacct ggaggcccc gtgatgtctg 540
atcgagaatg ccaaaaaaca gaacaaggaa aaagccacag gaattcctta tgtgtgaaat 600
ttgtgaaagt attcagccga atttttgggg aggtggcctg tgctactgtc atctgcaaag 660
acaagctcca ggggaatcag gtggggcact tcatgggagg ggacgtcggc atctacacca 720
atgttttaca atgttatcc tggattgaga acactgctaa ggacaagtga gacctactt 780
ctccctctgc attccactgg ctctgcatg gactatacaa gcagataatt ttccctctat 840
tcaaaataaa atctccaaat gaaaatttgg gaatgtagca aaaaaaaaaa 890

```

<210> 28
 <211> 1577
 <212> DNA
 <213> Homo sapiens

<220>
 <221> misc_feature
 <223> Incyte ID No: 5281209CB1

<400> 28

atgcagccca	cgggccgcga	gggttccccg	gcgctcagcc	ggcggatatct	gcggcgtctg	60
ctgtctctgc	tactgtctgt	gctgtgcgg	cagcccgtaa	cccgccgga	gaccacgcgc	120
ggcgccccca	gagccctctc	cacgtctggc	tccccagcc	tcttcaccac	gcccgggtgc	180
ccagcgcccc	tactaccccc	aggcctcact	acgccaggca	cccccaaac	cctggacctt	240
cggggctgcg	cgcaggccct	gatgcggagt	ttccactcg	tggacggcca	caatgacctg	300
ccccagggtcc	tgagacagcg	ttacaagaat	gtgcttcagg	atgttaacct	gcgaaatttc	360
agccatggtc	agaccagcct	ggacaggctt	agagacggcc	tctgtgggtg	ccagttcttg	420
tcagcctccg	tctcatgcca	gtcccaggac	cagactgccg	tgcgcctcgc	cctggagcag	480
attgacctca	ttcacgcgat	gtgtgcctcc	tactctgaac	tcgagcttgt	gacctcagct	540
gaaggctctga	acagctctca	aaagctggcc	tgctctattg	gcgtggaggg	tggtcactca	600
ctggacagca	gcctctctgt	gctgcgcagt	ttctatgtgc	tgggggtgcg	ctacctgaca	660
cttaccttca	cctgcagtac	accatgggca	gagagtcca	ccaagtccag	acaccacatg	720
tacaccaacg	tcagcggatt	gacaagcttt	ggtgagaaag	tagtagagga	ggtgaaccgc	780
ctgggcatga	tgatagattt	gtcctatgca	tccgacacct	tgataagaag	ggtcctggaa	840
gtgtctcagg	ctctgtgat	cttctccac	tcagctgcc	gagctgtgtg	tgacaatttg	900
ttgaatgttc	ccgatgat	cctgcagctt	ctgaagaaga	acggtggcat	cgtgatggtg	960
acactgtcca	tgggggtgct	gcagtgcac	ctgcttgcta	acgtgtccac	tgtggcagat	1020
cactttgacc	acatcagggc	agtcattgga	tctgagttca	tccggattgg	tggaatttat	1080
gacgggactg	gccggttccc	tcaggggctg	gaggatgtgt	ccacataccc	agtcctgata	1140
gaggagttgc	tgagtcgtag	ctggagcgag	gaagagcttc	aaggtgtcct	tctggaaac	1200
ctgtctcggg	tcttcagaca	agtggaaaag	gtgagagagg	agagcagggc	gcagagcccc	1260
gtggaggctg	agtttccata	tgggcaactg	agcacatcct	gccactccca	cctcgtgctt	1320
cagaatggac	accagctac	tcattctggag	gtgaccaagc	agccaacca	tccggtcccc	1380
tgagggtcct	caaatgcctc	cccatacctt	gttccaggcc	ttgtggctgc	tgccaccatc	1440
ccaaccttca	ccaagtggct	ctgctgacac	agtcggctcc	cgcagaggtc	actgtggcaa	1500
agcctcacaa	agccccctct	cctagttcat	tcacaagcat	atgtctagaa	taaacatgtt	1560
acacatggaa	aaaaaaaa					1577

<210> 29

<211> 1958

<212> DNA

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 2256251CB1

<400> 29

aagcggtcga	gctcggcatt	cattgtaacg	gcgccatgtg	ctggaaagg	cgtgtggttt	60
ctgtctcgcat	ctctcggttg	agtggggctg	gtcgggggtg	gctgcagggc	tgtctcccc	120
accaccactg	tagtcagctc	gtaccttggg	agatgctctg	aggccatgaa	acaacctggc	180
cctcctcgaa	ctttctcccc	acagcgtccc	caccgtggcc	ctggaaccag	ctgggggctt	240
tgcggtgtgg	gagagccggt	gccccagccc	acacccgctg	cctatctata	aacatctctg	300
tctgtctaca	tcaccagctt	ccttccattc	agcccagtg	gcacactcca	tcaccagcac	360
aattatccag	ctcaggcaga	cccagggtgt	gccagtggtg	ctccagctga	cttctcttta	420
atttcttctt	aacttactga	ggtgaagttt	acagaagata	aagttacatt	atcaagcgtc	480
caattcagag	gccccgagca	ccttggacag	tgctgtctgc	ccccgcctcc	ccgaatttta	540
tccagcttca	aatatctcca	ccacccctga	aggaaaaccg	gggcccacta	ggcagccacc	600
cctagcacc	cgggccttct	cgggggctcc	accgttctga	gcccccttct	agccgcctag	660
gggccctctg	cagcctttcc	accgcctccg	ggagccctgg	ttagtttggt	gagcatgggt	720
cttagaaaag	acgacctgag	ccccaggcgc	gctcactgct	cctgagacgt	catttctgct	780
gcacccacga	tgcttctggg	gagagtctgg	cagacgagag	agctgaagag	caaagtcccc	840
aagaaggcag	ggagggtgtg	tcagggaagg	cttcatggag	gaagtgcagt	gggcttcttg	900
ggatccccac	caggcacccc	ttctctcttc	gacttagggt	gtggccggcc	gcagggttctg	960
gatgcaggcg	gccggatcgt	ggggggctac	gctgccccgg	ccggcgcatg	gccatggcag	1020
gccagcctcc	gcctgcggag	ggtgcacgtg	tgccgggggt	cactgctcag	ccccagtggt	1080
gtgctcacag	ctgcccactg	cttctccggg	tccctgaact	catccgacta	ccagggtgcac	1140
ctgggggaac	tgagatcac	tctgtctccc	cacttctcca	ccgtgaggca	gatcatcctg	1200
cactccagcc	cctcaggaca	gccggggacc	agcggggaca	tcgcccgtgt	ggagctcagt	1260
gtccccgtga	ccctcttcag	ccggatcctg	cccgtctgcc	tcccggagge	ctcagatgac	1320
ttctgccctg	ggatccggtg	ctgggtgacc	ggctggggct	atacgcgga	gggagagcct	1380
ctgccacccc	cgtacagcct	gcgggaggtg	aaagtctccg	tggtggacac	agagacctgc	1440
cgccgggact	atcccggccc	cgggggcagc	atccttcagc	ccgacatgct	gtgtgcccgg	1500
ggccccgggg	atgctcgcca	ggcgaactcc	ggggggcctc	tggtctgcca	ggtgaacggt	1560
gcctgggtgc	aggctggcat	tgtgagctgg	ggtgagggtc	gcggccgccc	caacaggccg	1620
ggagtctaca	ctcgtgtccc	tgcctacgtg	aactggatcc	gccgccacat	cacagcatca	1680

```

gggggctcag agtctgggta ccccaggctc cccctcctgg ctggcttatt cctccccggc 1740
ctcttccttc tgctagtctc ctgtgtcctg ctggccaagt gcctgtctgca cccatctgcg 1800
gatgggtactc ccttccccgc ccttgactga tggcaggaat ccaagtgcac ttcttaata 1860
agttactatt tattccgctc cgccccctcc ctctcccttg agaagctgag tcttctgcat 1920
cagattattg caacatttaa cctgaattta acgacacc
1958

```

<210> 30

<211> 3106

<212> DNA

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 7160544CB1

<400> 30

```

gtcccgaggc caaggccgct gctactgccg ccgctgcttc ttagtgccgc gttcgccgcc 60
tgggttggtca ccggcgccgc cgctgaggaa gccactgcaa ccaggaccgg agtggaggcg 120
gcgcagcatg aagcggcgca ggcccgtccc atagcgcacg tcgggacggg ccgggcgggg 180
ccggggggaa ggaaaatgca acatggcagc agcaatggaa acagaacagc tgggtgttga 240
gatatttgaa actcgggact gtgaggagaa tattgaatca caggatcggc ctaaatggga 300
gcctttttat gttgagcggg attcctggag tcagcttaa aagctgcttg ccgataccag 360
aaaaatcatg ggctacatga tggctaaggc accacatgat ttcatgtttg tgaagaggaa 420
tgatccagat ggacctcatt cagacagaat ctattacctt gccatgtctg gtgagaacag 480
agaaaataca ctgttttatt ctgaaattcc caaaactatc aatagagcag cagtcttaat 540
gctctcttgg aagcctcttt tggatctttt tcaggcaaca ctggactatg gaatgtattc 600
tcgagaagaa gaactattaa gagaaagaaa acgcattgga acagtcggaa ttgcttctta 660
cgattatcac caagggaagt gaacatttct gtttcaagcc ggtagtggaa tttatcacgt 720
aaaagatgga gggccacaag gatttacgca acaaccttta aggcccaatc tagtggaac 780
tagttgtccc aacatacggg tggatccaaa attatgccct gctgatccag actggattgc 840
ttttatacat agcaacgata tttggatctc taacatcgta accagagaag aaaggagact 900
cacttatgtg cacaatgagc tagccaacat ggaagaagat gccagatcag ctggagtcgc 960
tacctttggt ctccaagaag aatttgatag atattctggc tattggtggt gtccaaaagc 1020
tgaaacaact cccagtgggt gtaaaattct tagaattcta tatgaagaaa atgatgaatc 1080
tgaggtggaa atttctcatg ttacatcccc tatgttgaa acaaggaggg cagattcatt 1140
ccgttatcct aaaacaggta cagcaaatcc taaagtcaat tttaagatgt cagaaataat 1200
gattgatgct gaagggaagg tcatagatgt catagataag gaactaatc aaccttttga 1260
gattctatct gaaggagtgt aatatattgc cagagctgga tggactcctg agggaaaata 1320
tgcttggtcc atctactag atcgctccca gactcgccta cagatagtgt tgatctcacc 1380
tgaattatct atcccagtag aagatgatgt tatggaaagg cagagactca ttgagtcagt 1440
gcctgattct gtgacgccac taattatcta tgaagaaaca acagacatct ggataaatat 1500
ccatgacatc ttctatgttt tccccaaag tcacgaagag gaaattgagt ttatttttgc 1560
ctctgaatgc aaaacagggt tccgtcattt atacaaaatt acatctatct taaaggaaag 1620
caaatataaa cgatccagtg gtgggctgcc tgctccaagt gatttcaagt gtctatcaa 1680
agaggagata gcaattacca gtgggtgaatg ggaagttctt ggccggcatg gatctaata 1740
ccaagttgat gaagtcagaa ggctggtata ttttgaaggc accaaagact cccctttaga 1800
gcatcacctg tacgtagtca gttacgtaaa tcttgagag gtgacaaggc tgactgaccg 1860
tggtactca cattcttgct gcatcagtca gcaactgtac ttctttataa gtaagtata 1920
taaccagaag aatccacact gtgtgtccct ttacaagcta tcaagtctg aagatgaccc 1980
aacttgcaa acaaagggaat tttgggccac cattttggat tcagcaggtc ctcttctga 2040
ctatactcct ccagaaattt tctcttttga aagtactact ggatttacat tgtatgggag 2100
gctctacaag cctcatgac tacagcctgg aaagaaatat cctactgtgc tgttcatata 2160
tggtgtcctc caggtgcagt tgggtgaataa tcggtttaaa ggagtcaagt atttccgctt 2220
gaatacccta gcctctctag gttatgtggt tgtagtata gacaacaggg gatcctgtca 2280
ccgagggctt aaatttgaag gcgcctttaa atataaaatg ggtcaaatag aaattgacga 2340
tcagggtgaa ggaactccat atctagcttc tcgatatgat ttcatgact tagatcgtgt 2400
gggcatccac ggctggtcct atggaggata cctctccctg atggcattaa tgcagagggtc 2460
agatatcttc aggggttgcta ttgctggggc ccagtcact ctgtggatct tctatgatac 2520
aggatacacg gaacgttata tgggtcaccc tgaccagaat gaacagggtc attacttagg 2580
atctgtggcc atgcaagcag aaaagttccc ctctgaacca aatcgtttac tgctcttaca 2640
tggtttcctg gatgagaatg tccattttgc acataccagt atattactga gttttttagt 2700
gagggtctga aagccatatg atttacagat ctactctcag gagagacaca gcataagagt 2760
tcctgaatcg ggagaacatt atgaactgca tcttttgac taccttcaag aaaaccttgg 2820
atcacgtatt gctgctctaa aagtgatata attttgacct gtgtagaact ctctgggtata 2880
cactggctat ttaaccaaat gaggagggtt aatcaacaga aaacacagaa ttgatcatca 2940
cattttgata cctgccatgt aacatctact cctgaaaata aatgtggtgc catgcagggg 3000
tctacggttt gtggtagtaa tctaatacct taacccaca tgctcaaaat caaatgatac 3060

```

atattcctga gagaccagc aataccataa gaattactaa aaaaaa

3106

<210> 31

<211> 3567

<212> DNA

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 7477386CB1

<400> 31

```

atggctccac tccgcgcgct gctgtcctac ctgtctgcctt tgcactgtgc gctctgcgcc 60
gccgcgggca gccggacccc agagctgcac ctctctggaa agctcagtga ctatggtgtg 120
acagtgcctt gcagcacaga ctttcgggga cgcttcctct cccacgtggt gtctggccca 180
gcagcagcct ctgcagggag catggtagtg gacacgccac ccacactacc acgacactcc 240
agtcacctcc ggggtggctcg cagccctctg caccacaggag ggacctgtg gcctggcagg 300
gtggggcgcc actccctcta cttcaatgtc actgttttcg ggaaggaact gcacttgccg 360
ctgcggccca atcggaggtt ggtagtgcca ggatcctcag tggagtggca ggaggatttt 420
cgggagctgt tccggcagcc cttacggcag gagtgtgtgt acactggagg tgtcactgga 480
atgcctgggg cagctgttgc catcagcaac tgtgacggat tggcgggcct catccgcaca 540
gacagcaccg acttcttcat tgagcctctg gagcggggcc agcaggagaa ggaggccagc 600
gggaggacac atgtggtgta ccgcccggag gccgtccagc aggagtgggc agaacctgac 660
ggggacctgc acaatgaagc ctttggcctg ggagaccttc ccaacctgct gggcctgggt 720
ggggaccagc tgggacgacac agagcgggaag cggcggcatg ccaagccagg cagctacagc 780
atcgaggtgc tgctgggtgt ggacgactcg gtggttcgct tccatggcaa ggagcatgtg 840
cagaactatg tctcaccct catgaatatc gtggtagatg agattacca cgatgagtcc 900
ctgggggttc atataaatat tgccctcgtc cgcttgatca tgggtggcta cgcacagcag 960
tccctgagcc tgatcgagcg cgggaacccc tcacgcagcc tggagcaggt gtgtcgctgg 1020
gcacactccc agcagcgcca ggaccccagc cacgctgagc accatgacca cgttgtgttc 1080
ctcaccggcg aggactttgg gccctcaggt gggtatgcac ccgtcactgg catgtgtcac 1140
cccctgagga gctgtgccct caacctagag gatggcttct cctcagcctt cgtgatagct 1200
catgagaccg gccacgtgct cggcatggag catgacggtc aggggaatgg ctgtgcagat 1260
gagaccagcc tgaggcagct catggcgccc ctggtgcagg ctgccttcca ccgcttccat 1320
tggctccgct gccgcaagct ggagctcagc cgtacctcc cgtcctacga ctgcctctc 1380
gatgacctct ttgatcctgc ctggccccag cccccagagc tgccctggat caactactca 1440
atggatgagc agtgccgctt tgactttggc agtggtacc agacctgctt ggcattcagg 1500
acctttgagc cctgcaagca gctgtggtgc agccatcctg acaaccgta cttctgcaag 1560
accaagaagg ggcctccgct ggatgggact gactgtgcac ccggcaagtg gtgcttcaa 1620
ggtcactgca tctggaagtc gccggagcag acatatggcg aggatggagg ctggagctcc 1680
tggaccaagt ttgggtcatg ttcgcggtca tbtgggggag ggggtgcgac ccgcagccgg 1740
agctgcaaca acccctcgcc agcctatgga ggcgcctgt gcttagggcc catgttcgag 1800
taccaggtct gcaacagcga ggagtgcctt cagaatgcca agcacagctg ggtgcctac 1860
tgtgccaagc gcaactcta ctatgtgcac cagatctgcc agtcggcgga cacgggggac 1920
gagcctgacg atgacgccc gaagtgtgag ctgactctgc gctgctgcc cagcggggac 1980
gtggtgttca tgaaccaggt ggttcacgat gggacacgct gcagctaccg ggaccatac 2040
agcgtctgtg cgcgtggcga gtgtgtgcct gtcggtgtg acaaggaggt ggggtccatg 2100
aaggcggatg acaagtgtgg agtctgcggg ggtgacaact cccactgcag gactgtgaag 2160
gggacgctgg gcaaggcctc caagcaggca ggtgctctca agctggtgca gatccagca 2220
ggtgccaggc acatccagat tgaggcactg gagaagtccc cccaccgcat tgtggtgaag 2280
aaccaggtca ccggcagctt catcctcaac cccaagggca aggaagccac aagccggacc 2340
ttcaccccca tgggcctgga gtgggaggat gcggtggagg atgccaagga aagcctcaag 2400
accagcgggc ccctgcctga agccattgcc atcctggctc tcccccaac tgagggtggc 2460
ccccgcagca gcctggccta caagtacgtc atccatgagg acctgctgcc ccttatcggg 2520
agcaacaatg tgctcctgga aggatggac acctatgagt gggcgctcaa gagctgggac 2580
ccctgcagca aggcctgtgg aggaggtatc cagtccacca aatacggctg ccggcgacga 2640
cgagaccacc acatggtgca gcgacacctg tgtgaccaca agaaggaggc caagccatc 2700
cgccggcgct gcaaccagca cccgtgctct cagcctgtgt ggggtgacgga ggagtggggt 2760
gcctgcagcc ggagctgtgg gtgcagacac ggtcagacac gggggataca gtgcctgctg 2820
cccctctcca atggaaccca caaggtcatg ccggccaaag cctgcgccgg ggaccggcct 2880
gaggcccgac ggcctgtct tggagagggc atccagcagc agtggaggct gggagcctgg 2940
tcccagtgtc ctgccacctg ttgcgagggg gataggccag ggcagggtgt gtgcaggacc 3000
aacgccaaca gcctcgggca gtggagcgga acactgtcca ggtctgcagc 3060
ctgcccgct gtggagcgga gccctgcacg ggagacaggt ctgtctctct ccagatggaa 3120
gtgctcgatc gctactgtc catccccggc taccaccggc tctgtgtgt gtctcgatc 3180
aagaaggcct cctggccca gccctgggac caacctcact gccccttc 3240
tccactcctg gaagccctt accaggacc caggacctg cagatgctgc agagcctcct 3300

```

ggaaagccaa	cgggatcaga	ggaccatcag	catggccgag	ccacacagct	cccaggagct	3360
ctggatacaa	gctccccagg	gacccagcat	ccctttgccc	ctgagacacc	aatccctgga	3420
gcatcctgga	gcatctcccc	taccaccccc	ggggggctgc	cttggggctg	gactcagaca	3480
cctacgccag	tccctgagga	caaagggcaa	cctggagaag	acctgagaca	tcccgccacc	3540
agcctccctg	ctgcctcccc	ggtgaca				3567

<210> 32

<211> 2930

<212> DNA

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 7473089CB1

<400> 32

cacgcagacc	gcggcagcgg	ccgagagccc	ggcccagccc	cttcccacag	cgccggcgttg	60
cgctgcccgg	cgccatgctt	ctgctgggca	tccaaacct	ggctttcgcc	ggcggaaccg	120
ctggaggctc	tgagccagag	cgggaggtag	tcgttcccat	ccgactggac	ccggacatta	180
acggcccgccg	ctactactgg	cggggtcccc	aggactccgg	ggatcaggga	ctcatttttc	240
agatcacagc	atttcaggag	gacttttacc	tacacctgac	gccggatgct	cagttctctg	300
ctcccgccctt	ctccactgag	catctgggcg	tccccctcca	ggggctcacc	gggggctctt	360
cagacctgcg	acgtgcttc	tattctgggg	acgtgaacgc	cgagccggac	tcgttcgctg	420
ctgtgagcct	gtgcgggggg	ctccgcggag	ccctttggcta	ccgaggcgcc	gagtatgtca	480
ttagcccgct	gcccaatgct	agcgcgccgg	cggcgccagc	caacagccag	ggcgccacacc	540
ttctccagcg	ccggggtggt	ccgggcgggc	cttccggaga	ccccacctct	cgctgcgggg	600
tggcctcggg	ctggaaacccc	gccatccctac	gggccttgga	cccttacaag	ccgcggcggg	660
cgggcttcgg	ggagagtctg	agccggcgca	ggctctggcg	cgccaagcgt	ttcgtgtcta	720
tcccgcggta	cgtggagacg	ctggtggtcg	cggacgagtc	aatggtcaag	ttccacggcg	780
cggacctgga	acattatctg	ctgacgctgc	tggcaacggc	ggcgcgactc	taccgccatc	840
ccagcatcct	caaccccatc	aacatcggtg	tggccaaggt	gctgcttctt	agagatcggt	900
actccggggcc	caaggtcacc	ggcaatgcgg	ccctgacgct	gcgcaacttc	tgtgcctggc	960
agaagaagct	gaacaaagtg	agtgaacagc	accccgagta	ctggggacact	gccatcctct	1020
tcaccaggca	ggacctgtgt	ggagccacca	cctgtgacac	cttgggcatg	gctgatgtgg	1080
gtaccatgtg	tgaccccaag	agaagctgct	ctgtcattga	ggacgatggg	cttccatcag	1140
ccttcaccac	tgccacagag	ctgggccaag	tggtcaacat	gccccatgac	aatgtgaaag	1200
totgtgagga	ggtgtttggg	aagctccgag	ccaaccacat	gatgtccccg	accctcatcc	1260
agatcgaccg	tgccaacccc	tggtcagcct	ccagtgtctg	catcatcacc	gacttccctg	1320
acagcgggca	cggtgactgc	ctcctggacc	aaccagcaa	gccccatctc	ctgcccagg	1380
atctgcccgg	cgccagctac	accctgagcc	agcagtgcga	gctggctttt	ggcggtgggt	1440
ccaagccctg	cgccctgcca	cagtactgca	ccaagctgtg	gtgcaccggg	aaggccaagg	1500
gacagatggt	gtgccagacc	cgccacttcc	cctgggcccga	tgccaccagc	tgtggcgagg	1560
gcaagctctg	cctcaaaggg	gcctgcgtgg	agagacacaa	cctcaacaag	cacagggtgg	1620
atggttccct	ggccaaatgg	gatccctatg	gccccctgct	gcgcacatgt	ggtggggggc	1680
tgcagctggc	caggaggcag	tgacccaacc	ccaccctgct	caacgggggg	aagtactcg	1740
agggagtgag	ggtgaaatac	cgatcctgca	atctggagcc	ctgccccage	tcagcctccg	1800
gaaagagctt	ccgggaggag	cagtgtgagg	ctttcaacgg	ctacaaccac	agcaccaacc	1860
ggctcactct	cgccgtggca	tgggtgcccc	agtaactccg	cgtgtctccc	cgggacaagt	1920
gcaagctcat	ctgccgagcc	aatggcactg	gctacttcta	tgtgctggca	cccaagggtg	1980
tgggtggacg	cacgctgtgc	tctcctgact	ccacctccgt	ctgtgtccaa	ggcaagtgca	2040
tcaaggctgg	ctgtgatggg	aacctgggct	ccaagaagag	attcgacaag	tgtgggggtg	2100
gtggggggaga	caataagagc	tgcaagaagg	tgactggact	cttcaccaag	cccatgcatg	2160
gctacaattt	cgtggtggcc	atccccgcag	gcgcctcaag	catcgacatc	cgcagcgcg	2220
gttacaagag	gctgatccgg	gatgacaact	acctggctct	gaagaacagc	caaggcaagt	2280
acctgctcaa	cgggcatttc	tggtgttcgg	cgggtggagc	ggacctgggt	gtgaaggcca	2340
gtctgctgcg	gtacagcggc	acgggcacag	cgggtggagag	cctgcaggct	tcccggccca	2400
tcctggagcc	gctgaccgtg	gaggtcctct	ccgtggggaa	gatgacaccg	ccccgggtcc	2460
gctactcctt	ctatctgccc	aaagagcctc	gggaggacaa	gtcctctcat	ccccgcacc	2520
cccggggagg	accctctgtc	ttgcacaaca	gcgtcctcag	cctctccaac	caggtggagc	2580
agccggacga	caggccccct	gcacgctggg	tggtggcag	ctggggggccg	tgctcccgca	2640
gctgcggcag	tggcctgcag	aagcggggcg	tggactggcg	gggctccgcc	gggcagcgca	2700
cggctccctgc	ctgtgatgca	gccccctggc	ccgtggagac	acaagcctgc	ggggagccct	2760
gccccacctg	ggagctcagc	gcctgggtcac	cctgtcccaa	gagctgcggc	cggggatttc	2820
agaggcgctc	actcaagtgt	gtggggccacg	gaggccggct	gctggcccg	gaccagtgca	2880
acttgcaccg	caagccccag	gagctggact	tctgcgtcct	gaggccgtgc		2930

<210> 33

<211> 4230

<212> DNA

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 7604035CB1

<400> 33

agcgagggttg	cctggagaga	gcgcctgggc	gcagaagggt	taacggggcca	ccgggggctc	60
gcagagcagg	aggggtgctct	cggacgggtg	gtccccact	gcactcctga	acttggagga	120
cagggtcgcc	gcgagggacg	cagagagcac	cctccacgcc	cagatgcctg	cgtagttttt	180
gtgaccagtc	cgctcctgcc	tccccctggg	gcagtagagg	gggagcgatg	gagaactgga	240
ctggcaggcc	ctggctgtat	ctgctgctgc	ttctgtccct	ccctcagctc	tgettggatc	300
aggaggtgtt	gtccggacac	tctcttcaga	cacctacaga	ggagggccag	ggccccgaag	360
gtgtctgggg	accttgggtc	cagtgggcct	cttgtctcca	gcccgtgcgg	gtgggggtgc	420
agcgaggagc	ccggagcatgt	cagctcccta	cagtgcagct	ccacccagat	ctgccccctc	480
ctccccggcc	cccaagacat	ccagaagccc	tcctcccccg	gggccagggt	cccagacccc	540
agactttctc	agaaaccttc	cccttgtaca	ggacacagtc	tcgggggaagg	ggtggccccc	600
ttcagggtcc	cgcttcccac	ctaggggagag	aggagaccca	ggagattcga	gcggccagga	660
ggtcccggct	tcgagacccc	atcaagccag	gaatgttcgg	ttatgggaga	gtgccctttg	720
cattgtccat	gcaccgggaac	cgagggcacc	ctcggagccc	accagatctt	gagctgtccc	780
tgatctcttc	tagaggggaa	gagcctattc	cgccccctac	tccaagagca	gagccattct	840
ccgcaaacgg	cagcccccaa	actgagctcc	ctcccacaga	actgtctgtc	cacaccccat	900
ccccccaagc	agaacctcta	agccctgaaa	ctgtctagac	agaggtggcc	cccagaacca	960
ggcctgcccc	cctacggcat	caccccagag	cccaggcctc	tggcacagag	ccccctctac	1020
ccacgcactc	cttagggagaa	ggtggcttct	tcctgtcatc	ccctcagcca	cgaaggccaa	1080
gttcccaggg	ttgggccagt	ccccaggtag	cagggagacg	ccctgatcct	tttctctcgg	1140
tcctctgggg	ccgaggccag	cagggccaaag	ggccttgggg	aacggggggg	actcctcacg	1200
ggccccgcct	ggagcctgac	cctcagcacc	cgggcgccctg	gctgccccctg	ctgagcaacg	1260
gcccccatgc	cagctccctc	tggagcctct	ttgtctccag	tagccctatt	ccaagatgtt	1320
ctggggagag	tgaacagcta	agagcctgca	gccaaagccc	ctgccccctt	gagcagccag	1380
acccccgggc	cctgcagtgc	gcagccttta	actcccagga	attcatgggc	cagctgtatc	1440
agtggggagc	cttcaactgaa	gtccagggtc	cccagcgctg	tgaactgaac	tgccggcccc	1500
gtggcttccg	cttctatgtc	cgtcacactg	aaaaggtcca	ggatgggacc	ctgtgtcagc	1560
ctggagcccc	tgacatctgt	gtggctggac	gctgtctgag	ccccggctgt	gatgggatcc	1620
ttggctctcg	cagggtcctc	gatggctgtg	gagtcgttgg	gggtgatgat	tctacctgtc	1680
gccttgtttc	ggggaacctc	actgaccgag	ggggccccct	gggctatcag	aagatcttgt	1740
ggattccagc	gggagccttg	cggctccaga	ttgccagctc	ccggcctagc	tccaactacc	1800
tggcacttcg	tggccctggg	ggccggtcca	tcacaaatgg	gaactgggct	gtggatcccc	1860
ctgggtccta	ggggccgggc	gggaccgtct	ttcgatataa	ccgtccctcc	agggaggagg	1920
gcaaagggga	gagtctgtcg	gctgaaggcc	ccaccacca	gcctgtggat	gtctatatga	1980
tctttcagga	ggaaaaccca	ggcgtttttt	atcagtatgt	catctcttca	cctcctccaa	2040
tccttgagaa	ccccacccca	gagccccctg	tccccagctc	tcagccggag	attctgaggg	2100
tggagccccc	acttgtctcg	gcacccccgc	cagccccggc	cccaggcacc	ctccagcgtc	2160
aggtgcggat	ccccagatg	cccgccccgc	cccatcccag	gacacccctg	gggtctccag	2220
ctgcgtactg	gaaacgagtg	ggacactctg	catgctcagc	gtcctgcggg	aaaggtgtct	2280
ggcgccccat	tttctctctg	atctcccgctg	agtcgggaga	ggaactggat	gaacgcagct	2340
gtgcgcgggg	tgccaggccc	ccagcctccc	ctgaaccctg	ccacggcacc	ccatgcccc	2400
catactggga	ggctggcgag	tggacatcct	gcagccgctc	ctgtggcccc	ggcaccacgc	2460
accgccagct	gcagtgcctg	caggaatttg	gggggggtgg	ctcctcggtg	cccccgagc	2520
gctgtggaca	tctcccccg	cccaacatca	ccagtccttg	ccagctgcgc	ctctgtggcc	2580
attgggaagt	tggctctcct	tggagccagt	gtccctgtcg	gtgcggcccg	ggccagagaa	2640
gccggcaggt	tcgctgtgtt	gggaacaacg	gtgatgaagt	gagcgagcag	gagtggtcgt	2700
caggcccccc	acagcccccc	agcagagagg	cctgtgacat	ggggccctgt	actactgct	2760
ggttccacag	cgactggagc	tccaagtgtc	cagccgagtg	tgggacggga	atccagcggc	2820
gctctgtggt	ctgccttggg	agtggggcag	ccactcgggc	caggccaggg	ggaagcagga	2880
gcaggaaactg	ggcagagctg	tccaacagga	agccggcccc	ctgacatgcg	cgctgcagc	2940
ctggggccct	gtgagagaa	ttggcgctgg	tacacagggc	cctggggtga	gtgtctctcc	3000
gaatgtggct	ctggcacaca	gcgtagagac	atcatctgtg	tatccaaact	ggggacggag	3060
ttcaacgtga	cttctccgag	caactgttct	cacctcccca	ggccccctgc	cctgcagccc	3120
tgtcaagggc	aggcctggca	ggaccgatgg	atttccacgc	cctggagccc	atgttctcgc	3180
tcctgccaaag	ggggaacgca	gacacgggag	gtccagtgcc	tgagcaccaa	ccagaccctc	3240
agcaccgat	gcccctctca	actgcggccc	tccaggaagc	gccccgttaa	cagccaaccc	3300
tcagccagc	gccttgatga	tcaatgcaag	gacagctctc	cacattgccc	cctggtggta	3360
caggccccgc	tctgcgtcta	ccccactaac	acagccacct	gttgccgctc	ttgcgcacat	3420
gtcctggagc	ggtctcccca	ggatccctcc	tgaagggggt	ccggggcacc	ttcacggttt	3480

```

tctgtgccac catcggtcac ccattgatcg gccactctg aacccctgg ctctccagcc 3540
tgtcccagtc tcagcaggga tgtcctccag gtgacagagg gtggcaagg gactgacaca 3600
aagtgacttt caggcgtgtg gtcaggccca tgggtgggtg tgatgggtgt gtgcacatat 3660
gcctcagggtg tgcttttggg actgcatgga tatgtgtgtg ctcaaactgt tatcactttt 3720
caaaaagagg ttacacagac tgagaaggac aagacctgtt tccttgagac ttctctaggt 3780
ggaaaggaaa gcaagtctgc agttccttgc taatctgagc tacttagagt gtggtctccc 3840
caccaactcc agttttgtgc cctaagcctc atttctcatg ttcagacctc acatcttcta 3900
agccgcctcg tgtctctgac cccttctcat ttgcctagta tctctgcccc tgctcccta 3960
attagctagg gctggggtca gccactgcc aacctgcctt actcaggaag gcaggaggaa 4020
agagactgcc tctccagagc aaggcccagc tgggcagagg gtgaaaaaga gaaatgtgag 4080
catcgcctcc cccaccacc cgccagcccc ctgacccccc tccctgcctc ctgaaatggt 4140
tcccaccagc aactaattta ttttttatta aagatggtca tgacaaatga aaaaaaaaaa 4200
aaaaaaaaaa aaaaaacaaa aaaaaaataa

```

```

<210> 34
<211> 3699
<212> DNA
<213> Homo sapiens

```

```

<220>
<221> misc_feature
<223> Incyte ID No: 3473847CB1

```

```

<400> 34
cgcagtgtgc tggcaagct tgactttccc agcaggccta tgtcataggt actgtggtct 60
ctacaataca gcagagggtat ctgaggctcc gagagggtga gtgacttgct catggtctgca 120
caaccagtaa atattggagc ctgactgggg catgggggag ggggtggcct ttgggcaggg 180
attttttccc tcaatttatt ctgactgggg catgggggag ggggtggcct ttgggcaggg 240
ccaccaggag cgaccaggcc cgtagagagc tgggtgcagg tacagaggaa aacctgttgt 300
cgagtgtggc cgtagtctcc catttttgcc tgaatggcac atttgaaagt gttatataac 360
catgtgaata ataatagttg ccctatatga gttttttaat ttgctttttg gtcgcgcat 420
ggtaacttct ttatcatcta ctatactctg ttgtgtctct tttgttgtaa tttgtaagta 480
gggtgagat aaagtacacc tagggtttgc tgggtttctt ccatgtcatc atgttctctc 540
ttgcatgggg ccaggatccg tggaggttgc ctggcaccta cgtggtggtg ctgaaggagg 600
agaccacact ctgcagtcga gagcgcactg cccgcgcct gcaggcccag gctgcccgcc 660
ggggatacct caccaagatc ctgcatgtct tccatggcct tcttcttggc ttctgtgga 720
agatgagtgg cgacctgtcg gagctggcct tgaagtggc ccatgtcgac tacatcgagg 780
aggactcctc tgtctttgcc cagagcatcc cgtggaacct ggagcggatt accctccac 840
ggtaccgggc ggtatgaatac cagccccccg acggaggcag cctggtggag gtgtatctcc 900
tagacaccag catacagagt gaccaccggg aaatcgaggg cagggtcatg gtcaccgact 960
tcgagaatgt gcccaggagg gacgggaccc gcttccacag acaggccagc aagtgtgaca 1020
gtcatggcac ccacctggca ggggtggtca gcggccggga tgccggcgtg gccaaagggtg 1080
ccagcatgcg cagcctgcgc gtgctcaact gccaaaggaa gggcacggtt agcggcaccc 1140
tcataggcct ggaagtttatt cggaaaagcc agctggtcca gcctgtgggg ccactgtgtg 1200
tgetgtgccc ggtggcgggt ggtcgtgctg ggttacagcc cgtctctcaa cgccgcctgc cagcgcctgg 1260
cgagggtctg ggtcgtgctg gtcaccgctg cgggcaactt ccgggacgat gcctgcctct 1320
actccccagc ctcagctccc gaggtcatca cagttggggc caccaatgcc caggaccagc 1380
cgggtgaccct ggggactttg gggaccaact ttggccgctg tgtggacctc tttgccccag 1440
gggaggacat cattggtgcc tccagcgact gcagcacctg ctttgtgtca cagagtggga 1500
catcacaggg tgctgcccac gtggctggca ttgcagccat gatgctgtct gccgagccgg 1560
agctcaccct ggccgagttg aggcagagac tgatccactt ctctgccaaa gatgtcatca 1620
atgaggcctg gttccctgag gaccagcggg tactgacccc caacctggtg gccgccctgc 1680
ccccagcac ccatggggca ggttggcagc tgttttgtag gactgtgtgg tcagcacact 1740
cggggcctac acggatggcc acagccatcg cccgctgcgc ccagatgag gagctgctga 1800
gctgctccag ttctccagg agtgggaagc gggggggcga ggcagtgagg gcccaagggg 1860
gcaagctggt ctgcccggcc cacaacgctt ttgggggtga ggggtgtctac gccattgcca 1920
ggtgctgcct gctaccccag gccaaactgca gcgtccacac agctccacca gctgaggcca 1980
gcatggggac ccgtgtccac tgccaccaac agggccacgt cctcacaggg tgcagctccc 2040
actgggaggt ggagacactt ggcaccaca agccgctgt gctgaggcca cgaggtcagc 2100
ccaaccagtg cgtgggccac agggaggcca gcatccacgc ttctgtctgc catgccccag 2160
gtctggaatg caaagtcaag gagcatggaa tcccggcccc tcaggagcag gtgaccgtgg 2220
cctgcgagga gggctggacc ctgactggct gcagtgcctt cctgggacc tcccacgtcc 2280
tgggggccta gcctgtagac aacacgtgtg tagtcaggag ccgggacgtc agcactacag 2340
gcagcaccag cgaagaggcc gtgacagccc ttgccatctg ctgccggagc cggcacctgg 2400
cgcaggcctc ccaggagctc cagtgcagc cccatcccag gatgggtgtc tggggagggt 2460
caagggtctg ggctgagctt taaaatggtt cccctctcag cctccatgg 2520
cctggcacga ggggatgggg atgcttccgc ctttccgggg ctgctggcct ggccttgag 2580

```

tggggcagcc	tccttgccctg	gaactcactc	actctgggtg	cctcctcccc	aggtggaggt	2640
gccaggaagc	tccttccctc	actgtggggc	atttcacat	tcaaacaggt	cgagctgtgc	2700
tcgggtgctg	ccagctgtct	ccaatgtgcc	gatgtccgtg	ggcagaatga	ctttttattga	2760
gctcttgctt	cgtgccaggc	attcaatcct	caggtctcca	ccaaggaggc	aggattcttc	2820
ccatggatag	gggagggggc	ggtaggggct	gcagggacaa	acatcggttg	gggggtgagt	2880
tgaaggtgc	tgatggccct	catctccagc	taactgtgga	gaagccctg	ggggctccct	2940
gattaatgga	ggcttagctt	tctggatggc	atctagccag	aggctggaga	caggtgtgcc	3000
cctgggtggc	acaggtctgt	ccttggtttc	ctgagccacc	tttactctgc	tctatgccag	3060
gctgtgctag	caacacccaa	aggtggcctg	cggggagcca	tcacctagga	ctgactcggc	3120
agtgtgcagt	ggtgcatgca	ctgtctcagc	caaccgcctc	cactaccggg	caggggtacac	3180
attcgacccc	ctacttcaca	gaggaagaaa	cctggaacca	gagggggcgt	gcctgccaa	3240
ctcacacagc	aggaactgag	ccagaaacgc	agattgggct	ggctctgaag	ccaagcctct	3300
tcttacttca	cccggtctgg	ctcctcattt	ttacgggtaa	cagtgaaggc	gggaagggga	3360
acacagacca	ggaagctcgg	tgagtgatgg	agaacgatg	cctgcaggca	tggaaactttt	3420
tcogttatca	cccaggcctg	attcactggc	ctggcggaga	tgcttctaag	gcatggctcg	3480
gggagagggc	caacaactgt	ccctccttga	gcaccagccc	cacccaagca	agcagacatt	3540
tatcttttgg	gtctgtccct	tctgttgcct	ttttacagcc	aacttttcta	gacctgtttt	3600
gcttttgtaa	cttgaagata	tttattctgg	gttttgtagc	atattttatta	atatggtgac	3660
tttttaaaat	aaaaacaaac	aaacgttgtc	ctaaaaaaa			3699

<210> 35

<211> 2410

<212> DNA

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 3750004CB1

<400> 35

cctcagcagt	ggcccccttc	ctccacgggc	tgccccggag	ctcagtccca	ccccctccgc	60
cgatgaggcc	atgaggcacc	gaacggacct	gggccagAAC	ctcctgtctt	tctgtggggc	120
cctgtgaac	tgtggtttgg	gggtcagtgc	tcagggtccg	ggcgagtgga	ccccgtgggt	180
gtcctggacc	cgctgtccca	gctcctgcgg	gcgtggcgct	tctgtgcgca	gcggcgctgt	240
cctccggctt	cctggggaag	aaccgtgtct	gggagactcc	catgagtacc	gcctctgccA	300
gtttgccagac	tgccccccag	gggtgtgtcc	cttccgagac	ctacagtgtg	ccctgtacaa	360
tgcccgccct	gtcctgggca	cccagaagac	ctaccagtgg	gtgccccctc	atggggcgcc	420
caaccagtgc	gacctcaact	gcctgggtga	ggggcacgcc	ttctaccaca	gcttcggccg	480
cgtcctggac	ggcacccgct	gcagcccggg	tgcccagggg	gtctgcgtgg	ctggccgctg	540
ccttagcgcc	ggctgtgatg	ggttgtttgg	ctcgggtgcc	ctcgaggacc	gctgtggccg	600
ctgcgggagg	gccaacgact	cgtgcctttt	cgtgcagcgc	gtgtttctgt	acgcgggtgc	660
cttcgctggg	tactggaacg	tgacctgat	ccccgagggc	gccagacaca	tccgcgtgga	720
acacaggagc	cgcaaccacc	tgggtatcct	aggatcactg	atggggggcg	atggggcgcta	780
cgtgtctaat	gggcactggg	tggtcagccc	accagggacc	tacgaggcgg	ccggcacgca	840
tgtggtctac	acccgagaca	cagggcccca	ggagacattg	caagcagccg	ggccccctc	900
ccatgacctg	ctcctacagg	tcctcctgca	ggagcccaac	cctggcatcg	agttttgagtt	960
ctggctccct	cgggagcgct	acagccccct	ccaggctcgt	gtgcaggccc	tgggctggcc	1020
cctgaggcag	cctcagcccc	ggggggtgga	cgtcagccc	cccgcagccc	ctgctgtcac	1080
ccctgcacag	accccaacgc	tgcccccaga	ccccctgcca	ccctgccctg	acaccccgcg	1140
ccgcgcccac	cgactactcc	actattgcgg	cagtgaactt	gtgttccagg	cccgagtgtc	1200
gggccaccac	caccaggccc	aggagaccgc	ctatgaggtg	cgcatccagc	tcgtctacaa	1260
gaaccgctcg	ccactgcggg	cacgcgagta	cgtgtgggcg	ccaggecact	gccccctgcc	1320
gatgctggca	ccccaccggg	actacctgat	ggctgtccag	cgtcttgtca	gccccgacgg	1380
cacacaggac	cagctgtctg	tgccccacgc	cggctacgcc	cggccctgga	gcctgtcgga	1440
ggacagccgc	ataccgctga	ctgcccggcg	ctgtcctggc	tgagccctgt	caggagcccc	1500
ggccacacac	agcaagaaag	atacatctga	ccagcctcaa	cgtcaacgta	tttccccctc	1560
caccctggct	tccaggcgagc	tctgaaatac	gtcccacctg	tgcagctatg	tgactccctc	1620
ccacacacgc	ttaagacacc	tctgcatgca	gtcaaagcca	ctgtcacaa	ccggcaggca	1680
ctggtgagga	ggcactaagg	agactctgac	ttttatttcg	cctctctcct	tggctgccag	1740
gaagctcata	gctattttata	ctcagaaagt	ttaacgctgc	tttctttctc	tttgccgcgg	1800
tcacacttgc	ttggagacac	tgatcatgaa	gagcatgaca	ccctgctgcc	ctgggtaccc	1860
agaagatcat	ctgtttactt	cccagacact	gtgctgtctc	tgctctctgc	tactcacaca	1920
cacctcatg	tgtgaagggc	agagacactg	tcacaaacag	gcattgcccc	tagaagacat	1980
gcctaaccag	gcactgtaac	gtaccaacgt	accaatttcc	ccttttcccc	tggctaccag	2040
gaaactcgga	gacaatcttt	tcagcctcag	catttctggc	tggatttcca	cccatcaaca	2100
cgtgcttctg	cctccttttt	ttttttctct	aggtagacct	tgctctgtca	cctaggtgta	2160
agtgcgggtg	tgcaatcatg	gctcactgca	gcctcaaatt	cctgggctca	agcgatcctc	2220

ccactcagct	ccacagcagct	ggaactcacg	tgtgatcaca	tgccggctaa	tttaaatttt	2280
gtagagatgg	gcttgtagct	gccaaatgtc	tcactatggc	tcaacatctc	tgctgggtcc	2340
aagactgaat	aggatgacat	gatgggtgtac	cccttatcct	tatttcagct	ttaaaaatc	2400
taaaaaaaaa						2410

<210> 36
 <211> 549
 <212> DNA
 <213> Homo sapiens

<220>
 <221> misc_feature
 <223> Incyte ID No: 4904126CB1

<400> 36	gggaggagag	aaaagccatg	gccgacaagg	tcctgaagga	gaagagaaa	cagtttatcc	60
	gttcagtgagg	cgaagggtaca	ataaatggct	tactgggtga	attattggag	acaagggtgc	120
	tgagccagga	agagatagag	atagtaaaat	gtgaaaatgc	tacagttatg	gataaggccc	180
	gagctttgct	tgactctgtt	attcggaaag	gggctccagc	atgccaaatt	tgcatcacat	240
	acatttgtga	agaagacagt	cacctggcag	ggacgctggg	actctcagca	ggccaacat	300
	ctggaaatca	ccttactaca	caagattctc	aaatagtact	tccttcctag	gtaatgctgt	360
	ttttaaagaa	agagcattct	ttgaaccgtg	gcttcccgtg	acattaatgt	tgtaggatga	420
	accacagtta	aaggggctat	gaagaattcc	catagagtga	tcatacaatt	ttctttttgt	480
	aatctattct	gctttttag	caactgtcaa	aacagcttca	ctatctatgt	ctacattaaa	540
	atttggaaat						549

<210> 37
 <211> 2755
 <212> DNA
 <213> Homo sapiens

<220>
 <221> misc_feature
 <223> Incyte ID No: 71268415CB1

<400> 37	ttgctaggag	ggtggagttc	atccacttat	gatataaatg	tctcttttta	ttttttgcag	60
	gcaacttttt	gctccttcct	acacagaaac	ccattatact	tcaagtggta	accctcaaac	120
	caccacacgg	aaattggagg	atcactgctt	ttaccacggc	acgggtgagg	agacagaact	180
	gtccagcgct	acgctcagca	cttgccgagg	aattagagga	ctgattacgg	tgagcagcaa	240
	cctcagctac	gtcatcgagc	ccctccctga	cagcaagggc	caacacctta	tttacagatc	300
	tgaacatct	aagccgcccc	cgggaaactg	tgggttcgag	cactccaagc	ccaccaccag	360
	ggactgggct	cttcagttta	cacaacagac	caagaagcga	cctcgagga	tgaaaaggga	420
	agatttaaac	tccatgaagt	atgtggagct	ttacctcggt	gctgattatt	tagagtttca	480
	gaagaatcga	cgagaccagg	acgccaccaa	acacaagctc	atagagatcg	ccaactatgt	540
	tgataagttt	taccgatcct	tgaacatccg	gattgtctct	gtgggcttgg	aagtgtggac	600
	ccacgggaac	atgtgtgaag	tttcagagaa	tccatattct	accctctggt	cctttctcag	660
	ttggaggcgc	aagctgcttg	cccagaagta	ccatgacaa	gcccatttaa	tcacgggcat	720
	gtccttcac	ggcaccacca	tcggcctggc	ccccctcatg	gccatgtgct	ctgtgtacca	780
	gtctggagga	gtcaacatgg	accactccga	gaatgccatt	ggcgtggctg	ccaccatggc	840
	ccacgagatg	ggccacaact	ttggcatgac	ccatgattct	gcagattgct	gctcggccag	900
	tgcggtgat	ggtgggtgca	tcattggcagc	tgccactggg	cacccctttc	ccaaagtgtt	960
	caatggatgc	aacaggaggg	agctggacag	gtatctgcag	tcagggtggtg	gaatgtgtct	1020
	ctccaacatg	ccagacacca	ggatgttgta	tggaggccgg	aggtgtggga	acgggtatct	1080
	ggaagatggg	gaagagtgtg	actgtggaga	agaagaggaa	tgtaacaacc	cctgctgcaa	1140
	tgctctaat	tgtaacctga	ggccgggggc	ggagtgtgct	cacggctcct	gctgccacca	1200
	gtgtaagctg	ttggctcctg	ggaccctgtg	ccgcgagcag	gccaggcagt	gtgacctccc	1260
	ggagttctgt	acggggcaagt	ctccccactg	ccctaccaac	ttctaccaga	tggtatggtac	1320
	cccctgtgag	ggcggccagg	cctactgcta	caacggcatg	tgccctcacct	accaggagca	1380
	gtgccagcag	ctgtggggac	ccggagcccc	acctgcccc	gacctctgct	tcgagaagggt	1440
	gaatgtggca	ggagacacct	ttggaaactg	tggaaaaggac	atgaatgggtg	aacacaggaa	1500
	gtgcaacatg	agagatgcga	agtgtgggaa	gatccagtgt	cagagctctg	aggccccggc	1560
	cctggagtcc	aacgcggtgc	ccattgacac	cactatcatc	atgaatggga	ggcagatcca	1620
	gtgccggggc	acccacgtct	accgaggtcc	tgaggaggag	ggtgacatgc	tggaaccagg	1680
	gctggtgatg	actggaacca	agtgtggcta	caaccataat	tgctttgagg	ggcagtgcat	1740
	gaacacctcc	ttctttgaaa	ctgaaggctg	tgggaagaag	tgcaatggcc	atggggctctg	1800
	taacaacaac	cagaactgcc	actgctgcc	gggctgggcc	cgcctctct	gcaacacacc	1860

ggggcacggg	ggcagtatcg	acagtggggc	tatgccccct	gagagtgtgg	gtcctgtggg	1920
agctggagtg	ttggtggcca	tcttgggtgt	ggcggtccct	atgctgatgt	actactgctg	1980
cagacagaa	aacaaactag	gccaactcaa	gccctcagct	ctcccttcca	agctgaggca	2040
acagttcagt	tgtcccttca	gggttttctc	gaacagcggg	actggatcat	ccaacccaac	2100
tttcaagctg	cagacgcccc	agggcaagcg	aaaggtgatc	aacactccgg	aaatcctgcg	2160
gaagccctcc	cagcctcctc	cccggccccc	tccagattat	ctgcgtgggt	gggtccccacc	2220
tgaccactg	ccagctcacc	tgagcagggc	tgctaggaac	tccccagggc	ccgggtctca	2280
aatagagagg	acggagtcgt	ccaggaggcc	tctccaagc	cggccaattc	cccccgacc	2340
aaattgcatc	gtttcccgag	acttctccag	gcctcgcccg	ccccagaagg	cactcccgcc	2400
aaaccagtg	ccaggccgca	ggagcctccc	caggccagga	ggtgcatccc	cactgcccgc	2460
ccctggtgct	ggccctcagc	agtcggggcc	tctggcagca	cttgcccca	aggtgagtc	2520
acgggaagcc	ctcaaggtga	aagctgggtac	cagaggggctc	caggggggca	ggtgtagagt	2580
tgagaaaaca	aagcaattca	tgcttcttgt	ggtctggact	gaacttcag	aacaaaagcc	2640
aagggcaaaa	cattcatgtt	tcttgggtgc	cgcttgactg	tgaggtttg	gcttcattgt	2700
aaaggtgatt	cttagaatcc	tgagctgtgg	tggtctcagt	cctgccccct	cacct	2755

<210> 38
 <211> 2553
 <212> DNA
 <213> Homo sapiens

<220>
 <221> misc_feature
 <223> Incyte ID No: 7473301CB1

<400> 38						
atggacaaag	aaaacagcga	tgtttcagcc	gcacctgctg	acctgaaaat	atccaatatc	60
tcagttccaag	tggtcagtg	ccaaaagaag	ctgccagtga	gacgaccacc	gttgccaggg	120
agacgactac	cattgccagg	aagacgacca	ccacaaagac	ccattggcaa	agccaaaccc	180
aagaagcaat	ccaagaaaaa	agttcccttt	tggaaatgtac	aaaaataaaat	cattctcttc	240
acagtatttt	tattcatcct	agcagtcata	gcctggacac	ttctgtgggt	gtatatcagt	300
aagacagaaa	gcaaagatgc	tttttacttt	gctgggatgt	ttcgcatcac	caacatcgag	360
tttcttcccg	aataccgaca	aaaggagtc	agggaaatttc	tttcagtgtc	acggactgtg	420
cagcaagtga	taaacctggg	ttatacaaca	tctgcctctc	ccaaatttta	tgagcagtc	480
gttgttgca	atgtcagcag	caacaacaaa	ggcgccctcc	ttgtccactt	ttggattgtt	540
tttgtcatgc	cacgtgccaa	aggccacatc	ttctgtgaag	actgtgttgc	cgccatcttg	600
aaggactcca	tccagacaag	catcataaac	cggacctctg	tggggagctt	gcagggactg	660
gctgtggaca	tggaactctg	ggtaactaaat	ggtgattggt	ggtcattcct	aaaaaaaaag	720
aaaagaaaag	aaaatgggtg	tgtctccaca	gacaaaggct	gctctcagta	cttctatgca	780
gagcatctgt	ctctccacta	cccgtctggg	atctctgcag	cctcagggag	gctgatgtgt	840
cacttcaagc	tggtggccat	agtgggctac	ctgattcgtc	tctcaatcaa	gtccatccaa	900
atcgaagccg	actgaactgc	cactgactcc	ctgaccattt	acgactccct	tttggccatc	960
cggagcagca	tcttgtacag	aatttgtgaa	cccacaagaa	cattaatgtc	atttgtttct	1020
acaaataatc	tcattgttgt	gacatttaag	tctcctcata	tacggaggct	ctcaggaatc	1080
cgggcatatt	ttgaggtcat	tccagaacaa	aagtgtgaaa	acacagtgtt	ggtcaaaagc	1140
atcactggct	ttgaagggaa	aatttcaagc	ccatattacc	cgagctacta	tcttccaaaa	1200
tgcaagtgtg	cctggaaatt	tcagacttct	ctatcaactc	ttggcatagc	actgaaattc	1260
tataactatt	caataaccaa	gaagagtatg	aaaggctgtg	agcatggatg	gtgggaaatt	1320
tatgagcaca	tgtactgtgg	ctcctacatg	gatcatcaga	caatttttct	agtggccagc	1380
cctctgggtc	acattcagct	ccagtgcagt	tcaaggcttt	caggcaagcc	acttttggca	1440
gaatatggca	gttacaacat	cagtcaaccc	tgccctgtgg	gatcttttag	atgctcctcc	1500
ggtttatgtg	tccctcaggc	ccagcgtggg	gatggagtaa	atgactgctt	tgatgaaagt	1560
gatgaactgt	tttgctgag	ccctcaacct	gcttgcata	ccagctcctt	caggcagcat	1620
ggccctctca	tctgtgatgg	cttcagggac	tgtagaatg	gccgggatga	gcaaaaactgc	1680
actcaaagta	ttccatgcaa	caacagaact	tttaagtgtg	gcaatgatat	ttgctttagg	1740
aaacaaaatg	caaaatgtga	tgggacagtg	gattgtccag	atggaaagtga	tgaagaaggc	1800
tgcacctgca	gcaggagtgc	ctccgccctt	caccgcatca	tccgaggcac	agacacccctg	1860
gagggggggt	ggccgtggca	ggtcagcctc	cactttgttg	gatctgccta	ctgtggtgcc	1920
tcagtcattc	ccagggtgag	gcttctttct	gcagccact	gttttcatgg	aaacaggctg	1980
tcagatccca	caccatggac	tgcacacctc	gggatgtatg	ttcaggggaa	tgccaaagttt	2040
gtctccccgg	tgagaagaat	tgtgtgtccac	gagtactata	acagtacagc	ttttgattat	2100
gatattgctt	tgctacagct	cagtattgcc	tggtctgaga	ccctgaaaca	gctcattcag	2160
ccaatatgca	ttctctccac	tggtcagaga	gttcgcagtg	gggagaagtg	ctgggttaact	2220
ggctgggggc	gaagacacga	agcagataat	aaaggctccc	tctgtctgca	gcaagcggag	2280
gtagagctca	ttgatcaaac	gctctgtgtt	tccacctacg	ggatcatcac	ttctcggatg	2340
ctctgtgacg	gcataatgtc	aggcaagaga	gatgcctgca	aaggagattc	gggtggacct	2400
ttatcttgct	gaagaaaaag	tgatggaaaa	tggattttga	ctggcattgt	tagctgggga	2460

catggatgtg gacgacaaa ctttcctgggt gtttacacaa ggggtgtcaaa ctttgttccc 2520
 tggattcata aatatgtccc ttctcttttg taa 2553

<210> 39
 <211> 1041
 <212> DNA
 <213> Homo sapiens

<220>
 <221> misc_feature
 <223> Incyte ID No: 7473308CB1

<400> 39
 atgttcagcg gcaacacagg aaaaacccat attatcaatg ctcaaaaacc tggccacctc 60
 aggcttagcc agttattcgt gagcagagag gtgtgtcatc tacatggcag tcatggcctg 120
 gatgggtctg gaactgtggc aagaatcctt ccaggaaaca gccggtctcc ctctctgctc 180
 tcagaaggca agtttcctta tcacctgtct gctctcagaa ggcaagtttc cttatcacct 240
 gtgaatcaca aaccacaga gtggccaaac atactgatgc aagaccatag gaaggggaaa 300
 gctgcagttg gtgtctcctt tgatgatgat gacaagattg ttgggggcta caactgtgag 360
 gagaattctg tccccacca ggtgtccctg aattctggct accacttctg tgttggctcc 420
 ctcaacaggg aatactgcat ccagggtgaga ctgggagagc acaacatcga agtcctagag 480
 gggaatgaac agttcatcta tgcggtcaag atcatccgcc accccaaata caacagctgg 540
 actctggaca atgacatcct gctgatcaag ctctccacac ctgccatcat caatgcccat 600
 gtgtccacca tctcttgcc caccacccct ccagctgctg gcactgagtg cctcatctct 660
 ggctggggca cactctgag ttctggcgcc gactaccagc acgagctgca gtgcctggat 720
 gctcctgtgc tgagccaggc tgagtatgaa gcctctacc ctggaaagat taccaacaac 780
 gtgttttctg tgggtttcct tgagggaggc aaggattcct gccagattat tcctatcaaa 840
 gtgcagcagc tggttacctc aagccaagag acagacataa ggatccctat ggccttgag 900
 acagctgctt ccacctcta cctgggcccc ttagactctt tacacaggaa agtgagtcac 960
 cccactgaga agcgttgcca gcagaaacag ggcatgaaaa tcacagataa ccatgggatt 1020
 acttccaagt ggtcagtata a 1041

<210> 40
 <211> 1707
 <212> DNA
 <213> Homo sapiens

<220>
 <221> misc_feature
 <223> Incyte ID No: 7478021CB1

<400> 40
 atgtctgcgcg cctccatctt ccgtccgaca ctgtctctct gctggctggc tgctccctgg 60
 cccaccacgc ccgagagtct ctccacagc cgggaccgct cggacctgga gccgtcccca 120
 ctgcgccagg ccaagcccat tgccgacctc cacgctgctc ageggttcct gtccagatac 180
 ggctgggtcag ggggtgtggc ggccctggggg ccaggtcccg aggggcccgc ggagaccccc 240
 aaggggcgcg ccctggccga ggcggtgcgc aggttccagc gggcgaaacgc gctgccggcc 300
 agcggggagc tggacgcggc caccctagcg gccatgaacc ggccgcgctg cgggggtccc 360
 gacatgcgcc caccgcccc ctccgcccc ccttcgcccc cgggccccgc cccagagcc 420
 cgctccaggc gctccccgcg ggcgcgctg tccttgctcc ggcggggttg gcagccccgc 480
 ggctaccccg acggcggagc tgcccaggcc ttctccaaga ggacgctgag ctggcggctg 540
 ctgggcgagg ccctgagcag ccaactgtcc gtggccgacc ageggcgcat agaggcctg 600
 gccttcagga tgtggagcga ggtgacgcg ctggacttcc gcgaggacct ggccgcccc 660
 ggggcccgcg tcgacatcaa gctgggcttt gggagacggc acctgggctg tccgcgggcc 720
 ttcgatggga gcgggcagga gtttgacac gcctggcgcc taggtgacat tcaatttgac 780
 gacgacgagc acttcacacc tcccaccagt gacacgggca tcagccttct caagggtggc 840
 gtccatgaaa ttggccatgt cctgggcttg cctcacacct acaggacggg atccataatg 900
 caaccaaatt acattcccga ggagcctgcc tttagattgg actggtcaga caggaaagca 960
 attcaaaagc tgatatggtc ctgtgaggga tcatttgata ctgcgtttga ctggattcgc 1020
 aaagagagaa accaatatgg agaggtgatg gtgagattta gcacatatct cttccgtaac 1080
 agctggtact ggctttatga aaatcgaaac aataggacac gctatgggga ccctatccaa 1140
 atcctcactg gctggcctgg aatcccaaca cacaacatag atgcctttgt tcacatctgg 1200
 acatggaaaa gagatgaacg ttattttttt caaggaaatc aatactggag atatgacagt 1260
 gacaaggatc aggccctcac agaagatgaa caaggaaaaa gctatcccaa attgatttca 1320
 gaaggatttc ctggcatccc aagtccccca gacacggcgt tttatgaccg aagacagaag 1380
 ttaatttact tcttcaagga gtcccttgta ttgtcatttg atgtcaacag aaatcgagta 1440
 cttaattctt atccaaagag gattactgaa gttttccag cagtaatacc acaaaatcat 1500

```

cctttcagaa atatagattc cgcttattac tcctatgcat acaactccat tttctttttc 1560
aaaggcaatg catactggaa ggtagttaat gacaaggaca aacaacagaa ttcttggett 1620
cctgctaatt gcttatttcc aaaaagttt atttcagaga agtgggttga tgtttgtgac 1680
gtccatatct ccacactgaa catgttaa 1707

```

```

<210> 41
<211> 1262
<212> DNA
<213> Homo sapiens

```

```

<220>
<221> misc_feature
<223> Incyte ID No: 4333459CB1

```

```

<400> 41
aaaagatctt tgcgaaacac tacattcaga aacatcagat ggacatgctt gattcaccac 60
gtcttgggta atgaataaac ttgtttttaa ttggcttatt gctggctctc caaggcttcc 120
tatttttggt tgcttttagtc tctctaaaat ttcagggaaa aactatgagt ctcaaaatgc 180
ttataagcag gaacaagctg attttactac taggaatagt cttttttgaa cgaggtaaatt 240
ctgcaactct ttcgctcccc aaagctcccc gttgtgggca gagtctggtt aagggtacagc 300
cttgggaatta tttaacatt ttcatgtcga ttcttgagg aagccaagtg gagaagggtt 360
cctatccctg gcagggtatct ctgaaacaaa ggcagaagca tatttgtgga ggaagcatcg 420
tctcaccaca gtgggtgatc acggcggtc actgcattgc aaacagaaac attgtgtcta 480
ctttgaatgt tactgctgga gagtatgact taagccagac agaccagga gagcaaaact 540
tcaatttga aactgtcatc atacatccac atttctcac caagaaacca atggactatg 600
atattgccct tttaagatg gctggagcct tccaatttgg ccactttgtg gggcccatat 660
gtcttccaga gctgcgggag caatttgagg ctggttttat ttgtacaact gcaggctggg 720
gccgcttaac tgaaggtggc gtcctctcac aagtcttgca ggaagtgaat ctgcctattt 780
tgacctggga agagtgtgtg gcagctctgt taacactaaa gaggcccatc agtgggaaga 840
cctttctttg cacaggtttt cctgatggag ggagagacgc atgtcaggga gattcaggag 900
gttcaactcat gtgccggaat aagaaagggg cctggactct ggctgggtgt acttctctgg 960
gtttgggctg ttgtcgaggc tggagaaaca atgtgaggaa aagtgatcaa ggatcccctg 1020
ggatcttcac agacattagt aaagtgcctt cctggatcca cgaacacatc caaactggta 1080
actaagccat cacacaaggt taagaagctg ccattctgct agggccagag acagcatcag 1140
cagagtctct gcaaatcaga gcacctgaac caacaggctc tacctctgtt ctcagtgtag 1200
cacacaagga ttgtgagggt taccaagtct aaataaaaca agagttaaat atggtaaaaa 1260
aa 1262

```

```

<210> 42
<211> 3067
<212> DNA
<213> Homo sapiens

```

```

<220>
<221> misc_feature
<223> Incyte ID No: 6817347CB1

```

```

<400> 42
gcactgtgaa cggttggtgc atccaaatct gaattttgtc tgggaccagg gtcagggacc 60
agaatacacc agagctgagg gccagcccta cctgagaacc atcaacaaac ttaccccaca 120
tcccattata cctcctcact ccctgcagcc tgctcagcttc cccaatctcc cacactcact 180
gtcacctggg gctctggtgc accagatgac actacttget cctgggtaca caggcccat 240
gatecccatg gatgttaatg agcccagctc cgtgaccagc gctcctaccc tcagctctag 300
cctgcagcat atctcctcat tcctggccac tggtaagaaa ctttccctcc atttttggtca 360
tccacgtgag tgtgaagtca ccaggattga tgacaaaaat agaagaggat tggagacag 420
tgagccagggt gccaaactct tcaataatga tggagtctgt tgttgccctgc aaaaacgggg 480
gccagtgaac attacatcag tgtgtgtgag tcccaggacc ttacaaatat cagtttttgt 540
gttatcagag aaatacgagg gtattgttaa atttgaatcg gatgaattac ctttttggtgt 600
aattggttct aatattggtg atgcacattt tcaagaattc agggctggaa tctcctggaa 660
gectgtggta gatectgatg accccattcc tcagttccct gattgctgca gcagcagcag 720
cagcaggatt ccttcagtga gtgtgctagt tgcagttcct ctggttgtag gccacaaagg 780
gcaggcattt attgaaagga tgctgggggtg cttcaaggaa ttgaagcaag agctgactca 840
ggaagggccg ggcgggggac accccaggte tgcgtggccc ccgcccggcc acgcccagtg 900
gccgcccgag cctgcgagc agggggagga gccgcccga gtggaggcgg aggaggtaga 960
ggaggcggag acggcggaga aggcggagag gaaggtggag gcggaggcga aggtggaggg 1020
gaaggcggag gcggcgggga aggcggaggc ggcggggaag gtggacgcca ccgagaaggt 1080
ggagacggcg gggaaggtgg acgcccgtgg gaaggtggag acggcggagg gtccggggccg 1140

```

ccgggctgag	ctcaagctgg	agcccgaacc	cgagccggtc	cgaggaggcg	agcaggagcc	1200
gaagcaggag	ctggaggatg	agaaccacgc	gaggagcggc	ggtggcgga	acagcgacga	1260
ggttcctccc	cccacccttc	cctccgatcc	accgcggccc	cccgatccct	ctccgcgtcg	1320
cagtcgtgcg	ccgcgcgcgc	gaccccgccc	ccggccccag	accgcgctcc	gtaccccgcc	1380
gcagcctagg	ccccggcccc	cgccccggcc	ccggccccgg	cgcgccctcg	ggggcggtatg	1440
cctggatgtg	gattttgcgg	tggggccacc	aggctgttct	cacgtgaaca	gctttaaggt	1500
gggagagaac	tggaggcagg	aactgcgggt	tatctaccag	tgcttcgtgt	ggtgtggaac	1560
cccagagacc	aggaaaagca	aggcaaagtc	ctgcattctg	catgtgtgtg	gcacccatct	1620
gaacagactc	cactcttgcc	tttctgtgt	cttctttggc	tgcttcacgg	agaaacacat	1680
tcacgagcac	gcagagacga	aacaacacaa	cttagcagta	gacctgtatt	acggagggtat	1740
atactgcttt	atgtgtaagg	actatgtata	tgacaaagac	attgagcaaa	ttgccaaaga	1800
agagcaagga	gaagctttga	aattacaagc	ctccacctca	acagagggtt	ctcaccagca	1860
gtgttcagtg	ccaggccttg	gtgagaaatt	cccaacctgg	gaaacaacca	aaccagaatt	1920
agaactgctg	gggcacaacc	cgaggagaag	aagaatcacc	tccagcttta	cgatcggttt	1980
aagaggactc	atcaatcttg	gcaacacgtg	ctttatgaac	tgcattgtcc	aggccctcac	2040
ccacacgccg	atactgagag	atctctttct	ctctgacagg	caccgatgtg	agatgccgag	2100
tcccagagtg	tgtctggtct	gtgagatgtc	gtcgctgttt	cgggagttgt	attctggaaa	2160
cccgctctct	catgtgccct	ataagttact	gcacctggtg	tggatacatg	cccgccattt	2220
agcagggtac	aggcaacagg	atgcccacga	gttcctcatt	gcagcgtag	atgtcctgca	2280
caggcactgc	aaaggtgatg	atgtcgggaa	ggcgggccaac	aatcccaacc	actgtaactg	2340
catcatagac	caaactctca	caggtggcct	gcagtctgat	gtcacctgtc	aagcctgcca	2400
tggcgctctc	accacgatag	acccatgctg	ggacattagt	ttggacttgc	ctggctcttg	2460
cacctccttc	tggcccatga	gcccaggagg	ggagagcagt	gtgaacgggg	aaagccacat	2520
accaggaatc	accacctca	cggactgctt	gaggagggtt	acgaggccag	agcacttagg	2580
aagcagtgcc	aaaatcaaat	gtggtagtgt	ccaaagctac	caggaatcta	ccaaacagct	2640
cacaatgaat	aaattacctg	tcgttgccct	ttttcatttc	aaacggtttg	aacattcagc	2700
gaaacagagg	cgcaagatca	ctacatacat	ttcctttcct	ctggagctgg	atatgacgcc	2760
gtttatggcc	tcaagtaaag	agagcagaat	gaatggacaa	ttgcagctgc	caaccaatag	2820
tggaaacaac	gaaaataagt	attccttggt	tgtgtgtgtt	aatcaccaag	gaaccttgga	2880
gagtggccac	tataccagct	tcattccggca	ccacaaggac	cagtgggttca	agtgtgatga	2940
tgccgtcatc	actaaggcca	gtattaagga	cgtactggac	agtgaagggt	atttactgtt	3000
ctatcacaaa	caggtgctag	aacatgagtc	agaaaaagtg	aaagaaatga	acacacaagc	3060
ctactga						3067